

Notes on Relativity and Gravitation

J.L.F. BARBÓN

Instituto de Física Teórica IFT UAM/CSIC
C/ Nicolás Cabrera 13
C.U. Cantoblanco, E-28049 Madrid, Spain
`jose.barbon@csic.es`

Abstract

Introductory notes on Relativity (Special and General) for the Course “Gravitation”, delivered at the Master de Física Teórica, IFT UAM/CSIC.

Contents

1	Introduction and Preliminaries	5
1.1	Particles, Fields, Lagrangians and Symmetries	11
1.2	Crisis ‘fin de siècle’	15
1.2.1	Newton	16
1.2.2	Maxwell–Lorentz	18
1.2.3	Einstein	20
2	Relativistic Particles and Fields	25
2.1	Lorentz Covariance	26
2.1.1	The Lorentz Group	26
2.1.2	Tensor Representations	28
2.1.3	Integral Invariants	31
2.1.4	Spinors	33
2.2	Relativistic Particles	34
2.2.1	Free Particles	34
2.2.2	Interacting Particles	36
2.2.3	The Local Energy-Momentum Tensor	39
2.3	Relativistic Fields	42
2.3.1	Scalar Fields	43
2.3.2	Spinor Fields	45
2.3.3	Vector Fields	45
2.3.4	Higher Rank Tensor Fields	50
2.4	Symmetry and the Field	51
2.4.1	Noether	51
2.4.2	Energy-Momentum Tensor in Field Theories	54
3	The Principle of Equivalence	59
3.1	From Galileo to Einstein	60
3.1.1	Relativistic Inertial Fields	62
3.1.2	A Tale of Two Frames: a First Look at General Covariance	65
3.2	Riemannian Geometry and the Principle of Equivalence	72
3.2.1	Particle Probes	72
3.2.2	Gravitational Forces	76
3.3	Minimal Coupling	78
3.4	An Interlude: Vernacular Riemannian Geometry	83

4	Curvature	91
4.1	Relativistic Tidal Acceleration	91
4.1.1	The Riemann Tensor	93
4.1.2	Ricci, Einstein and Weyl Tensors	96
5	Einstein's Equations	99
5.1	The Matter-Gravitation Dynamics	99
5.1.1	Einstein's Equations	100
5.1.2	Energy-Momentum 'Local' Conservation	103
5.1.3	Isometries and Conservation Laws	104
5.2	Finding Solutions	106
5.2.1	More on Isometries	106
5.2.2	Spherically Symmetric Vacuum Solutions	108
5.2.3	The Schwarzschild Solution	110
5.3	The Classic Tests	111
5.3.1	Orbital Precession	111
5.3.2	Gravitational Lensing	114
5.3.3	Gravitational Redshift	117
5.3.4	Echo Delay	118
6	Faraway GR	121
6.1	Weak Gravitational Fields	121
6.1.1	Linearized GR	121
6.1.2	Newtonian and Post-Newtonian	123
6.1.3	Dipole Tails from a Little Rotation	126
6.1.4	Gravitomagnetism	127
6.1.5	Gravitational Energy	130
6.2	Gravitational Radiation	133
6.2.1	Generation	134
6.2.2	Detection	137
7	Cosmological Spacetimes	141
7.1	FLRW spacetimes	141
7.2	The Standard Cosmological Model	142
8	Black Holes	149
8.1	The Schwarzschild Black Hole	149
8.1.1	The Kruskal Extension	152
8.1.2	Global Causal Structure	155
8.2	Gravitational Collapse	158
8.2.1	Imploding Shells	158
8.2.2	Stars and Their Limits	161
8.3	Rotating Black Holes	164
8.3.1	The Kerr Metric	164
8.3.2	The Ergosphere	167
8.3.3	The Penrose Process and the Second Law	169

8.4	Bekenstein and Hawking	171
8.4.1	Black Hole Entropy	171
8.4.2	Hawking Radiation	172

Chapter 1

Introduction and Preliminaries

We assume prior knowledge of mechanics and electrodynamics at the level of the first few chapters of Landau–Lifshitz vols I and II. In particular, we require certain familiarity with Lagrangian mechanics, Maxwell’s equations and Special Relativity (SR). There are many books which can be used as a complement to these notes. A biased list is the following.

- **Elementary level:** Schutz (Cambridge 1980, 2nd ed 2009). Hartle (Benjamin 2003). Coleman (Cambridge 2022). Blundell-Lancaster (Oxford 2025).
- **Comprehensive:** Weinberg (Wiley 1972). Misner-Thorne-Wheeler (Freeman 1973). Landau-Lifshitz (Reverte 1981). Carroll (Benjamin 2003). Zee (Princeton 2013).
- **Advanced level:** Hawking-Ellis (Cambridge 1973). Wald (Chicago 1984). Poisson (Cambridge 2004).
- **Web:** Carroll, gr-qc/9712019. Tong, <https://www.damtp.cam.ac.uk/user/tong/gr.html>

Conventions

The acronym GR is used for General Relativity, SR is used for Special Relativity. Frames adapted to freely falling observers are sometimes termed *frefos*, while generic frames corresponding to fiducial observers are called *fidos*.

Four-dimensional indices transforming in the Lorentz group are labeled with latin letters from the beginning of the alphabet: $a, b, c, \dots$. Four-dimensional indices transforming in the group of diffeomorphisms are labeled with greek letters: $\alpha, \beta, \dots, \mu, \nu, \dots$. Einstein’s summation convention is used in these cases. Three-dimensional *spatial* indices, or more generally indices which do not include time, are labeled with latin letters from the middle of the alphabet: $i, j, k, \dots$, and Einstein’s convention is *not* used. Brackets in indices denote antisymmetrization,

$$A_{[\mu\nu]} = \frac{1}{2}(A_{\mu\nu} - A_{\nu\mu}) , \quad A_{[\mu\nu\rho]} = \frac{1}{3!}(A_{\mu\nu\rho} + A_{\nu\rho\mu} + A_{\rho\mu\nu} - A_{\nu\mu\rho} - A_{\mu\rho\nu} - A_{\rho\nu\mu}) ,$$

and so on. Analogously, curved brackets denote symmetrization, $S_{(\mu\nu)} = \frac{1}{2}(S_{\mu\nu} + S_{\nu\mu})$ and so on.

The metric signature convention is $(-+++)$, so that the proper time element satisfies

$$d\tau^2 = -ds^2 = -g_{\mu\nu} dx^\mu dx^\nu .$$

When ds^2 is positive, ds is interpreted as a proper distance element. When $d\tau^2$ is positive, $d\tau$ is interpreted as a proper time interval. We will often use the notation of a dot over a variable that is being differentiated with respect to τ .

The Levi-Civita symbol ϵ_{abcd} is defined as a completely antisymmetric object in four covariant indices, defining a tensor with respect to proper Lorentz transformations with the convention $\epsilon_{0123} = 1$. This implies that the contravariant version ϵ^{abcd} is completely antisymmetric with $\epsilon^{0123} = -1$. The three-dimensional version appearing in vector calculus is $\epsilon_{ijk} = \epsilon_{0ijk}$ and $\epsilon^{ijk} = -\epsilon^{0ijk}$.

The following set of conventions is adopted when writing the norm of a general four-vector $(U^\mu) = (U^0, \vec{U})$:

$$U^2 = g_{\mu\nu} U^\mu U^\nu = g^{\mu\nu} U_\mu U_\nu = U_\mu U^\mu .$$

For a Lorentz four-vector we have similar expressions in terms of Lorentz indices and the Lorentz metric η_{ab} . In addition, we also have $U^2 = U_a U^a = -(U^0)^2 + \vec{U}^2 = -(U^0)^2 + \sum_{i=1}^3 (U^i)^2$. These conventions are also used for differential operators, such as the Laplacian $\nabla^2 \equiv g^{\mu\nu} \nabla_\mu \nabla_\nu$. Throughout the notes, the covariant derivatives ∇_μ are defined with respect to the Christoffel (also known as Levi-Civita) connection, the unique connection which is symmetric and metric compatible.

Riemann's tensor is defined with the appropriate sign so that the gravitational Lagrangian takes the form

$$\mathcal{L}_g = \frac{1}{16\pi G} \sqrt{-g} R ,$$

where $R = g^{\mu\nu} R_{\mu\nu}$ is the Ricci scalar, $g \equiv \det(g_{\mu\nu})$ and G is Newton's constant.

The integration measure which is invariant under diffeomorphisms is $d^4x \sqrt{-g}$. This generalizes for submanifolds in terms of the induced metric on the submanifold. For any scalar function $f(x)$, we often use the simplified notation

$$\int_X d^4x \sqrt{-g} f(x) = \int_X f .$$

Prelude

The Theory of Relativity, developed mainly by Einstein between 1905 and 1915, is the culmination of more than two millennia of reflections on the nature of space and time. The Einsteinian synthesis mixes space and time into a new entity, called *spacetime*, which is endowed with a dynamical geometry. This geometry is a generalization of both Euclidean and Riemannian geometry, preserving smoothness but allowing curvature. The dynamical content of the geometry of spacetime is identified with the physical phenomenon of gravitation. The whole picture is summarized in the famous *slogan* by J. A. Wheeler: *matter tells space how to curve, space tells matter how to move*.

The ancient Greeks took the first giant step when they discovered the ‘structure of space’, as characterized by the laws of Euclidean geometry. They were so fascinated by this finding that they fell into the Platonic illusion that Euclidean geometry is a logical necessity. This belief started to shatter when XIX century mathematicians discovered logically consistent, yet non-Euclidean geometries, a revolutionary idea which culminated with the Riemannian description of geometries which are smooth, i.e. approximately Euclidean on short distances, but curved on generic distances. By the end of the XIX century the geometry of space was again considered an empirical question, but no suggestion of a dynamical nature was made at this point.




Figure 1.1: The fundamental structure of Einstein’s spacetime is revealed by considering Pythagorean triangles extended in the time direction, representing the time lapses registered by moving clocks (ticking τ units) and static clocks (ticking t units). The purely spatial theorem $\ell^2 = x^2 + y^2$ is generalized to $-\tau^2 = -t^2 + \ell^2$, with time measured in, say, seconds and distances measured in light-seconds.

Greek geometry was extended by Einstein with the invention of the Theory of Special Relativity and its subsequent geometrical interpretation by Minkowski. Roughly, the behavior of light implies a generalization of the symmetries of classical mechanics. This generalization can be stated as a geometrical principle. Given a right triangle in space with sides x, y and ℓ , the Pythagorean theorem states $\ell^2 = x^2 + y^2$. Now, if we ‘lift’ this triangle into the time direction, by specifying two clocks, one at rest ticking to time t and another one in motion ticking to ‘proper time’ τ , one finds, in units where the speed of light is unity,

$$-\tau^2 = \ell^2 - t^2 = x^2 + y^2 - t^2, \quad (1.1)$$

that is to say, the Pythagorean theorem is lifted into *spacetime* by simply sticking a minus sign in

the time directions (see Figure 1.1). This simple geometrical structure is one of the two or three fundamental principles of Physics. Finally, in 1915 Einstein brought together the Riemannian and the Minkowskian generalizations of Euclidean geometry with a new synthesis: spacetime is the curved (Riemannian) generalization of (1.1), with the curvature being determined by the gravitational field.

In these notes we will distill this deep result following a physical path, by showing that this is the only answer to the problem of constructing a relativistic version of the Newtonian law of gravitation, famously stating

$$|F_G| = G \frac{m_1 m_2}{r^2} \quad (1.2)$$

for the gravitational attraction between two masses separated by a distance r . The potential energy at height r in the field of a mass M is given by $-GM/r$ per unit of test mass, up to an additive normalization constant. In orbital motion we have a kinetic energy of the same order of magnitude. The characteristic velocity of an orbit at height r is then $v(r) \sim \sqrt{GM/r}$, so that the relativistic character of this motion is measured by the ratio

$$\frac{v^2}{c^2} \sim \frac{GM}{c^2 r} \sim \frac{r_s}{r},$$

where c is the speed of light and we have defined a characteristic gravitational (so-called Schwarzschild) radius for the gravitational field of a mass M :

$$r_s(M) \equiv \frac{2GM}{c^2}.$$

The gravitational interaction becomes relativistic, i.e. non-Newtonian, for $\phi(r) \equiv r_s/r \sim 1$. The dimensionless potential ϕ also measures the order of magnitude of relativistic effects arising as corrections to the Newtonian theory.

For localized systems, the value of ϕ on Earth is of order $\phi_{\oplus} \sim 10^{-9}$. In the vicinity of the sun, $\phi_{\odot} \sim 10^{-6}$. On the surface of a white dwarf star we have $\phi_{\text{wds}} \sim 10^{-4}$, the same order of magnitude of relativistic effects as the hydrogen atom. Finally, relativistic stars such as neutron stars reach $\phi_{\text{ns}} \sim 0.1$ and black holes always have $\phi_{\text{bh}} \sim 1$.

For non-localized systems, such as a uniform-density distribution, we have $M(r) \sim \rho \cdot r^3$ for the mass enclosed by a sphere of size r . Then, we have $\phi \sim (Hr)^2$, with H defining the Hubble parameter¹

$$H^2(\rho) \equiv \frac{8\pi G\rho}{3c^2}.$$

Therefore, a cosmological model becomes relativistic at distances of order H^{-1} . This happens in our Universe for $H^{-1} \sim \text{Gigaparsecs (Gpc)}$.

In these notes we develop the current theory of the gravitational interaction in relativistic regimes where $\phi = O(1)$, essentially developed by Einstein in the decade prior to 1916. As indicated above, this theory can be tested to order 10^{-6} in solar system experiments, and it is essential to understand the dynamics of relativistic stars and the global properties of the universe on distance scales beyond Gigaparsecs. On the other hand, the effects of gravity are largely irrelevant at the scale of elementary particles. The reason is the extreme weakness of the gravitational interaction between subatomic particles.

¹The precise numerical factors in the definitions of r_s and H are irrelevant for our present discussion of order-of-magnitude estimates, but they conform to the more precise definitions to come.

The ratio of electric to gravitational force between two protons with mass m_p is

$$\frac{|F_{\text{em}}|}{|F_G|} = \frac{e^2}{4\pi G m_p^2} \sim 10^{36},$$

which makes gravitation utterly negligible at the subatomic level.² We can write this ratio in terms of dimensionless quantities characterizing the intrinsic strength of the interaction, the fine structure constant $\alpha = e^2/4\pi\hbar c$, and the gravitational analog $\alpha_G = Gm_p^2/\hbar c$. Since $\alpha \approx 1/137$ is a pure number, it turns out that $\alpha_G \approx 10^{-38}$ is also a pure number. Therefore we find that the quantity

$$m_{\text{Pl}} \equiv \sqrt{\frac{\hbar c}{G}} \approx 10^{19} m_p$$

has dimensions of mass and it is called the Planck mass. The associated Planck energy is $m_{\text{Pl}}c^2 \approx 10^{19}$ GeV and the Planck length $\ell_{\text{Pl}} = \sqrt{\hbar c/G} \approx 10^{-19} \lambda_{\text{proton}} \sim 10^{-33}$ cm, where $\lambda_{\text{proton}} \sim \hbar/m_p c$ is the size of the proton.

The Planck scale has a quantum significance, marking the regime where quantum effects make gravity as strong as the other forces. To see this, notice that a quantum state localized on distance scales ℓ has typical momentum $\hbar/\ell$, according to Heisenberg's principle. When such a momentum is larger than mc for all masses in the problem, the state is highly relativistic, having energy $E \sim \hbar c/\ell$ and effective gravitating mass $\hbar/\ell c$. The dimensionless intensity of the gravitational field for such a mass is

$$\alpha_G(\ell) \sim \frac{G}{\hbar c} \cdot \frac{\hbar^2}{c^2 \ell^2} \sim \frac{G\hbar}{c^3 \ell^2} \sim \left(\frac{\ell_{\text{Pl}}}{\ell}\right)^2,$$

which becomes of order unity when $\ell \sim \ell_{\text{Pl}}$. In fact, it equilibrates the electromagnetic force at scales of the order of tens of Planck lengths.

We conclude that significant quantum gravitational effects would require Planck-mass elementary quantum objects or Planck lengths in the quantum resolution of scales implied by Heisenberg's principle. This means in practice that quantum gravitational effects are irrelevant from the point of view of feasible experiments.

It is possible to imagine situations where this conclusion is significantly modified. For example, it could be that the world is $4 + n$ dimensional below some length scale ℓ_c . In this case Newton's force could actually scale like $1/r^{2+n}$ for $r \ll \ell_c$, which means a stronger growth at short distances. In practice this would imply a lower effective Planck mass and thus room for experimental hope. Currently there is no evidence for such scenarios, but they serve as indications that other possibilities exist.

More generally, the greatest revolutions in theoretical physics are associated to conceptual syntheses of seemingly incompatible theories, in a tight and essentially unique solution. Famous examples of this trend are the theory of quantum fields, the essentially unique unification of Special Relativity and quantum mechanics. Another example is Einstein's theory of gravitation, again the essentially unique theory of relativistic classical gravity. Many physicists hope that the final synthesis involving the three basic constants, $\hbar$, c and G , will be borne out in a conceptually 'tight' fashion, so that we may hope to 'corner' the answer even in the absence of crucial experimental input.

²In fact, on distance scales about a tenth of a millimeter and below, gravitation is already surpassed by Van der Waals forces.

While we wait for the fully fledged development of such a theory, we shall set $\hbar = 0$ for most of these lecture notes.

1.1 Particles, Fields, Lagrangians and Symmetries

Dynamical systems can be described classically by equations of motion or by specifying action functionals. Let the pair $(\mathcal{Q}(t), \mathcal{Q}_t(t))$ denote the *state* of a mechanical system (the data necessary to determine its future, usually initial positions and first time derivatives $\mathcal{Q}_t \equiv d\mathcal{Q}/dt$). Here we regard $\mathcal{Q}$ as an abstract notation for a set of position coordinates, particle label or local excitation of a field.

Given initial and final values of the configuration variables, $\mathcal{Q}(t_i) = \mathcal{Q}_i, \mathcal{Q}(t_f) = \mathcal{Q}_f$, we construct the action functional

$$S[\gamma] = \int_{\gamma} dt L(\mathcal{Q}, \mathcal{Q}_t), \quad (1.3)$$

where γ denotes a given *history*, i.e. the function $\gamma : t \rightarrow \mathcal{Q}(t)$, and L is called the Lagrangian. Then, the equations of motion follow from the extrema of the action functional, $\delta S = 0$, for all variations $\mathcal{Q} \rightarrow \mathcal{Q} + \delta\mathcal{Q}$ with fixed initial and final values. This can be expressed as

$$0 = \delta S = \int_{\gamma} dt \left(\frac{\partial L}{\partial \mathcal{Q}} \delta\mathcal{Q} + \frac{\partial L}{\partial \mathcal{Q}_t} \delta\mathcal{Q}_t \right).$$

Using that $\delta\mathcal{Q}_t = d(\delta\mathcal{Q})/dt$ and integrating by parts we find

$$0 = \int_{\gamma} dt \delta\mathcal{Q} \left(\frac{\partial L}{\partial \mathcal{Q}} - \frac{d}{dt} \frac{\partial L}{\partial \mathcal{Q}_t} \right) + 0, \quad (1.4)$$

where the last vanishing term corresponds to the contribution of total derivatives in time, vanishing because of the boundary conditions $\delta\mathcal{Q}(t_i) = \delta\mathcal{Q}(t_f) = 0$. The extremal trajectory must satisfy (1.4) for all values of $\delta\mathcal{Q}$, implying the vanishing of the term in parenthesis. Therefore we arrive at the so-called Euler–Lagrange equations of motion,

$$\frac{d}{dt} \frac{\partial L}{\partial \mathcal{Q}_t} - \frac{\partial L}{\partial \mathcal{Q}} = 0. \quad (1.5)$$

Lagrangians are defined up to a total derivative in time. Given L , then $L + df/dt$ leads to the same equations of motion.

The elementary example is that of an isolated system of Newtonian particles with generalized coordinates $\mathcal{Q}(t) \rightarrow \vec{X}_p(t)$. The Lagrangian reads

$$L = \sum_p \frac{1}{2} m_p \left(\frac{d\vec{X}_p}{dt} \right)^2 - U(\vec{X}_1, \dots, \vec{X}_p, \dots), \quad (1.6)$$

with U defining a potential energy function. This Lagrangian produces the well-known Newtonian system of equations

$$m_p \frac{d^2 \vec{X}_p}{dt^2} = - \frac{\partial U}{\partial \vec{X}_p}.$$

The Lagrangian formalism is very useful in the description of local degrees of freedom, their interactions and symmetries. It also provides one of the simplest jumping points from classical to quantum physics, since the fundamental quantum object is the probability amplitude for a

transition between initial and final configurations, which has a simple formal relation to the classical action,

$$\mathcal{A}_{\mathcal{Q}_i \rightarrow \mathcal{Q}_f} = \sum_{\text{histories } \gamma_{i \rightarrow f}} e^{iS[\gamma]/\hbar}, \quad (1.7)$$

according to Feynman's formula. In these notes, we will not deal with quantum theory, but we will make full use of the Lagrangian formalism as the most powerful tool in elucidating local dynamics, whose most fundamental idea is the concept of *fields*.

Fields

We may take a formal continuum limit for systems of the form (1.6) with degrees of freedom pinned to points of space, leading to the notion of field theories. Here, the role of the particle index p is played by an ‘approximately’ continuous label corresponding to the point of space $\mathcal{Q}(t) \rightarrow q_{\vec{x}}(t)$. Intuitively, we imagine a field as an assembly of ‘particles’ attached to points of space. The function $q_{\vec{x}}(t)$ denotes ‘oscillations’ of this ‘attached particle’, either in physical space or in some internal space.

Given an equilibrium configuration $q_{\vec{x}}^{(0)}$ of the potential energy function U , we may expand it in powers of the local displacement variable $\phi(\vec{x}, t) = \sqrt{m_{\vec{x}}} (q_{\vec{x}}(t) - q_{\vec{x}}^{(0)})$, which will be referred to as the *field*. In the continuum limit, the assumption of *locality* means that the functional $U[\phi(\vec{x}, t)]$ is the integral of a *density*, admitting an expansion around $\phi = 0$ in powers of derivatives of the field $\phi(\vec{x}, t)$:

$$U[\phi] = \int d^3x \left(V(\phi) + \frac{1}{2} c_s^2 (\vec{\partial}\phi)^2 + \text{higher derivatives} \right),$$

where we have further assumed translational and rotational invariance. Local stability of the equilibrium point requires that the so-called ‘squared speed of sound’, c_s^2 , be positive and the non-derivative term, $V(\phi)$, be well approximated by a positive parabola near $\phi = 0$.

Notice that long distance physics is controlled by the terms with the smallest number of spatial derivatives. Consider for example two contributions with two and four derivatives respectively, $(\vec{\partial}\phi)^2 + \lambda^2 (\vec{\partial}^2\phi)^2$. By dimensional analysis, λ must necessarily be a length scale, so that a field configuration with scale of variation of order ℓ contributes an amount of order $\ell^{-2}(1 + C(\lambda/\ell)^2)$ for some constant C . Hence, at long distances $\ell \gg \lambda$ the higher-derivative terms give a subleading contribution and in any case their effects can be gradually introduced by perturbation theory in the corresponding dimensionless effective couplings, such as $\lambda_{\text{eff}}(\ell) = \lambda/\ell$. This type of ‘long-distance’ expansion is called *Effective Field Theory* and it is the mathematical embodiment of the Principle of Locality.

Collecting all terms together we are led to a field theory Lagrangian

$$L = \int d^3x \mathcal{L}[\phi], \quad (1.8)$$

where the Lagrangian *density* has the following structure

$$\mathcal{L}[\phi] = \frac{1}{2} (\partial_t \phi)^2 - \frac{1}{2} c_s^2 (\vec{\partial}\phi)^2 - V(\phi) + \text{higher derivatives} + \text{boundary terms}. \quad (1.9)$$

In most applications, boundary terms are neglected assuming appropriate boundary conditions for the fields, so that we can integrate by parts at will. The physics of small fluctuations around

equilibrium can be studied by polynomial approximations of the potential function $V(\phi)$. A linear term of the form $J\phi$, called a ‘source term’, is often considered for a general function $J(\vec{x}, t)$ as a way of ‘driving’ the system, to study perturbative response of the equilibrium state. The quadratic term is often called the ‘mass’ of the field. The Euler–Lagrange equations stemming from the long-distance approximation (1.9) are

$$\left(-\partial_t^2 + c_s^2 \vec{\partial}^2\right) \phi = V'(\phi) = J c_s^2 + m^2 c_s^4 \phi + \text{nonlinear terms} , \quad (1.10)$$

where the factors of c_s in the terms proportional to J and m^2 are conventional definitions. The elementary solution of the homogeneous equation for zero mass,

$$(-\partial_t^2 + c_s^2 \vec{\partial}^2) \phi = 0 \quad (1.11)$$

are waves $\phi_w(t, \vec{x})$ propagating at speed c_s . The elementary solution for a static (time-independent) field created by a source J is the solution of Poisson’s equation,

$$\phi(\vec{x})_{\text{static}} = \frac{1}{\vec{\partial}^2} J = - \int d^3 y J(\vec{y}) \int \frac{d^3 k}{(2\pi)^3} \frac{e^{i\vec{k}(\vec{x}-\vec{y})}}{\vec{k}^2} = - \int \frac{d^3 y}{4\pi} \frac{J(\vec{y})}{|\vec{x}-\vec{y}|} . \quad (1.12)$$

Time-dependent disturbances created by a time-dependent source $J(\vec{x}, t)$ are obtained from the static ones by simply recalling that these disturbances travel at speed c_s . Hence the solution is formally the same as (1.12) with the source replaced by its value at the ‘retarded time’ $t - |\vec{x} - \vec{y}|/c_s$,

$$\phi(\vec{x}, t)_{\text{retarded}} = \phi(\vec{x}, t)_{\text{wave}} - \int \frac{d^3 y}{4\pi} \frac{J\left(t - \frac{|\vec{x}-\vec{y}|}{c_s}, \vec{y}\right)}{|\vec{x} - \vec{y}|} , \quad (1.13)$$

where ϕ_{wave} is a general solution of the homogeneous equation (1.11), which might be interpreted as a sort of ‘primordial’ field, coming from the remote past, which was there even before any currents existed that contributed to the field excitations. We can use the same argument to write a formal solution of the general non-linear equation with arbitrary potential:

$$\phi(\vec{x}, t)_{\text{retarded}} = \phi_{\text{wave}} + \frac{1}{-\partial_t^2 + c_s^2 \vec{\partial}^2} \Big|_{\text{ret}} V'(\phi) = \phi_{\text{wave}} - \int \frac{d^3 y}{4\pi c_s^2} \frac{V'\left[\phi\left(t - \frac{|\vec{x}-\vec{y}|}{c_s}, \vec{y}\right)\right]}{|\vec{x} - \vec{y}|} . \quad (1.14)$$

Rather than an explicit solution, this is an integral equation for any potential with quadratic terms or higher non-linearities. This equation may be solved iteratively in the powers of the function $V'[\phi]$. For example, for a mass term we have $V'(\phi) = m^2 c_s^4 \phi$. The term of order m^{2n} in the iterative solution represents a wave sourced by J and averaged over n ‘kicks’ at which the wave is regenerated. This construction is analogous to Huygens’ method of wave dispersion, so that one says that the mass term causes ‘wave dispersion’, as a result of which the effective propagation velocity over distances larger than $1/(mc_s)$ is *smaller* than c_s . Alternatively, we can look for solutions of definite frequency ω and momentum $\vec{k}$,

$$\phi(t, \vec{x})_{\omega, \vec{k}} \sim e^{-i\omega t + i\vec{k}\vec{x}}$$

and the equation of motion implies the so-called ‘dispersion relation’

$$\omega^2 = m^2 c_s^4 + \vec{k}^2 c_s^2 .$$

The basic phenomenon of field dynamics is wave propagation. Locality translates into the finite propagation velocity c_s for any physical perturbations modeled by the field $\phi(t, \vec{x})$.

Symmetries

A classical system is said to enjoy a symmetry action when some group of transformations $\mathcal{G}$ acts on the space of solutions of the equations of motion, i.e. given one solution $\mathcal{Q}$, the transformed $g(\mathcal{Q})$ under $\mathcal{G}$ is also a solution of the equations of motion. One way to ensure this is to demand that the action functional be invariant under the action of $\mathcal{G}$:

$$S[\gamma] = S[g(\gamma)] , \quad (1.15)$$

for any history γ and any group element $g \in \mathcal{G}$. This is slightly more than strictly necessary in the classical realm, because one could do with an action of $\mathcal{G}$ just on the space of extrema of S , rather than the whole domain of definition of S . However, in quantum mechanics one really explores all possible trajectories and symmetries have to respect them all.

Invariance of the action is not always borne out by invariance of the Lagrangian. Because of the ambiguity in Lagrangians by a total derivative, it is enough to demand that a Lagrangian is invariant up to a total time derivative,

$$L[g(\mathcal{Q})] = L[\mathcal{Q}] + \frac{df_g}{dt} . \quad (1.16)$$

For field theories, and under the technical assumption that boundary terms do not contribute to the dynamics, the Lagrangian density may be invariant up to a total derivative in time and space:

$$\mathcal{L}[\phi] \rightarrow \mathcal{L}[g(\phi)] = \mathcal{L}[\phi] + \partial_t f^t + \vec{\partial} \vec{f} , \quad (1.17)$$

for some functions f^t and $\vec{f}$.

The main use of Lagrangian methods is to implement economically the requirements of symmetry. At the same time, this formalism provides a very general link between symmetries and conservation laws, via a famous theorem by Noether.

Let the symmetry $\mathcal{G}$ act on the space of trajectories $\mathcal{Q}(t)$ as a continuous group and work near the identity,

$$g(\mathcal{Q}) = \mathcal{Q} + \delta_g \mathcal{Q} = \mathcal{Q} + \epsilon s[\mathcal{Q}] + O(\epsilon^2) , \quad (1.18)$$

where $s[\mathcal{Q}]$ might be a rather general functional of $\mathcal{Q}$. Since Lagrangians are defined up to a total time derivative, an invariant action is compatible with a variation of the Lagrangian

$$\delta_s L = \epsilon \frac{df_s}{dt} , \quad (1.19)$$

for some f_s . Rewriting this in differential form,

$$\delta_s L = \frac{\partial L}{\partial \mathcal{Q}} \delta_s \mathcal{Q} + \frac{\partial L}{\partial \mathcal{Q}_t} \delta_s \mathcal{Q}_t = \left(\frac{\partial L}{\partial \mathcal{Q}} - \frac{d}{dt} \frac{\partial L}{\partial \mathcal{Q}_t} \right) \delta_s \mathcal{Q} + \frac{d}{dt} \left(\frac{\partial L}{\partial \mathcal{Q}_t} \delta_s \mathcal{Q} \right) . \quad (1.20)$$

Using the equations of motion and equating (1.19) and (1.20), we establish the conservation of the Noether charge Q_s :

$$\frac{dQ_s}{dt} = 0 , \quad Q_s = s[\mathcal{Q}] \frac{\partial L}{\partial \mathcal{Q}_t} - f_s . \quad (1.21)$$

Some examples of importance include the usual definitions of energy and momentum for a free Newtonian particle with Lagrangian $L = \frac{1}{2} m \vec{v}^2$, associated to translation symmetry in time and space, respectively

$$\vec{p} = \frac{\partial L}{\partial \vec{v}} = m \vec{v} , \quad E = \vec{v} \cdot \frac{\partial L}{\partial \vec{v}} - L = \frac{1}{2} m \vec{v}^2 = \frac{\vec{p}^2}{2m} . \quad (1.22)$$

For field theories with Lagrangians of functional form $\mathcal{L}(\partial_t\phi, \vec{\partial}\phi, \phi)$, the statement of a continuous symmetry is that $\delta_s \mathcal{L} = \epsilon \partial_t f_s^t + \epsilon \vec{\partial} \vec{f}_s$ under a field transformation $\delta_s \phi = \epsilon s[\phi]$, where $s[\phi]$ is a functional of $\phi(\vec{x}, t)$ with similar structure as the Lagrangian. Repeating the previous derivation (1.20), neglecting total spatial derivatives, one finds a local ‘continuity equation’

$$\partial_t J_s^t + \vec{\partial} \vec{J}_s = 0 \quad (1.23)$$

for the Noether current defined by

$$J_s^t = s[\phi] \cdot \frac{\partial \mathcal{L}}{\partial(\partial_t \phi)} - f_s^t, \quad \vec{J}_s = s[\phi] \cdot \frac{\partial \mathcal{L}}{\partial(\vec{\partial} \phi)} - \vec{f}_s. \quad (1.24)$$

Whenever we encounter (1.23) we can interpret it as a statement of local conservation of charge, by assigning the role of *charge density* to J^t and the role of *flux-density* to $\vec{J}$. Namely, we define the Noether charge inside a three-dimensional region V^3 as the integral of the density,

$$Q_s(V^3) = \int_{V^3} J_s^t, \quad (1.25)$$

and the flux across the boundary ∂V^3

$$\text{Flux}_s(\partial V^3) = \oint_{\partial V^3} (J_s)_n, \quad (1.26)$$

where $(J_s)_n$ denotes the component of $\vec{J}_s$ along the outward pointing normal to ∂V^3 . Then, upon integration over V^3 , (1.23) is equivalent to

$$\frac{d}{dt} Q_s(V^3) = -\text{Flux}_s(\partial V^3), \quad (1.27)$$

i.e. the variation of charge equals the inward flux through the boundary ∂V^3 , without any creation or destruction of charge inside V^3 .

1.2 Crisis ‘fin de siècle’

Circa 1900 one could summarize the Lagrangian of the ‘fundamental’ physics in terms of two components

$$L_{1900} \sim L_{\text{Newton}} + L_{\text{Maxwell-Lorentz}}, \quad (1.28)$$

where L_{Newton} stands for the model of pointlike masses interacting by instantaneous gravitational forces. The Maxwell–Lorentz sector concerns the electromagnetic phenomena and their coupling to charged massive particles. A mechanical interpretation was missing in this sector, although drawing inspiration from the theory of optics, it was natural to postulate a material medium, called *ether*, that would support the electromagnetic waves. This prompted an active search for mechanical models of the ‘ethereal’ substance. The continued failure of such models led to a conceptual crisis at the turn of the century.

1.2.1 Newton

A model of matter in terms of pointlike objects characterized by positions $\vec{x}_p$ and velocities $\vec{v}_p = d\vec{x}_p/dt$, interacting by a Newtonian gravitational force is described by the Lagrangian

$$L_{\text{Newton}} = \sum_p \frac{1}{2} m_p \vec{v}_p^2 + G \sum_{p < q} \frac{\mu_p \mu_q}{|\vec{x}_p - \vec{x}_q|} . \quad (1.29)$$

The most salient features of this Lagrangian are the instantaneous nature of gravitational forces (the potential energy only depends on positions of particles) and the equality of inertial and gravitational masses, $\mu_p = m_p$, the so-called “Principle of Equivalence”. It also satisfies the Galilean principle of relativity, being invariant under a ten-parameter group of transformations

$$x_i \rightarrow \sum_j R_i^j (x_j + a_j + V_j t) , \quad t \rightarrow t + a_t , \quad (1.30)$$

including boosts by a velocity $\vec{V}$ as well as $\mathbb{R}^3$ -rotations $R_i^j \in SO(3)$ and translations in space and time.³ This formulation of the symmetry principles of Newtonian mechanics assumes the existence of ‘inertial frames’ in which the dynamics actually takes the form expressed in (1.29). The pragmatic rule is that inertial frames are those in which test particles that are isolated (free from any forces) move in straight lines at constant velocity.

The equality of gravitational and inertial masses is famously exemplified by the (perhaps false) story of Galileo throwing objects from the top of Pisa’s tower. In fact, if we denote m the inertial mass and μ the ‘gravitational charge’, it is enough to ensure that the ratio m/μ is a universal constant, equal for all possible bodies, since then this ratio can be set to unity by a rescaling of Newton’s constant.

In the nineteenth century, Eötvös was able to check this universality with great accuracy in a series of experiments comparing the balance of a gravitational force (responding to μ) and a purely inertial force (responding to m). A good example is a body at rest on the surface of the Earth, subject to gravitational and centrifugal forces. Tiny differences of the ratio m/μ between two objects would result in a slight misalignment of the net force acting on them. For the latitude of Budapest, home of Eötvös, one finds

$$\Delta_{m/\mu} \approx 600 |\sin \alpha| , \quad (1.31)$$

where α is the misalignment angle. Eötvös was able to bound $\Delta_{m/\mu}$ to less than one part in 10^8 . The bound was later improved by Dicke in the 60’s to one part in 10^{11} and today we know it to be true to one part in 10^{15} . The conclusion is, therefore, that such a precision test should not be based upon a simple coincidence, but some deep principle. The interpretation of the Principle of Equivalence as the fundamental key to the theory of gravitation is the main contribution of Einstein.

Problem: Eötvös Experiment

Verify expression (1.31) by computing the misalignment of the total gravitational+centrifugal force for two objects of gravitational masses μ_1, μ_2 and inertial masses m_1, m_2 . Give an expression for $|\sin \alpha|$ in

³This is an example of the case that the Lagrangian is invariant up to a total derivative in time.

terms of the surface gravity $g_{\oplus} = GM_{\oplus}/R_{\oplus}^2$, the Earth radius $R_{\oplus}$ and the Earth rotation angular velocity $\Omega_{\oplus}$.

Anticipating its use in the following, we note that Newton's gravitational law can be formally written as a field theory coupled to the mass density of matter. The equations of motion following from (1.29) are

$$m_p \frac{d\vec{v}_p}{dt} = \vec{F}_N = -Gm_p \sum_{q \neq p} \frac{m_q (\vec{x}_p - \vec{x}_q)}{|\vec{x}_p - \vec{x}_q|^3}, \quad (1.32)$$

where we have already implemented the equality of gravitational and inertial masses, so that m_p drops from this equation. The force term on the right hand side can be written in terms of a gravitational potential, $\vec{F}_N = -m_p \vec{\partial} \phi_N$, where

$$\phi_N(t, \vec{x}) = - \sum_q \frac{Gm_q}{|\vec{x} - \vec{x}_q(t)|}. \quad (1.33)$$

This form of the potential, defined up to an additive constant, is the solution of the so-called Poisson equation sourced by the mass density

$$\vec{\partial}^2 \phi_N(\vec{x}, t) = 4\pi G \sum_q m_q \delta^{(3)}(\vec{x} - \vec{x}_q(t)) \equiv 4\pi G \rho_m(\vec{x}, t). \quad (1.34)$$

We see that Newtonian theory can be recast in the form of a field theory with a gravitational potential ϕ_N interacting with mass density through the equations

$$\frac{d\vec{v}_p}{dt} = -\vec{\partial} \phi_N, \quad \vec{\partial}^2 \phi_N = 4\pi G \rho_m, \quad (1.35)$$

which are equivalent to the Lagrangian

$$L_{\text{Newton}} = \sum_p \frac{1}{2} m_p \vec{v}_p^2 - \frac{1}{8\pi G} \int d^3x (\vec{\partial} \phi_N)^2 - \int d^3x \rho_m \phi_N. \quad (1.36)$$

Solving for the potential ϕ_N using Poisson's equation yields precisely (1.33), and substituting back into (1.36) produces the 'action-at-a-distance' form of the Newtonian system (1.29). Notice that the absence of time derivatives from Poisson's equation is directly related to the instantaneous nature of the interactions in (1.29). In principle, a finite speed of propagation for the gravitational interaction can be introduced into this formalism by a simple replacement of Poisson's equation by a more conventional wave equation such as

$$\left(-\frac{1}{c_G^2} \partial_t^2 + \vec{\partial}^2 \right) \phi_N = 4\pi G \rho_m,$$

the standard Newtonian theory thus arising in the limit $c_G \rightarrow \infty$. At any rate, the extreme weakness of the gravitational interaction makes it very difficult to detect dynamical effects of the gravitational field, such as velocity-dependent forces. In comparing with the more familiar optical and magnetic effects, notice that the nominal intensity of gravity is only significant on human scales for planet-sized objects. If c_G is assumed to be in the ballpark of $c \approx 3 \cdot 10^{10}$ cm/s, then any v/c_G effects visible to XIX-century physicists would require 'light-fast' planetary-sized objects hurtling around. This is the kind of situation which finally led to the detection of gravitational waves as late as 2015.

1.2.2 Maxwell–Lorentz

Electrostatic phenomena can be described by a system of equations entirely analogous to (1.35) and (1.36), with the replacement of the Newtonian potential by the Coulomb potential produced by an electric charge e ,

$$\phi_C(\vec{x}, t) = \frac{e}{4\pi|\vec{x}|} .$$

Magnetic phenomena may be incorporated by the introduction of velocity-dependent forces. However, it was found experimentally by Faraday and others that electrodynamics was very efficiently represented by the notion of ‘electromagnetic fields’ interacting with charged matter particles.

Maxwell found the set of local equations that explained all electromagnetic phenomena in terms of electric $\vec{E}$ and magnetic $\vec{B}$ fields in the presence of charges and currents,

$$\begin{aligned} \vec{\partial} \vec{E} &= \rho_e, & c \vec{\partial} \times \vec{B} - \partial_t \vec{E} &= \vec{J}_e , \\ \vec{\partial} \vec{B} &= 0, & c \vec{\partial} \times \vec{E} + \partial_t \vec{B} &= 0 . \end{aligned} \quad (1.37)$$

The charge density and current are defined by

$$\rho_e(\vec{x}, t) = \sum_p e_p \delta^{(3)}(\vec{x} - \vec{x}_p(t)) , \quad \vec{J}_e(\vec{x}, t) = \sum_p e_p \vec{v}_p(t) \delta^{(3)}(\vec{x} - \vec{x}_p(t)) \quad (1.38)$$

and satisfy the conservation law

$$\partial_t \rho_e + \vec{\partial} \vec{J}_e = 0 . \quad (1.39)$$

This conservation law for electric charge acquires a special character in this theory. Since electric charge sources the electric field, one can *detect* the charge inside a large region V^3 by measuring the electric field on the boundary of the region, ∂V^3 . This follows from Gauss’s law, which is the first of the Maxwell equations in (1.37). Integrating this equation over V^3 one finds

$$Q_e(V^3) = \oint_{\partial V^3} E_n , \quad (1.40)$$

where E_n is the normal component of the electric field at ∂V^3 . If $V^3 = \mathbb{R}^3$, this integral computes the total electric charge and the electric flux can be regarded as piercing the so-called ‘celestial sphere at infinity’ S_∞^2 .

Maxwell’s equations depend on a numerical parameter, c , which can be given a physical interpretation by solving the second pair of equations in terms of potentials,

$$\vec{B} = \vec{\partial} \times \vec{A} , \quad \vec{E} = -\frac{1}{c} \partial_t \vec{A} - \vec{\partial} \phi , \quad (1.41)$$

defined up to the *gauge* ambiguity,

$$\phi \rightarrow \phi - \frac{1}{c} \partial_t f , \quad \vec{A} \rightarrow \vec{A} + \vec{\partial} f , \quad (1.42)$$

with $f(t, \vec{x})$ an arbitrary smooth function. Fixing this ambiguity (partially) by imposing the Lorenz condition ⁴

$$\frac{1}{c} \partial_t \phi + \vec{\partial} \vec{A} = 0 , \quad (1.43)$$

⁴Lorenz the *Danish* instead of Lorentz the *Dutch*.

the first pair of Maxwell equations can be rewritten as wave equations

$$\begin{aligned}\left(\vec{\partial}^2 - \frac{1}{c^2} \partial_t^2\right) \phi &= -\rho_e, \\ \left(\vec{\partial}^2 - \frac{1}{c^2} \partial_t^2\right) \vec{A} &= -\frac{\vec{J}_e}{c},\end{aligned}\tag{1.44}$$

which reveals the interpretation of c as the speed of electromagnetic waves. These waves also provide an electromagnetic theory of optics, since oscillating solutions of the electric and magnetic fields exist even outside charged matter and indeed the numerical value of c happens to coincide precisely with the velocity of light *in vacuo*.

The interaction between electromagnetic fields and charged particles is determined by the Lorentz force law

$$m_p \frac{d\vec{v}_p}{dt} = e_p \left(\vec{E} + \frac{\vec{v}_p}{c} \times \vec{B} \right), \tag{1.45}$$

which in turn derives from the Lagrangian

$$\sum_p \frac{1}{2} m_p \vec{v}_p^2 - \sum_p e_p \left(\phi(t, \vec{x}_p) - \frac{\vec{v}_p}{c} \cdot \vec{A}(t, \vec{x}_p) \right). \tag{1.46}$$

The second term may be rewritten in terms of currents and charge densities to obtain the action principle behind Maxwell's equations (1.37),

$$S_{\text{Maxwell-Lorentz}} = \frac{1}{2} \int dt d^3x \left(\vec{E}^2 - \vec{B}^2 \right) - \frac{1}{c} \int dt d^3x \left(c \rho_e \phi - \vec{J}_e \cdot \vec{A} \right), \tag{1.47}$$

with $\vec{E}$ and $\vec{B}$ implicitly written in terms of the potentials $\phi, \vec{A}$ via eq. (1.41). Notice that the potential formalism is very useful in writing the interactions with charges, and essential in the Lagrangian formalism.

The main problem of principle with this theory is the interpretation of the coupling constant c as a velocity of propagation of electromagnetic waves. Immediately the question arises: velocity with respect to what particular frame? Indeed, a Galilean transformation $\vec{x} \rightarrow \vec{x} + \vec{v}t$ cannot possibly be a symmetry of the Maxwell equations, because such a transformation would change the velocity of light to $\vec{c} + \vec{v}$. Hence, if we insist on adhering to a mechanical interpretation of electromagnetism (ether), it seems that Maxwell equations are only valid in the particular frame that sits at rest with respect to the ether.

In this situation, it becomes interesting to measure the velocity of the Earth with respect to the ether. The famous experiments of Michelson–Morley failed to reveal any such drift velocity. At the same time, stellar aberration and other optical data disfavored the possibility that the Earth could be dragging the ether along with its motion. So, there was a real problem that was tackled by Lorentz and others, with a number of *ad hoc* hypotheses.

Despite the fact that Lorentz had developed most of the right formulas of Special Relativity, it is notable that he was unable to interpret them correctly. Poincaré was also close to the solution, as he proposed that the Galilean group of transformations should be replaced by a different group and, in doing so, gave privilege to the electromagnetic part of the total Lagrangian. One would need to replace S_{Newton} with something different that would be compatible with the strange behavior of light. One could say that the Lorentz–Poincaré combination was bound to find the right answer sooner or later... but the first complete solution was given by Einstein in his famous 1905 work.

1.2.3 Einstein

Einstein's starting point was to take for granted the Galilean principle of relativity, i.e. that motion at constant velocity could not have absolute meaning, and at the same time assume as a feature of nature that the velocity of light is a constant, independent of the frame of reference.

Heuristically, one could argue that one simply needs the existence of a maximum velocity of propagation of signals. If one assumes the existence of this limiting velocity, the principle of relativity requires it to be the same in all frames in relative uniform motion, because otherwise one could assign an intrinsic velocity to a frame of reference by quoting the maximal velocity of signals in that frame. From this point of view, one could have Special Relativity even if light was not propagating right at the limiting velocity. For simplicity, let us assume in any case that c , the light's velocity *in vacuo*, is the maximal one. Then, the principle of relativity says that there must exist a group of transformations that leaves the physics invariant and that preserves the condition of uniform rectilinear motion. Using orthogonal Cartesian coordinates, any such transformation,

$$(t, \vec{x}) \rightarrow (t', \vec{x}') = \tilde{L}(t, \vec{x}) \quad (1.48)$$

must send straight trajectories $\vec{x}(t)$ into straight trajectories $\vec{x}'(t')$, and so it must be a *linear* map. Furthermore, the invariance of light's velocity means that

$$\vec{x}^2 - c^2 t^2 = 0 = \vec{x}'^2 - c^2 t'^2 . \quad (1.49)$$

This equation, together with the linearity property, implies that $\vec{x}^2 - c^2 t^2$ is invariant up to a constant factor,

$$\vec{x}^2 - c^2 t^2 \rightarrow f(\vec{V}) (\vec{x}^2 - c^2 t^2) , \quad (1.50)$$

where $f(\vec{V})$ can only depend on the relative velocity of the frames $\vec{V}$. Spatial isotropy further imposes the condition that f must be a function of the modulus $|\vec{V}| = V$. Introducing now two boosts with velocities $\vec{V}_1$ and $\vec{V}_2$, and relative velocity $\vec{V}_{12}$, we find, upon performing two iterated transformations,

$$f(V_2) = f(V_1)f(V_{12}) , \quad \text{equivalently} \quad f(V_{12}) = \frac{f(V_2)}{f(V_1)} , \quad (1.51)$$

but V_{12} depends on the relative angle between $\vec{V}_1$ and $\vec{V}_2$, unlike the right hand side of the previous equation. Hence, we conclude that f does not depend on $\vec{V}$ and in fact $f = 1$. We thus define the Lorentz group as those linear transformations that leave the quadratic form

$$c^2 t^2 - \vec{x}^2 = \text{invariant} . \quad (1.52)$$

This group is called $O(1,3)$. When supplemented by the spacetime translations it becomes the Poincaré group, a ten-parameter group which contracts to the Galilean group in the limit $V/c \rightarrow 0$. Factoring out the subgroup of $SO(3)$ rotations, we can align the boost velocity in a given direction, say x , and we are left with a two-dimensional problem of linear transformations leaving $c^2 t^2 - x^2 = (ct - x)(ct + x)$ invariant. From this point on, we choose units so that $c = 1$. The required linear transformations (maintaining the origin fixed) are

$$x \pm t \rightarrow e^{\pm\psi} (x \pm t) ,$$

or, in matrix notation

$$\begin{pmatrix} t' \\ x' \end{pmatrix} = \begin{pmatrix} \cosh \psi & \sinh \psi \\ \sinh \psi & \cosh \psi \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix}. \quad (1.53)$$

The parameter ψ is called *rapidity*. It can be related to the relative velocity by monitoring the transformation of the point $x = 0$, leading to $\tanh \psi = x'/t' = V$. Hence, we have the usual, so-called *boost* transformations⁵

$$\begin{pmatrix} t' \\ x' \end{pmatrix} = \begin{pmatrix} \gamma & \gamma V \\ \gamma V & \gamma \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix}, \quad (1.54)$$

with $\gamma = (1 - V^2)^{-1/2}$. We see that, as expected, we obtain the Galilean transformations $t' \approx t$ and $\vec{x}' \approx \vec{x} + \vec{V}t$ whenever the relative frame velocities are small compared to the velocity of light, $|\vec{V}| \ll 1$.

Problem: Ships on Top of Ships

Using the rapidity parametrization, find a relativistic version of the formula for the addition of velocities when two Lorentz transformations are iterated. Imagine you construct a huge spaceship capable of maximal velocity $V_M < 1$. If one builds another spaceship with identical capabilities, to be launched from the first one in the forward direction, what is the net speed that the second ship can reach? If this procedure is iterated, what is the maximum velocity attainable in the limit of an infinite series of ships launched on top of ships?

Minkowskian Geometry

The geometrical interpretation of Special Relativity was given by Minkowski in 1908. We can think of $\mathbb{R}^{1+3} \equiv \mathbb{R} \times \mathbb{R}^3$ as the spacetime, where the first factor is ‘time’ and the second factor is ‘space’. We have the set of ‘events’ $(t, \vec{x}) \in \mathbb{R}^{1+3}$, equipped with a metric

$$\mathbf{g}(\Delta t, \Delta \vec{x}) = \Delta \vec{x}^2 - \Delta t^2 \quad (1.55)$$

between pairs of events.⁶ This generalizes the standard Euclidean metric of $\mathbb{R}^3$, which computes the rotationally invariant length-squared of the vector $\Delta \vec{x}$, by the addition of the term $-\Delta t^2$ which makes the Minkowski metric indefinite (not positive definite). Lorentz transformations introduce motions on $\mathbb{R}^{1+3}$ that preserve this metric, inducing a causal structure. To explain the physical meaning of the metric, we notice that any point in $\mathbb{R}^{1+3}$ must fall into one of three Lorentz-invariant sets, depending on the sign of $\mathbf{g}$.

If $\mathbf{g} < 0$, there exists a frame in which $\Delta \vec{x}^2 = 0$ and the events ‘occur’ at the same point in space. In this case we say that the interval between such events is ‘timelike’ because in this rest frame, the interval can be written as $\mathbf{g} = -\Delta \tau^2$, where τ is the time as measured by a clock in that rest frame, called ‘proper time’.

⁵The boost with rapidity ψ is the analytic continuation of a standard Euclidean rotation by angle $\theta = i\psi$, leaving the Euclidean interval $I_E = (it)^2 + x^2$ invariant.

⁶A common notation for infinitesimal intervals is $\mathbf{g}(dt, d\vec{x}) \equiv ds^2 = -dt^2 + d\vec{x}^2$.




Figure 1.2: The orbits of the Lorentz group in the (t, x) plane form a set of hyperbolae, defined by $\mathbf{g} = -t^2 + x^2 = \text{constant}$. Orbits in the spacelike region $\mathbf{g} > 0$ intersect the $t = 0$ axis, whereas orbits in the timelike region, $\mathbf{g} < 0$ intersect the $x = 0$ axis and preserve the sign of the times. The hyperbolae degenerate to straight lines in the light-like region, $\mathbf{g} = 0$ given by all the points connected by light rays from the origin. Rotating this picture into the rest of $\mathbb{R}^3$ gives the past and future light-cones for the $\mathbf{g} = 0$ ‘null’ set.

On the other hand, for $\mathbf{g} > 0$ there is a frame where $\Delta t^2 = 0$, and the events can be considered as ‘simultaneous’. In this situation we speak of ‘spacelike’ separated events. In the special frame of simultaneity the metric equals the standard Euclidean metric $\mathbf{g} = \Delta \vec{x}^2$.

Finally, the events with $\mathbf{g} = 0$ are said to be ‘lightlike’ or ‘null’ separated and correspond to those that can be connected by light rays. Generally speaking, the physical content of the metric is that $\sqrt{\pm \mathbf{g}}$ can be interpreted as either a standard time lapse or a standard spatial distance.

It turns out that the sign of Δt is Lorentz invariant for timelike intervals, making it possible to establish causal relationships between such events. In particular, the velocity of any inertial motion between timelike events, $\vec{v} = \Delta \vec{x} / \Delta t$, is subluminal, $|\vec{v}| < 1$. On the other hand, the ordering of times for spacelike events is *not* Lorentz invariant, so that in that case we cannot agree on a Lorentz-invariant notion of causal influence. Effective velocities in that case would be superluminal. Hence, we conclude that only sub-luminal velocities can correspond to causal processes. Given a point P in spacetime, the set of events that can be reached by a causal path (one whose velocity never exceeds that of light) lie on the future light-cone or in its interior. At the same time, all possible causal paths that can reach P from its past lie either on the past light cone or in its interior.

Minkowskian geometry is a generalization of Euclidean geometry which expresses the fact that the rotations of $\mathbb{R}^3$ generalize into ‘rotations’ of $\mathbb{R}^{1+3}$ in such a way that physics is independent of ‘direction’. In the purely spatial directions we have the usual property of *isotropy* saying that no particular axis or orientation angle in $\mathbb{R}^3$ is special. In the spacetime directions mixing space and time by the boost velocity $\vec{V}$, the symmetry means that physics is independent of $\vec{V}$.

Problem: Twin Paradox

Given two timelike separated events, A and B, prove that any traveler making the journey from A to B will maximize her/his proper travel time by moving at constant velocity (i.e. along the straight worldline joining A and B). Show that the subjective travel time can be made arbitrarily short if we tolerate arbitrary accelerations (Langevin's twin paradox).

Free Particle Dynamics

With the simple elements introduced so far we are ready to study the relativistic dynamics of free particles. Let us consider a causal path γ of a particle of mass m , specified as a function $\vec{x}(t)$ in such a way that the velocity $\vec{v} = d\vec{x}/dt$ is sub-luminal at all times, $|\vec{v}| < 1$. Following the general rules of Lagrangian dynamics, we define an action

$$S_P = \int dt L_P(\vec{x}(t), \vec{v}(t))$$

with the sole requirement that it gives the correct equations of motion for a free particle, i.e. $\vec{v} = \text{constant}$, and that it be Lorentz invariant. To find the right action, let us see the path as the limit of fine polygonal approximations. Then we can write the action as

$$S_P \approx \sum_{\text{segments}} \Delta S_{\text{segment}}$$

On each straight segment, we can pick a frame with respect to which the particle is at rest. The clock on that comoving frame ticks to 'proper time' τ . Since the state of the particle in this frame is that of 'rest', we can apply the standard policy from elementary mechanics and, defining the Lagrangian as 'kinetic energy minus potential energy', we can write

$$\Delta S_{\text{segment}} = -C \Delta \tau_{\text{segment}} , \quad (1.56)$$

where C is a constant specifying the 'potential energy' of the particle at rest. Taking now the limit we find the action for a free particle to be

$$S_P = -C \int_{\gamma} d\tau . \quad (1.57)$$

From the definition of proper time we have

$$d\tau^2 = dt^2 - d\vec{x}^2 = dt^2 \left(1 - \left(\frac{d\vec{x}}{dt} \right)^2 \right) ,$$

where $(t, \vec{x})$ are coordinates in an external, fixed, but arbitrary Lorentz frame. We thus rewrite (1.57) in an arbitrary Lorentz frame in the form

$$S_P = -C \int_{\gamma} dt \sqrt{1 - \vec{v}^2} , \quad (1.58)$$

and the value of the constant C can be obtained by expanding the Lagrangian at low velocities and matching to the Newtonian form: $-C\sqrt{1-\vec{v}^2} \approx -C + \frac{1}{2}C\vec{v}^2 \approx -C + \frac{1}{2}m\vec{v}^2$, which determines $C = m$ and the final form

$$S_P = -m \int_{\gamma} d\tau = -m \int_{\gamma} dt \sqrt{1-\vec{v}^2} , \quad (1.59)$$

of the free particle action in Special Relativity. From here, we get the free equation of motion, $d\vec{v}/dt = 0$ and the relativistic versions of the conserved quantities in the motion, the momentum and the energy

$$\vec{p} = \frac{\partial L_P}{\partial \vec{v}} = \frac{m \vec{v}}{\sqrt{1-\vec{v}^2}} , \quad E = \vec{v} \cdot \vec{p} - L_P = \frac{m}{\sqrt{1-\vec{v}^2}} = \sqrt{\vec{p}^2 + m^2} . \quad (1.60)$$

In particular, a particle has divergent energy as it approaches the speed of light, giving another, more dynamical incarnation of the limiting character of the speed of light.

Notice that Lorentz invariance of the dynamics leads naturally to the notion of a potential energy at zero velocity, given by the rest mass, or the famous $E = mc^2$ if the conventional units were restored. Writing the last of (1.60) equations in the form $E^2 - \vec{p}^2 = m^2$, we notice that the four-vector $(E, \vec{p})$ behaves under Lorentz transformations just as the coordinates $(t, \vec{x})$.

Chapter 2

Relativistic Particles and Fields

Spacetime as described in the previous chapter, particularly in the geometrical picture of Minkowski, lies at the very foundation of contemporary physics. The only other pillar of comparable importance is Quantum Mechanics. Both foundational structures were laid down in the first quarter of the XX century.

As regards the spacetime pillar, the basic concepts are the symmetry principles on one side and the dynamical Principle of Locality on the other side. The first one is of a more kinematical nature: one assumes homogeneity of space and time and the relativity principle embodied mathematically in the Lorentz group. Geometrically this is couched by saying that the metric of spacetime, as well as the action principle determining the dynamics, is invariant under the Poincaré group, a semi-direct product of the Lorentz group and the translations in space and time. Anticipating what is to come, the fact that gravitation will generalize the geometry of spacetime *does not* eliminate the importance of this principle, since it will be still true at sufficiently short distances, being part of the basic assumption of *smoothness* of spacetime.

The Principle of Locality is incorporated by the requirement that no action-at-a-distance exists, at least on scales accessible to experiment. In a relativistic theory this is equivalent to having a maximal speed of propagation of signals. Locality takes slightly different forms depending on the model of matter and forces that we adopt. For particles, we specify that their interactions occur at points in spacetime, either with other particles or with fields. Fields are the very embodiment of the notion of *local* degrees of freedom, since they are a mathematical abstraction from ‘particles attached to points in space’. In their case, the Principle of Locality means that interactions of fields are specified at infinitesimally close points of spacetime or, in other words, the Lagrangian density is specified as a function at each point in spacetime, containing at most a finite number of derivatives of the fields. The expansion in derivatives is actually a systematic expansion, called *Effective Field Theory*, in which short-distance physics gives power-like corrections to long-distance physics. This notion of Effective Field Theory extends perfectly well into the quantum domain (Quantum Field Theory) and remains one of the backbones of our current description of nature in the XXI century.

In this chapter we will construct the actions specifying the dynamics in accordance with these principles, for both particles and fields. The particle-field dichotomy is crucial in classical physics. It was certainly a requirement of experiment at the end of the XIX century. On the other hand, the existence of *both* particles and fields became a source of notorious mathematical troubles, caused by the fact that pointlike sources necessarily introduce singularities in field configurations. The classic example was the divergent electromagnetic self-energy of a particle

due to the singularity of the Coulomb field. Despite attempts at unification in the framework of non-linear classical field theories, this basic dichotomy was not transcended until the arrival of Quantum Field Theory, and in some ways this conundrum still plagues the foundations of physics today.¹ These developments are, however, beyond the scope of these notes. Here we will introduce the Lorentz-covariant formalism to deal with particles interacting through the simplest field theories, guided by pedagogical simplicity and relevance to the actual description of nature.

2.1 Lorentz Covariance

The procedure outlined at the end of the previous chapter for free particles can be repeated for more complicated systems. In postulating dynamical principles based on Lorentz symmetry, we must always start by guessing a Lorentz-invariant action. Further constraints are brought by the principles of locality and the matching to known limits, such as the Newtonian approximation.

2.1.1 The Lorentz Group

In order to proceed with this program, it is useful to introduce some powerful notation to help in the search of Lorentz-invariant combinations of physical quantities.

The Lorentz group $O(1, 3)$ was defined as the set of linear maps of $\mathbb{R}^{1+3}$ onto itself with the condition of keeping the Minkowski interval invariant. Working in differential notation

$$ds^2 = -d\tau^2 = -dt^2 + d\vec{x}^2 \quad (2.1)$$

is left invariant. In the notation $(t, \vec{x}) = (x^0, \vec{x}) = (x^a)$, with $a = 0, 1, 2, 3$, we have

$$x^a \rightarrow x'^a = \sum_b L^a{}_b x^b \equiv L^a{}_b x^b, \quad (2.2)$$

where we have used Einstein's convention of implicit summation of repeated indices in top-down pairs.² Then, the Minkowski metric η_{ab} ,

$$ds^2 = \eta_{ab} dx^a dx^b, \quad (\eta_{ab}) = \text{diag}(-1, 1, 1, 1) \quad (2.3)$$

is left invariant by the Lorentz matrices,

$$\eta_{ab} L^a{}_c L^b{}_d = \eta_{cd} \quad (2.4)$$

or, in matrix notation $\eta = L^t \eta L$, where L^t stands for the transpose of the L matrix. Inverting this relation and noting that $\eta = \eta^{-1}$ we have $\eta = L^{-1} \eta (L^{-1})^t$ or, in index notation,

$$\eta_{ab} (L^{-1})^a{}_c (L^{-1})^b{}_d = \eta_{cd}. \quad (2.5)$$

It is common practice to write

$$(L^{-1})^a{}_b = L_b{}^a,$$

¹Under the modern name of *Hierarchy Problem*.

²In the following, we will use Einstein's summation convention for any set of four-dimensional indices, keeping the explicit sum notation for vector-like three-dimensional indices.

so that a switch in the ordering of upper and lower indices is tantamount to inversion. This is the Lorentzian version of the matrix equation $L^t = L^{-1}$, which is familiar for standard orthogonal matrices.

The transformations $x \rightarrow x' = Lx$ can be viewed as a change of coordinates, in what is called the *passive* view. There is also an *active* view, in which we interpret $x \rightarrow x'$ as a motion in the spacetime, a map between different points of Minkowski spacetime, expressed in the *same* coordinate system. In this active view, the invariance of the metric (2.5) expresses the fact that any geometrical statement based on lengths or angles is actually invariant under this motion. This is usually called an *isometry* in the mathematical literature.

The full group of isometries of Minkowski spacetime is known as the Poincaré group, a semi-direct product of the Lorentz group and the spacetime translations. More explicitly,

$$x^b \longrightarrow x'^b = a^b + L^b_c x^c, \quad (2.6)$$

a ten-parameter group of transformations which becomes the Galilean group at velocities small compared to the speed of light.

Setting $c = d = 0$ in (2.4) we have

$$-1 = \eta_{00} = L^a_0 L^b_0 \eta_{ab} = -(L^0_0)^2 + \sum_i (L^i_0)^2, \quad (2.7)$$

so that $(L^0_0)^2 \geq 1$ and one cannot change continuously the sign of L^0_0 . Similarly, from $\det \eta = \det \eta (\det L)^2$ we find that $\det L = \pm 1$ and we cannot smoothly change the sign of the determinant either. This divides the set of L matrices into four connected components according to the signs of L^0_0 and $\det(L)$. The subgroup of Lorentz transformations with both L^0_0 and $\det(L)$ positive are continuously connected to the identity and denoted $SO_+(1,3)$, and we will sometimes refer to them as *small* Lorentz transformations. We can further characterize the small transformations by expanding near the identity. Let $L^a_b = \delta^a_b + \theta^a_b + \dots$ with $|\theta^a_b| \ll 1$. Then, the isometry condition (2.5) is equivalent to requiring that the matrix $\eta_{ab} \theta^b_c$ be antisymmetric in the indices (a, c) .

We can reach the other components of $O(1,3)$ by the action of $\mathcal{P}$ and $\mathcal{T}$ on the small transformations, where $\mathcal{P}$ stands for parity, $(t, \vec{x}) \rightarrow (t, -\vec{x})$, and $\mathcal{T}$ is time reversal, $(t, \vec{x}) \rightarrow (-t, \vec{x})$. In what follows, we will tacitly refer to *the* Lorentz transformations as the *small* ones, and state it explicitly whenever we are considering parity and time-reversal transformations. A good reason to restrict the Lorentz group to the small component is that, actually, neither $\mathcal{P}$ nor $\mathcal{T}$ are fundamental symmetries of nature, being violated in the quantum theory of elementary particles.

Upon analytic continuation $(t, x) \rightarrow (it, x)$ the Lorentzian metric becomes Euclidean $ds^2 = d(it)^2 + d\vec{x}^2$. The resulting connected group is $SO(4)$, which is locally isomorphic to $SU(2)_L \times SU(2)_R$. The operation of analytic continuation, being continuous, should not change the discrete structure of group representations. This means that all representations of the Lorentz group can be constructed as the pairs (j_L, j_R) , with $j_{L,R}$ the standard spins of $SU(2)$ irreducible representations. This form is very useful to describe spinor degrees of freedom, although most of the applications in classical physics can be exhausted by the so-called *tensor* representations, which appear in the tensor products of vector representations.

We thus conclude that a relativistic model is given by a Poincaré-invariant Lagrangian defined over Minkowski spacetime, i.e. the pair $(\mathbb{R}^{1+3}, \eta)$, with dynamical variables in some representations of the small Lorentz group $SO_+(1,3)$.

2.1.2 Tensor Representations

The defining (vector) representation of the Lorentz group corresponds to any quantity that transforms as the coordinates:

$$U^a \rightarrow L^a{}_b U^b. \quad (2.8)$$

Any such quantity is called a ‘contravariant’ vector. Given two contravariant vectors, U, V , we can form a two-index object $T^{ab} = U^a V^b$ by ‘tensor’ product. It transforms as

$$T^{ab} \rightarrow L^a{}_c L^b{}_d T^{cd}.$$

Other combinations with the same transformation law are linear combinations of tensor products, $U^a V^b + X^a Y^b + \dots$. Hence, any set of quantities transforming as above may be defined as a second-rank contravariant tensor, and the generalization to arbitrary rank is obvious.

A different transformation law is followed by the differential operators

$$\partial_a \equiv \frac{\partial}{\partial x^a}.$$

If we view x'^a as functions of x^a , we have

$$\frac{\partial x'^a}{\partial x^b} = L^a{}_b, \quad \text{and the inverse} \quad \frac{\partial x^a}{\partial x'^b} = (L^{-1})^a{}_b. \quad (2.9)$$

We can now see why it was a good idea to introduce the notation

$$L_b{}^a \equiv (L^{-1})^a{}_b. \quad (2.10)$$

This notational trick dovetails nicely with the summation convention since, using the chain rule we may write

$$\partial'_a = \frac{\partial x^b}{\partial x'^a} \partial_b = L_a{}^b \partial_b. \quad (2.11)$$

Such a transformation law, with the inverse Lorentz matrix L^{-1} , is conventionally called ‘covariant’. General covariant tensors can be defined following the previous steps of their contravariant cousins, by tensor product and addition. For example, a rank-two covariant tensor transforms as

$$A_{ab} \rightarrow L_a{}^c L_b{}^d A_{cd} \quad (2.12)$$

and so on. Mixed tensors with p contravariant and q covariant indices can be constructed in straightforward fashion. They can be given an intrinsic definition by regarding the collections of indexed numbers as components in an abstract basis formed by the contravariant differentials dx^a and the covariant derivative operators ∂_b , i.e. we require the formal expression

$$\mathbf{T} = \sum_{a_1, \dots, b_1, \dots} T^{a_1 \dots a_p}{}_{b_1 \dots b_q}(x) dx^{b_1} \otimes \dots \otimes dx^{b_q} \otimes \partial_{a_1} \otimes \dots \otimes \partial_{a_p} \quad (2.13)$$

to be invariant under a change of basis

$$x^a \rightarrow x'^a = L^a{}_b x^b, \quad dx^a \rightarrow L^a{}_b dx^b, \quad \partial_a \rightarrow L_a{}^b \partial_b. \quad (2.14)$$

The transformation laws of the components follow suit:

$$T^{a_1 \dots}{}_{b_1 \dots}(x) \longrightarrow T'^{a_1 \dots}{}_{b_1 \dots}(x') = L^{a_1}{}_{c_1} \dots L_{b_1}{}^{d_1} \dots T^{c_1 \dots}{}_{d_1 \dots}(x), \quad (2.15)$$

where we have written the functional dependence on the coordinates to expose how the functions transform, in the case the tensor quantities are defined locally point by point. For instance, a scalar function with no indices transforms as $\phi'(x') = \phi(x)$. Therefore, the value of the function is invariant but the function itself does transform.

This construction also shows that any pair of indices that are identified and summed up with Einstein's convention become irrelevant as far as the transformation rules are concerned. That is, the 'contracted' product $U^a V_a$ of a contravariant and a covariant vector is a Lorentz invariant, whereas the contraction $V_a T^{ab}$ transforms as a contravariant vector, according to the free index b . We can also construct invariant differential operators, such as

$$\partial^2 \equiv \partial_a \partial^a = \eta^{ab} \partial_a \partial_b = -\partial_t^2 + \vec{\partial}^2, \quad (2.16)$$

usually called Laplacian or d'Alembertian.

Invariant Tensors

The task of constructing Lorentz invariants is greatly aided by the definition of *invariant* tensors. These are tensors that retain their value under Lorentz transformations. The most obvious case is the Kronecker delta δ_b^a with upper-and-lower indices. It transforms as an invariant mixed tensor of rank two:

$$\delta_b^a = L^a{}_c L_b{}^d \delta_d^c. \quad (2.17)$$

Another important invariant tensor is the metric itself. According to (2.5) we have

$$\eta_{ab} L_c{}^a L_d{}^b = \eta_{cd}, \quad (2.18)$$

as corresponds to the transformation law of a rank-two covariant tensor. It is convenient to define the inverse metric (equal to itself) with upper indices

$$(\eta^{ab}) = \text{diag}(-1, 1, 1, 1). \quad (2.19)$$

Hence, $\eta^{ab} \eta_{bc} = \delta_c^a$, and given the tensor character of η_{ab} and δ_b^a , it follows that η^{ab} is an invariant contravariant tensor of rank two.

The invariant tensors constructed from the metric are especially interesting because they can be used to make covariant tensors out of contravariant tensors and vice versa. By the rules already stated it is very easy to check that, given a contravariant tensor, the contraction

$$\eta_{ab} T^{bc\dots} = \bar{T}_a{}^{c\dots} \quad (2.20)$$

transforms as a mixed tensor with the indices as indicated. We say that the metric can be used to 'lower' an index. It is also common to simplify the notation and use the same letter to denote the new tensor, relying on the position of the indices to distinguish it from the previous fully contravariant one, i.e. we usually write $\bar{T}_a{}^{c\dots} = T_a{}^{c\dots}$. Analogously, we can use the upper-index version to 'raise' covariant indices into contravariant ones. When playing the game of raising and lowering indices, the reader should be aware of the relative positions of the indices, for instance $T_a{}^b = \eta_{ac} T^{cb} \neq T^a{}_b = \eta_{bc} T^{ac}$ unless T^{ab} happens to be symmetric.

A final interesting object is the Levi-Civita tensor. Let us define the *permutation symbol* as the completely antisymmetric object $[abcd]$ with $[0123] = 1$. In other words, it is given by the even/odd parity of the permutation $\pi : (0, 1, 2, 3) \rightarrow (a = \pi(0), b = \pi(1), c = \pi(2), d = \pi(3))$:

$$[0123] = 1, \quad [\pi(0)\pi(1)\pi(2)\pi(3)] = (-1)^{\text{parity}(\pi)}. \quad (2.21)$$

The Levi–Civita tensor is defined as a covariant object with the same numerical value as the permutation symbol ³

$$\epsilon_{abcd} = [abcd] . \quad (2.22)$$

The identity

$$\epsilon_{abcd} L^a{}_e L^b{}_f L^c{}_g L^d{}_h = \det(L) \epsilon_{efgh} \quad (2.23)$$

implies that the Levi–Civita symbol is a true invariant tensor for small Lorentz transformations (more generally, all $SO(1, 3)$ Lorentz transformations with $\det(L) = 1$), and flips its sign under parity or time reversal. The familiar Levi–Civita 3-tensor of vector algebra is

$$\epsilon_{ijk} = \epsilon_{0ijk} .$$

We can also define associated objects by raising/lowering indices with the metric η_{ab} . In particular, the completely contravariant version ϵ^{abcd} is also completely antisymmetric with

$$\epsilon^{0123} = \eta^{0a} \eta^{1b} \eta^{2c} \eta^{3d} \epsilon_{abcd} = \det^{-1}(\eta) \epsilon_{0123} = -1 ,$$

so that it differs from the permutation symbol by a crucial minus sign coming from the Lorentzian signature of spacetime:

$$\epsilon^{abcd} = -[abcd] . \quad (2.24)$$

A useful formula for contractions of arbitrary d -dimensional Levi–Civita tensors is

$$\epsilon^{c_1 \dots c_n a_{n+1} \dots a_d} \epsilon_{c_1 \dots c_n b_{n+1} \dots b_d} = (-1)^{\text{sig}} n! \det(\delta^{a_j}{}_{b_k}) = (-1)^{\text{sig}} n!(d-n)! \delta^{a_{n+1}}_{b_{n+1}} \dots \delta^{a_d}_{b_d} , \quad (2.25)$$

where $(-1)^{\text{sig}}$ is the signature, +1 for Euclidean metrics and -1 for Lorentzian ones, and the square brackets stand for the complete anti-symmetrization of indices. These give the Lorentzian generalization of the useful identities of vector calculus

$$\sum_k \epsilon_{ijk} \epsilon_{mnk} = \delta_{im} \delta_{jn} - \delta_{in} \delta_{jm} , \quad \sum_{jk} \epsilon_{ijk} \epsilon_{mjk} = 2\delta_{im} , \quad \sum_{ijk} \epsilon_{ijk} \epsilon_{ijk} = 6 . \quad (2.26)$$

Having one more invariant tensor at our disposal we can use it to construct new kinds of Lorentz invariants. Here it is convenient to work in arbitrary dimensions to see the emergence of combinatorial patterns.

Given a rank- p antisymmetric tensor $A_{[p]}$ with components $A_{a_1 \dots a_p} = A_{[a_1 \dots a_p]}$ (also known as a p -form) we can construct a so-called Poincaré dual tensor $\tilde{A}^{a_{p+1} \dots a_d}$ which is completely antisymmetric in $d - p$ indices and has components

$$\tilde{A}^{a_{p+1} \dots a_d} = \frac{1}{p!} \epsilon^{a_1 \dots a_p a_{p+1} \dots a_d} A_{a_1 \dots a_p} . \quad (2.27)$$

It contains the same information as the original tensor (this is similar in spirit to the raising and lowering of indices). Poincaré duality is involutive up to a sign:

$$\tilde{\tilde{A}}_{[p]} = (-1)^{\text{sig}} (-1)^{p(d-p)} A_{[p]} . \quad (2.28)$$

³The reader should be aware of the fact that sign conventions for Levi–Civita tensors differ significantly across the literature.

If $p = d/2$ we can now write an invariant contraction between A and $\tilde{A}$. More generally, given a p -form A and a different $(d - p)$ -form A' we can construct the Lorentz invariant

$$\frac{1}{p!(d-p)!} \epsilon^{a_1 \dots a_p a_{p+1} \dots a_d} A_{a_1 \dots a_p} A'_{a_{p+1} \dots a_d} . \quad (2.29)$$

When including Levi-Civita tensors in Lagrangians, we must remember that every time it appears we introduce a violation of parity and time reversal.

2.1.3 Integral Invariants

So far we have discussed invariants that can be defined locally, point by point in spacetime. This is the main requirement to construct a local action. There are, however, other invariants with a non-local character and with important physical interpretation. Two immediate examples coming to mind are the circulation of the vector potential $\oint \vec{A} d\vec{x}$ or the electric or magnetic flux $\oint \vec{E} d\vec{S}$ in electrodynamics. These are rotation invariants which are defined for extended lines and surfaces respectively. One important property they satisfy is that they depend on the particular submanifold of interest, but *do not* depend on the concrete parametrization we may use, i.e. they are intrinsically associated to the geometrical set. The prototypical example is the vector circulation, which admits an immediate generalization to a Lorentz-invariant line integral:

$$I[\gamma] = \int_{\gamma} A \equiv \int_{\gamma} A_a dx^a \equiv \int_{\gamma} A_a(x(\sigma)) \frac{dx^a}{d\sigma} d\sigma , \quad (2.30)$$

which we have written in increasing degree of detail, to show that the integral is not only Lorentz-invariant but also invariant with respect to choices of path parameter σ . On the other hand, this is slightly non-trivial because, trying an obvious generalization to a two-tensor integrated over a two-surface $x^a(\sigma^1, \sigma^2)$

$$\int_{\Sigma^2} \frac{1}{2} T_{ab} dx^a dx^b = \int_{\Sigma^2} \frac{1}{2} T_{ab}(x(\sigma^1, \sigma^2)) \sum_{m,n=1}^2 \frac{\partial x^a}{\partial \sigma^n} \frac{\partial x^b}{\partial \sigma^m} d\sigma^n d\sigma^m , \quad (2.31)$$

the terms containing $(d\sigma^1)^2$ and $(d\sigma^2)^2$ do not correspond to a correct two-dimensional integration measure. Getting rid of these terms by restricting T_{ab} to be antisymmetric produces a standard reparametrization-invariant two-dimensional integral, where $d\sigma^n d\sigma^m$ can be replaced with $[n, m] d\sigma^1 d\sigma^2$, in terms of the permutation symbol in two indices. This pattern generalizes and any covariant antisymmetric tensor, a p -form, can be integrated intrinsically over a p -surface Σ^p immersed in spacetime by functions $x^a(\sigma^1, \dots, \sigma^p)$. The integral invariant is

$$I[\Sigma^p] = \int_{\Sigma^p} A_{[p]} = \int_{\Sigma^p} \frac{1}{p!} A_{a_1 \dots a_p} d\sigma^{a_1 \dots a_p} , \quad (2.32)$$

where

$$d\sigma^{a_1 \dots a_p} = [n_1 \dots n_p] \frac{\partial x^{a_1}}{\partial \sigma^{n_1}} \dots \frac{\partial x^{a_p}}{\partial \sigma^{n_p}} d^p \sigma , \quad (2.33)$$

and we have used the permutation symbol in p indices $[n_1 \dots n_p]$ and $d^p \sigma = d\sigma^1 d\sigma^2 \dots d\sigma^p$. In this form we recognize (2.33) as the volume element on the p -hyperplane tangent to Σ^p , since the vectors $E_r^a \Delta \sigma^r = (\partial x^a / \partial \sigma^r) \Delta \sigma^r$ are tangent to Σ^p and (2.33) computes their determinant,

i.e. the volume of a small polytope having these vectors on its edges. Using Poincaré duality (2.27) we can also write (2.32) as

$$I[\Sigma^p] = (-1)^{\text{sig}} \int_{\Sigma^p} \frac{1}{(d-p)!} \tilde{A}_{a_{p+1} \dots a_d} \tilde{d\sigma}^{a_{p+1} \dots a_d}. \quad (2.34)$$

In this form, the tensor being integrated has indices transverse to the manifold Σ^p , so that we are representing a generalized flux. For $p = d-1$ this formula takes the familiar form of a vector flux over a codimension-one hypersurface,

$$I[\Sigma^{d-1}] = (-1)^{\text{sig}} \int_{\Sigma^{d-1}} \tilde{A}_a dS^a \quad (2.35)$$

with $\tilde{d\sigma}^a = dS^a = n^a dS$ the oriented hypersurface element normal to $\Sigma^{(d-1)}$.

A fundamental object in p -form theory is the *exterior derivative*, which takes a p -form to a $(p+1)$ -form by an antisymmetric derivation

$$(dA_{[p]})_{a_0 a_1 \dots a_p} \equiv (p+1) \partial_{[a_0} A_{a_1 \dots a_p]}. \quad (2.36)$$

It can only act once, since $d^2 = 0$ identically. One of its *raison d'être* is the generalized Stokes' theorem: if $\Sigma^p = \partial V^{p+1}$ is a boundary, then

$$\int_{V^{p+1}} dA_{[p]} = \int_{\Sigma^p} A_{[p]}. \quad (2.37)$$

A mundane example of all these definitions and theorems is the magnetic flux formula through a disc in ordinary space, $D \in \mathbb{R}^3$:

$$\int_D d\vec{S} \cdot \vec{B} = \int_D \frac{1}{2} F_{ij} dx^i dx^j = \oint_{\partial D} d\vec{x} \cdot \vec{A}, \quad (2.38)$$

where one uses (2.35) in the first equality and Stokes' theorem (2.37) in the second equality. We also assume the standard notation $F_{ij} = \partial_i A_j - \partial_j A_i = \sum_k \epsilon_{ijk} B_k$.

The Poincaré dual of the exterior derivative is the *divergence*, which we can denote by $d^\dagger$ and maps p -forms to $(p-1)$ -forms by contracting one index with a derivative:

$$(d^\dagger A_{[p]})_{a_2 \dots a_p} \equiv -\partial^{a_1} A_{a_1 a_2 \dots a_p}. \quad (2.39)$$

The extra sign in this expression and the dagger notation is chosen so that $d^\dagger$ indeed acts as the adjoint of d , in the sense $(A, dA') = (d^\dagger A, A')$, with respect to the inner product

$$(A_{[p]}, A'_{[p]}) \equiv \frac{1}{p!} \int d^d x A^{a_1 \dots a_p} A'_{a_1 \dots a_p}. \quad (2.40)$$

With this definition, one finds that $d^\dagger d^\dagger = 0$ and

$$\tilde{dA}_{[p]} = (-1)^p d^\dagger \tilde{A}_{[d-p]}, \quad (2.41)$$

which allows us to derive analogous versions of Stokes' theorem for divergences of antisymmetric tensors, otherwise known as Gauss' integration theorems. The simplest and most important for

the purposes of these notes is the volume integral of divergence of a current, which is usually written in the form

$$\int_{V^{1+3}} \partial_a J^a = \int_{\partial V^{1+3}} J_a n^a, \quad (2.42)$$

where V^{1+3} is the interior of a tubelike surface of finite extent in time, namely its boundary ∂V^{1+3} has cylinder shape with future and past caps, $\partial V^{1+3} = V_+^3 \cup V_-^3 \cup B^{1+2}$. The caps $V_\pm^3$ are spacelike (normal vector is timelike) and the tube B^{1+2} has everywhere spacelike normal vector. The formula (2.42) is slightly subtle in that the normal vector n^a is outward pointing when it is spacelike ($n_a n^a = 1$) and *inward* pointing when it is timelike ($n_a n^a = -1$).




Figure 2.1: Any increment of a locally conserved charge registered between $V_\pm^3$ must be due to net inward flow through B^{1+2} . Notice the conventions for normal unit vectors.

Then, Gauss' theorem applied to $\partial_a J^a = 0$ gives

$$Q(V_+^3) - Q(V_-^3) = -\text{Flux}(B^{1+2}), \quad (2.43)$$

where $Q(V^3) = -\int_{V^3} J_a \tau^a$, with τ^a the future-directed normal to V^3 , and $\text{Flux}(B^{1+2})$ is the outward flux of J^a through the cylindrical-shaped boundary.⁴ When B^{1+2} is formally taken to infinity, so that $V^3 = \Sigma^3$ is a spacelike, codimension-one submanifold without boundary, we learn that the charge is independent of the choice of Σ^3 (assuming as usual trivial field configurations at infinity). On the other hand, when the $V_\pm^3$ collapse into one another we obtain the form (1.27).

2.1.4 Spinors

In known examples in nature, spinor representations are associated to fermionic fields. These are fundamentally more ‘quantum’ than their bosonic counterparts. By the spin-statistics theo-

⁴The easiest way of checking the (admittedly tricky) sign conventions is to write down (2.42) directly for a straight hypercube.

rem of Quantum Field Theory, we know that the bosonic fields come in representations of integer Lorentz spin, i.e. of tensor type. For this reason we will not dwell on spinor representations in these notes, except for a quick review of standard notation.

The simplest spinor representation is carried by the so-called Dirac spinor, ψ , which transforms in the representation $(1/2, 0) \oplus (0, 1/2)$, of complex dimension four. To construct the Lorentz action in this representation, one usually starts from the four-dimensional Dirac matrices γ^a , satisfying the Clifford algebra,

$$\{\gamma^a, \gamma^b\} = -2\eta^{ab} ,$$

so that

$$\sigma^{ab} = \frac{i}{4}[\gamma^a, \gamma^b]$$

generate the $SO(1, 3)$ Lie algebra. Then, the spinor representation is given by $\psi \rightarrow \mathcal{D}(L)\psi$, where $\mathcal{D}(L)$ is the four-dimensional matrix

$$\mathcal{D}(L) = \exp \left(\frac{i}{2} \sum_{a,b} \theta_{ab} \sigma^{ab} \right) ,$$

and the antisymmetric matrix of parameters θ_{ab} is defined in terms of the Lorentz transformation matrix L , by the relation $L = \exp(\theta')$, where $\theta'^a{}_b = \eta^{ac}\theta_{cb}$. The irreducible components are reached acting with the projectors $P_{\pm} = \frac{1}{2}(1 \pm \gamma_5)$, with $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$. The resulting two-component spinors $\psi_{\pm} = P_{\pm}\psi$ are the two Weyl spinors of opposite chirality (whether ψ_+ is called right- or left-handed depends on the sign convention for γ_5). The two projectors are mapped into one another by parity. Hence, insisting on parity as a good symmetry forces the use of the four-component Dirac spinors.

2.2 Relativistic Particles

In general, a particle in classical physics is any system whose internal structure can be ignored on the scales of interest. For instance, planets are particles if we are interested in the long-range properties of their orbits. Therefore, the minimal description of a particle is given by its path in spacetime, a curve γ embedded in spacetime by functions $x^a(\sigma)$, parametrized by a real variable σ . It is often useful to use a proper-time parametrization $x^a(\tau)$, but this is only possible for massive particles. The only global constraint on the path γ is that it be *causal*, in the sense that the velocity is sub-luminal at all times for massive particles, and exactly at the speed of light when they are massless. In the following sections we describe the basic Lagrangian formalism for particles and their interactions, including their coupling to fields.

2.2.1 Free Particles

We can now recast the particle dynamics in a more covariant language. Let $x^a(\tau)$ be the trajectory of a particle in proper-time parametrization. The four-velocity defined as

$$u^a = \frac{dx^a}{d\tau}$$

is a contravariant four-vector under the Lorentz group, and the free equation of motion may be written as

$$\frac{d^2 x^a}{d\tau^2} = \frac{du^a}{d\tau} = 0 ,$$

solved by straight lines $x^a(\tau) = x_0^a + u^a \tau$, determined by initial conditions x_0^a and u^a .

Conserved energy and momentum can also be assembled into a contravariant four-vector, called the four-momentum $p^a = mu^a$, where $(p^0, \vec{p}) = (E, \vec{p})$. The so-called ‘dispersion’ or ‘mass-shell’ relation $E^2 - \vec{p}^2 = m^2$ may be written in condensed form as $p^2 + m^2 = 0$.⁵ Equivalently, the four-velocity is constrained to satisfy $u^2 = u_a u^a = -1$.

The free particle action over a trajectory γ can be written in the following variety of equivalent forms:

$$S_P = -m \int_{\gamma} d\tau = -m \int_{\gamma} dt \frac{d\tau}{dt} = -m \int_{\gamma} dt \sqrt{1 - \vec{v}^2} = -m \int_{\gamma} dt \sqrt{-\eta_{ab} \frac{dx^a}{dt} \frac{dx^b}{dt}} . \quad (2.44)$$

In particular, the second form shows that the coordinate time t is just an integration parameter which might be redefined at will. In many situations, the freedom to reparametrize the time variable does offer some rewards. So we consider the action in arbitrary parametrization

$$S_P = -m \int d\tau = -m \int d\sigma \frac{d\tau}{d\sigma} = -m \int d\sigma \sqrt{-\eta_{ab} \frac{dx^a}{d\sigma} \frac{dx^b}{d\sigma}} . \quad (2.45)$$

There is a useful trick that simplifies the action by linearizing it and, at the same time, permitting a smooth massless limit, a convenient fact for the discussion of light propagation.

Let us consider the action

$$S_P[x^a, e] = \frac{1}{2} \int d\sigma \left(\frac{1}{e(\sigma)} \eta_{ab} \frac{dx^a}{d\sigma} \frac{dx^b}{d\sigma} - m^2 e(\sigma) \right) , \quad (2.46)$$

where the function $e(\sigma)$ is considered as an independent variable. Its equation of motion $\delta S_P / \delta e = 0$ leads to

$$\frac{dx^a}{d\sigma} \frac{dx_a}{d\sigma} = -m^2 e(\sigma)^2 . \quad (2.47)$$

From this equation we can solve for $e(\sigma)$ and substitute back into (2.46), obtaining the original action (2.45). In this way, we show that both actions are physically equivalent.

The action (2.46) is invariant under reparametrizations of the path (just as (2.45) was) provided we let $e(\sigma)$ transform in such a way that $e(\sigma)d\sigma$ is left invariant. Worldline parameters for which $e(\sigma) = \text{constant}$ are called *affine*. Choosing $e(\sigma) = 1$ and taking the limit $m \rightarrow 0$ produces the equations of motion of a massless particle,

$$\frac{d^2 x^a}{d\sigma^2} = 0 , \quad \frac{dx^a}{d\sigma} \frac{dx_a}{d\sigma} = 0 .$$

The first equation says that the trajectory is ‘straight’ in the σ parametrization, and the second equation says that it lies on the light cone.

⁵Experimental measurements of dispersion relations, $p^2 + m^2 = 0$, provide the most stringent tests of Lorentz symmetry, down to the 10^{-20} level.

For massive particles, it is convenient to choose the proper time as the path parameter, $\sigma = \tau$, so that (2.47) sets $e(\tau) = 1/m$. This means that we can simply use the linearized version of the action

$$S_P = \frac{1}{2} m \int d\tau \left(\eta_{ab} \frac{dx^a}{d\tau} \frac{dx^b}{d\tau} - 1 \right) , \quad (2.48)$$

which is actually the most *naïve* generalization of the Newtonian action. The resulting equations of motion, $d^2x^a/d\tau^2 = 0$, must be supplemented by the field equation of $e(\tau)$, which was lost when setting it to a constant in the action. This equation of motion is simply the mass-shell condition $u^2 = -1$, or equivalently $p^2 + m^2 = 0$. The energy of the particle in a ‘laboratory’ frame is $E = p^0 = -\eta_{ab} p^a u_{\text{lab}}^b$, with $(u_{\text{lab}}^b) = (1, \vec{0})$, so that the Lorentz-invariant expression for the energy measured by an observer at four-velocity u^a is

$$E_u(p) = -\eta_{ab} p^a u^b = -p_a u^a . \quad (2.49)$$

2.2.2 Interacting Particles

Introducing interactions in a relativistic theory of particles is a rather non-trivial issue, given the fact that a limit to the speed of information transfer is built into the very fabric of Special Relativity. This means that standard generalizations of many-particle interaction Lagrangians from Newtonian mechanics are bound to be problematic, since specifying potentials involving particles located at different points in spacetime is a sort of ‘action at a distance’.

There are two basic solutions to this problem. The first solution is to take a phenomenological attitude and consider *contact* interactions in spacetime. In this view, particles are free except for sharply defined collisions at spacetime points, whose physical effect is to change the energy-momentum of each particle $p_{\text{in}}^a \rightarrow p_{\text{out}}^a$ at the collision point, with the constraint of energy-momentum conservation

$$\sum_{\text{colliding particles}} p_{\text{in}}^a = \sum_{\text{colliding particles}} p_{\text{out}}^a . \quad (2.50)$$

A clear shortcoming of this prescription is the arbitrariness of the detailed interaction law, since (2.50) does not determine the precise values of outgoing momenta, given the incoming ones.⁶ A rather more satisfactory solution to the interaction problem is to follow the blueprint of the Maxwell–Lorentz theory and prescribe the particle interactions to be *mediated* by fields. Fields are assumed to satisfy *local* relativistic equations, ensuring that the solutions are waves traveling at most at the speed of light, thus automatically incorporating the required retardation effects between particles. Since the interaction of a fixed background field configuration with a given particle must be locally specified at the particle trajectory, we can make contact with the formal Lagrangian framework by describing such interaction with a world-line relativistic potential of the form

$$S_P = -m \int d\tau - \int d\tau \mathcal{V}(x(\tau), \dot{x}(\tau)) , \quad (2.51)$$

where $\dot{x}^a = dx^a/d\tau$ and we have allowed for the potential to depend on the particle’s four velocity as well as the particle coordinate. In order to find the equations of motion implied by

⁶It is remarkable that this picture reemerges in Quantum Field Theory, where the indefiniteness of the interaction is just an embodiment of the quantum mechanical uncertainty, and the contact interaction described here is literally the so-called *vertex* in a Feynman diagram.




Figure 2.2: *Interaction of particles through fields. Particle on path γ has a kink (change in momentum) which locally produces disturbances on the field $\phi(\vec{x}, t)$ propagating inside the future-directed spacetime cone of bounded velocities, and intersecting path γ' of the second particle at a later time, thereby ‘transmitting’ part of the momentum kick that was carried by the field in the interim. All interactions in the picture are local.*

this action we must be careful to use a general parametrization of the path, i.e. we write (2.51) in the form

$$S_P = -m \int d\sigma \frac{d\tau}{d\sigma} \left[1 + \frac{1}{m} \mathcal{V} \left(x(\sigma), \frac{d\sigma}{d\tau} \frac{dx}{d\sigma} \right) \right] , \quad (2.52)$$

with $d\tau/d\sigma = \sqrt{-\eta_{ab} \frac{dx^a}{d\sigma} \frac{dx^b}{d\sigma}}$.

Varying this action under fluctuations of the trajectory function $x^a(\sigma) \rightarrow x^a(\sigma) + \delta x^a(\sigma)$ and requiring it to be stationary we obtain the equations of motion in the form of a relativistic Newton’s law

$$m \frac{d^2 x^a}{d\tau^2} = F^a , \quad (2.53)$$

where the generalized force four-vector F^a depends on the potential $\mathcal{V}$ in a very complicated and non-linear way. Even for the case of a potential without dependence on four-velocities, $\mathcal{V}_0 = q_0 \phi(x)$, we find the non-trivial velocity-dependent force

$$F_a = -q_0 \partial_a \Phi - q_0 \dot{x}_a \dot{x}^b \partial_b \Phi , \quad (2.54)$$

with

$$\Phi = \frac{m}{q_0} \log \left(1 + \frac{q_0}{m} \phi \right) . \quad (2.55)$$

The result is *very different* from the naive guess for the relativistic generalization of the Newtonian force, which would be $-q_0 \partial_a \phi$.

We can generalize such purely scalar coupling by expanding the general potential $\mathcal{V}(x, \dot{x})$ in powers of four-velocities, obtaining a set of couplings of the particle’s worldline to generalized

fields transforming in symmetric tensor representations of the Lorentz group:

$$S_P = -m \int d\tau - q_0 \int d\tau \phi(x(\tau)) - q_1 \int d\tau \dot{x}^a \phi_a(x(\tau)) - \frac{1}{2} q_2 \int d\tau \dot{x}^a \dot{x}^b \phi_{ab}(x(\tau)) + \dots \quad (2.56)$$

where we have separated conventional numerical coefficients, q_i , called ‘charges’ according to $q_n \phi_{a_1 \dots a_n} = \dot{\partial}_{a_1} \dots \dot{\partial}_{a_n} \mathcal{V}|_{\dot{x}=0}$, and we use the notation $\dot{\partial}_a \equiv \frac{\partial}{\partial \dot{x}^a}$. These terms can be interpreted as the interaction of the particle with an *external* scalar field ϕ , a vector field ϕ_a , a symmetric tensor field ϕ_{ab} , etc. In general, only symmetric tensors can be coupled in this way, unless the particle is decorated with some other degrees of freedom (such as for example a spin four-vector S^a).

The generic relativistic force obtained in this way is extremely non-linear in the external potentials, except for the case of a world-line potential linear in the four-velocities, i.e. $\mathcal{V}(x, \dot{x}) = q_1 \dot{x}^a \phi_a$. In this case, the interaction potential has an intrinsic meaning as a purely geometrical line integral over the trajectory:

$$\int_{\gamma} d\tau \dot{x}^a \phi_a = \int_{\gamma} dx^a \phi_a ,$$

and the relativistic force is linear in velocities,

$$F_a = -q_1 \Phi_{ab} \dot{x}^b , \quad (2.57)$$

and *also linear* in the potentials, through the field strength $\Phi_{ab} \equiv \partial_a \phi_b - \partial_b \phi_a$. In fact, it is only for the coupling to vector fields that we find such simple force laws. The general case is similar to the situation with scalar fields (2.54) in the sense that we are led to extremely non-linear force laws. We shall see later that the vector-field force corresponds to the special case of electromagnetic theory.

The coupling of a system of particles to external fields can be summarized in field-theory Lagrangians by defining appropriate currents. For scalar couplings we have a scalar ‘density’

$$J_S(x) = \sum_p \int d\tau_p q_p \delta^{(4)}(x - x(\tau_p)) , \quad (2.58)$$

which is defined so that ⁷

$$\int d^4x J_S(x) \phi(x) = \sum_p q_p \int d\tau_p \phi(x(\tau_p)) .$$

Vector fields are always coupled via standard vector currents

$$J_V^a(x) = \sum_p \int d\tau_p q_p \dot{x}^a \delta^{(4)}(x - x(\tau_p)) . \quad (2.59)$$

Vector currents are always associated to conservation laws. To see this, we pass to the non-covariant representation by performing the substitution $d\tau_p = dt d\tau_p/dt$ in the proper time

⁷The four-dimensional delta function $\delta^{(4)}(x - x(\tau)) = \delta(t - t(\tau)) \delta^{(3)}(\vec{x} - \vec{x}(\tau))$ is Lorentz-invariant, since it transforms with $|\det L|^{-1} = 1$.

integrals and using the temporal component of the delta function $\delta(t - t(\tau_p))$ to compute the integral over t . The result is the usual expression for the currents

$$J^0(t, \vec{x}) = \sum_p q_p \delta^{(3)}(\vec{x} - \vec{x}_p(t)) , \quad \vec{J}(t, \vec{x}) = \sum_p q_p \vec{v}_p \delta^{(3)}(\vec{x} - \vec{x}_p(t)) . \quad (2.60)$$

The covariant conservation law $\partial_a J^a = 0$ translates then into the usual $\partial_t J^0 + \vec{\partial} \cdot \vec{J} = 0$.

We may also consider higher tensorial generalizations. The most interesting one is the two-index symmetric object

$$J_T^{ab}(x) = \sum_p \int d\tau_p q_p \dot{x}^a \dot{x}^b \delta^{(4)}(x - x(\tau_p)) , \quad (2.61)$$

which actually contains the scalar density above in the trace component, since $\dot{x}^a \dot{x}_a = -1$ implies $\eta_{ab} J_T^{ab} = -J_S$. The symmetric tensor (2.61) acquires an important physical interpretation when $q_p = m_p$, the rest mass of the particles.

2.2.3 The Local Energy-Momentum Tensor

The second order tensor

$$T^{ab}(x) = \sum_p \int d\tau_p m_p \dot{x}_p^a \dot{x}_p^b \delta^{(4)}(x - x(\tau_p)) \quad (2.62)$$

is called the energy-momentum tensor. According to the previous discussion, it measures the response of the system of particles to the introduction of an external field transforming as a symmetric tensor of rank two, and coupling proportionally to the mass of the particles.

The name ‘energy-momentum’ tensor is justified as follows. Using repeatedly the relation

$$\int d\tau F(x(\tau), \dot{x}(\tau)) \frac{dt}{d\tau} \delta^{(4)}(x - x(\tau)) = F(x(t), \dot{x}(t)) \delta^{(3)}(\vec{x} - \vec{x}_p) \quad (2.63)$$

we find that

$$T^{00} = \sum_p E_p \delta^{(3)}(\vec{x} - \vec{x}_p) \quad (2.64)$$

is the energy density of the particle system. In turn,

$$T^{i0} = \sum_p E_p v_p^i \delta^{(3)}(\vec{x} - \vec{x}_p) = \sum_p p_p^i \delta^{(3)}(\vec{x} - \vec{x}_p) , \quad (2.65)$$

which we can interpret either as the energy current, or as the momentum density. On the other hand

$$T^{ij} = \sum_p p_p^i v_p^j \delta^{(3)}(\vec{x} - \vec{x}_p) = \sum_p p_p^j v_p^i \delta^{(3)}(\vec{x} - \vec{x}_p) \quad (2.66)$$

is either the flux of i -th momentum in the j -th direction, or vice versa.

Since T^{ab} characterizes the density and flow of energy, it should be associated to a local conservation law, i.e. to a current that satisfies a ‘continuity’ equation $\partial_a J^a = 0$, just like

the electromagnetic current. The natural current to be associated with the energy-momentum tensor is the four-momentum flux as measured by a local observer with four-velocity u^a ,

$$\mathcal{P}_u^a = -u_b T^{ab} . \quad (2.67)$$

To justify this expression, notice that it reduces to the correct T^{0a} for an observer at rest and, being a tensor equation, it is valid in all Lorentz frames. We define the local current at all points by giving a family of fiducial observers with four-velocity field u^a . If these observers are stationary with respect to one another, all u^a are parallel, and $\partial_a u^b = 0$. Under these conditions, we obtain conservation of the local energy-momentum flux, $\partial_a \mathcal{P}_u^a = 0$, if the energy-momentum tensor satisfies $\partial_a T^{ab} = 0$.

A direct computation for the particle system yields

$$\partial_a T^{ab} = \sum_p m_p \int d\tau_p \dot{x}_p^a \dot{x}_p^b \partial_a \delta^{(4)}(x - x(\tau_p)) . \quad (2.68)$$

Acting on the delta function

$$\dot{x}^a \partial_a \delta^{(4)}(x - x(\tau)) = -\dot{x}^a \frac{\partial}{\partial x^a(\tau)} \delta^{(4)}(x - x(\tau)) = -\frac{d}{d\tau} \delta^{(4)}(x - x(\tau)) ,$$

so that we can write

$$\partial_a T^{ab} = -\sum_p m_p \int d\tau_p \frac{d}{d\tau_p} \left(\dot{x}_p^b \delta^{(4)}(x - x(\tau_p)) \right) + \sum_p m_p \int d\tau_p \ddot{x}_p^b \delta^{(4)}(x - x(\tau_p)) . \quad (2.69)$$

The first term is supported on the endpoints and thus vanishes, whereas the last term can be written

$$\partial_a T^{ab} = \sum_p \int d\tau_p \frac{dp_p^b}{d\tau_p} \delta^{(4)}(x - x(\tau_p)) = \sum_p \frac{dp_p^b}{dt} \delta^{(3)}(\vec{x} - \vec{x}_p) \quad (2.70)$$

as the density of momentum *non-conservation*. Hence, the *local* energy-momentum conservation requires that particles are free $\ddot{x}^a = 0$, or perhaps that their interactions are localized in spacetime (in this case, $\sum_{\text{in}} p = \sum_{\text{out}} p$ at each interaction point). We conclude that, up to regions of measure zero in spacetime, the local conservation of the energy momentum tensor is equivalent to the equation of motion in a system of free particles. This property will be recurrent in all well-defined matter systems. The local conservation of the energy-momentum tensor is of paramount importance in relativistic theories, since it embodies the basic requirements of locality and Lorentz symmetry.

Angular Momentum and Spin

The local conservation of the energy-momentum tensor allows us to define the conserved four-momentum of a system of particles enclosed in a three-dimensional region V^3 (which could be all of $\mathbb{R}^3$).

$$P^a = \int_{V^3} T^{0a} , \quad (2.71)$$

which behaves as the conserved four-momentum of a particle. In particular, we can define the total mass $M = \sqrt{-P^2}$ and the global four-velocity $U^a = P^a/M$. Analogously, we can define the total orbital angular momentum with respect to the origin as

$$L^{ab} = X^a P^b - X^b P^a , \quad (2.72)$$

where $X^a = (t, \vec{X})$ is the “center of mass” trajectory, where

$$\vec{X} = \frac{1}{M} \int_{V^3} \vec{x} T^{00} .$$

The intrinsic angular momentum, or *spin*, is defined as the angular momentum with respect to the center of mass

$$S^{ij} = \int_{V^3} ((x^i - X^i) T^{j0} - (x^j - X^j) T^{i0}) , \quad (2.73)$$

which differs from the total angular momentum with respect to the origin

$$J^{ij} = \int_{V^3} (x^i T^{j0} - x^j T^{i0}) ,$$

by

$$J^{ij} = L^{ij} + S^{ij} .$$

In vector notation, $S^i = \frac{1}{2} \sum_{jk} \epsilon^{ijk} S^{jk}$, i.e. $S^{jk} = \sum_i S^i \epsilon^{ijk}$. We can generalize this to a four-vector by stating that $S^0 = 0$ in the rest frame of the center of mass. Hence, a Lorentz-invariant equation that reduces to these in the rest frame is

$$S_a = -\frac{1}{2} \epsilon_{abcd} S^{bc} U^d . \quad (2.74)$$

The vanishing of the temporal component in the rest frame translates into $U_a S^a = 0$.

The local tensor object

$$M^{abc} \equiv x^a T^{bc} - x^b T^{ac} \quad (2.75)$$

is conserved provided the energy-momentum tensor is *symmetric*, $\partial_c M^{abc} = T^{ba} - T^{ab} = 0$, and defines an angular momentum density $\mathcal{J}_u^{ab} = -u_c M^{abc}$ associated to a spacelike surface orthogonal to the field of four-velocities u^a . On a $t = \text{constant}$ surface it defines the conserved angular-momentum tensor

$$J^{ab} = -J^{ba} = \int_{V^3} M^{ab0} = \int_{V^3} (x^a T^{b0} - x^b T^{a0}) , \quad (2.76)$$

whose spatial components give J^{ij} above. We can project the spin four-vector out of J^{ab} by the formula (Pauli–Lubanski)

$$S_a = -\frac{1}{2} \epsilon_{abcd} J^{bc} U^d , \quad (2.77)$$

since the orbital part L^{ab} drops out from this expression.

Conservation of P^a , J^{ab} and S^a implies that

$$\frac{dU^a}{d\tau_{\text{CM}}} = \frac{dS^a}{d\tau_{\text{CM}}} = 0 , \quad (2.78)$$

where τ_{CM} is the proper time associated to the trajectory of the center of mass. If we abstract the internal structure of the system, we can regard the system as an effective particle with four velocity U^a and spin S^a . Hence, adding a spin degree of freedom is a minimal extension of the notion of point-like particle.

Fluids

It is interesting for applications to take the fluid limit. This is a particle system with very short-ranged collisions, so that a comoving observer sees the system as characterized by an energy density, a pressure, and perhaps gradients of the net velocity field, after one averages over the short-distance and short-time collisions. The fluid is called *perfect* if the comoving observer sees the fluid as isotropic, characterized solely by the density and the pressure:

$$\rho = \left\langle \sum_p E_p \delta^{(3)}(\vec{x} - \vec{x}_p) \right\rangle_{\text{fluid}}, \quad p = \frac{1}{3} \sum_i \left\langle \sum_p p_p^i v_p^i \delta^{(3)}(\vec{x} - \vec{x}_p) \right\rangle_{\text{fluid}}. \quad (2.79)$$

Hence, in the fluid limit, we have $T^{00} = \rho$ and $T^{ij} = p \delta^{ij}$ in the comoving frame.⁸ This determines the compact expression

$$T^{ab}(x)|_{\text{fluid}} = p(x) \eta^{ab} + (p(x) + \rho(x)) U^a(x) U^b(x) \quad (2.80)$$

in a general inertial frame. Here, U^a is the (spacetime dependent) field of four-velocities of the fluid. The conservation equation $\partial_a T^{ab} = 0$ implies, in the non-relativistic limit, $p \ll \rho$, $|\vec{v}| dp/dt \ll |\vec{\partial} p|$, the two Euler equations for the dynamics of a perfect fluid,

$$\partial_t \rho + \vec{\partial}(\rho \vec{v}) = 0, \quad \rho \left(\partial_t \vec{v} + (\vec{v} \cdot \vec{\partial}) \vec{v} \right) = -\vec{\partial} p. \quad (2.81)$$

2.3 Relativistic Fields

A local Lagrangian of fields $\Psi(x)$ in arbitrary representations of the Lorentz group must be constructed as a Lorentz scalar (all indices must be contracted) and, by the Principle of Locality, may be organized by its dependence on derivatives $\mathcal{L}(\Psi, \partial_a \Psi, \partial_a \partial_b \Psi, \dots)$. The terms with fewer derivatives will dominate the long-distance physics, and the terms with progressively higher derivatives will be gradually introduced in a perturbative expansion to recover effects at shorter distances, in the ‘Effective Field Theory’ philosophy outlined in the introduction to these notes.

Once the Lagrangian is specified, the corresponding equations of motion are obtained by requiring the action

$$S = \int \mathcal{L}[\Psi] \quad (2.82)$$

to be stationary under the variations $\Psi \rightarrow \Psi + \delta\Psi$. After repeated application of the identity $\delta(\partial_a \Psi) = \partial_a \delta\Psi$, integration by parts and dropping of boundary terms coming from integrating total derivatives, one finds that stationarity of the action is equivalent to

$$0 = \frac{\partial \mathcal{L}}{\partial \Psi} - \partial_a \frac{\partial \mathcal{L}}{\partial (\partial_a \Psi)} + \partial_a \partial_b \frac{\partial \mathcal{L}}{\partial (\partial_a \partial_b \Psi)} - \dots, \quad (2.83)$$

where the signs alternate according to the number of derivatives acting on Ψ . In most applications, one can restrict to Lagrangians depending polynomially upon one single derivative of the

⁸The fluid averages are defined as $\langle A \rangle_{\text{fluid}} = \Delta V^{-1} \int_{\Delta V} A$ for a small volume element ΔV , still large enough to contain a macroscopic number of particles. Using kinetic theory, one can argue that $p_p^i v_p^i$ is the momentum transfer per unit area and time (i.e. the contribution to pressure) exerted by a particle over a surface orthogonal to the i th direction.

field, $\partial_a \Psi$ (perhaps after repeated application of partial integrations). In this case we obtain the more standard form of the Euler–Lagrange equations, which reduces to the first two terms in (2.83). As indicated above, terms in the Lagrangian with more than two derivatives must be safely considered as perturbative corrections in the sense of the local expansion, and their effect is only to be trusted when they are not dominant.

In what follows, we list the most common types of field theories, as established by their pedagogical features or their importance in the actual description of nature.

2.3.1 Scalar Fields

The prototype relativistic field theory with minimal derivative content is the scalar model (sometimes known as the Klein–Gordon field theory)

$$\mathcal{L}_{\text{KG}} = -\frac{1}{2} \eta^{ab} \partial_a \phi \partial_b \phi - V(\phi) . \quad (2.84)$$

The first term is the only Lorentz-invariant combination of two derivatives of the field, whose detailed structure

$$-\frac{1}{2} (\partial \phi)^2 = \frac{1}{2} (\partial_t \phi)^2 - \frac{1}{2} (\vec{\partial} \phi)^2 , \quad (2.85)$$

shows that the time derivative is the strict ‘kinetic’ term and the spatial derivative is really part of the ‘potential energy’. Up to a total derivative, we also have

$$\mathcal{L}_{\text{KG}} = \frac{1}{2} \phi \partial^2 \phi - V(\phi) + \partial(\dots) , \quad (2.86)$$

which leads to the field equation:

$$\partial^2 \phi - V'(\phi) = 0 . \quad (2.87)$$

One usually separates the linear and quadratic parts of the potential $V(\phi) = J\phi + \frac{1}{2} m^2 \phi^2 + \dots$ to interpret them as the coupling to a scalar ‘source’ and a ‘mass’ term.

To investigate basic properties of the solutions of the field equation, we start with the massless free field, $V(\phi) = 0$. The solution of the field equation $\partial^2 \phi = 0$ is given by a linear superposition of plane waves $\phi_k(x) = \tilde{\phi}_k \exp(ikx) + \text{c.c.}$ of wave vector $\vec{k}$ and frequency $\omega = k^0 = |\vec{k}|$. These waves are defined on all of $\mathbb{R}^{1+3}$, traveling at the speed of light from the infinite past to the infinite future.

The wave vector k^a is orthogonal to the constant-phase surfaces $k^a x_a = \text{constant}$. Consider the integral curves $x^a(\sigma)$ that solve the linear differential equation $dx^a/d\sigma = k^a$ and have, by definition, tangent vector k^a . These curves are orthogonal to the constant-phase surfaces and can be interpreted as trajectories of zero-mass particles, by the condition $k^2 = (dx^a/d\sigma)(dx_a/d\sigma) = 0$. The frequency measured in a rest frame is $\omega = k^0$, so that the frequency measured by an observer with four-velocity u^a is

$$\omega = -u_a k^a . \quad (2.88)$$

We notice the analogy with the formula for the energy of a particle, $E = -p_a u^a$. In the quantum theory, $E = \hbar\omega$ and $\vec{p} = \hbar\vec{k}$ for a photon, so that the particle ‘model’ for light propagation is no longer an analogy but a strict fact.⁹ For the purposes of Special and General Relativity, we

⁹Although we couch this discussion for the case of scalar fields and scalar particles, the propagation properties of massless scalar waves generalize readily to higher spin cases, including electromagnetic waves and the ‘photon’ particles.

can formally treat the propagation of localized, massless wave packets in the ‘ballistic’ picture, as the zero-mass limit of particle propagation with the operator replacement

$$m \frac{d}{d\tau} \rightarrow \frac{d}{d\sigma},$$

where σ is an affine parameter along the ‘light’ ray.

The next level of complication is the case of an external ‘source’ $V(\phi) = J\phi$, with field equation $\partial^2 \phi = J$, solved formally as

$$\phi = \phi_{\text{wave}} + \frac{1}{\partial^2} J, \quad (2.89)$$

where the first term is a solution of the massless wave equation in *vacuo*, $\partial^2 \phi_{\text{wave}} = 0$, and the second term is a formal representation of a particular solution with structure similar to the Newtonian potential, except for the replacement of the operator $\vec{\partial}^2$ by the operator $\partial^2 = -\partial_t^2 + \vec{\partial}^2$, which takes the Poisson equation into the Klein–Gordon equation. Given a purely static solution of the Poisson equation $\vec{\partial}^2 \phi = J$,

$$\phi_{\text{static}}(\vec{x}) = \frac{1}{\vec{\partial}^2} J = - \int \frac{d^3 y}{4\pi} \frac{J(\vec{y})}{|\vec{x} - \vec{y}|},$$

with no time dependence, we can obtain a solution of the relativistic equation by introducing time-dependence through the trick of the retarded potential, i.e. evaluating the source at the retarded time $t - |\vec{x} - \vec{y}|$, to represent the fact that disturbances on the field ϕ travel at the speed of light, as vacuum waves.¹⁰

$$\phi(t, \vec{x})_{\text{retarded}} = \frac{1}{\partial^2} \Big|_{\text{ret}} J = - \int \frac{d^3 y}{4\pi} \frac{J(t - |\vec{x} - \vec{y}|, \vec{y})}{|\vec{x} - \vec{y}|}. \quad (2.90)$$

We can now use this same intuition to interpret the general solution. For an arbitrary potential we may rewrite (2.87) as

$$\phi(t, \vec{x})_{\text{retarded}} = \frac{1}{\partial^2} V'(\phi) \Big|_{\text{retarded}} = - \int \frac{d^3 y}{4\pi} \frac{V'[\phi(t - |\vec{x} - \vec{y}|, \vec{y})]}{|\vec{x} - \vec{y}|}. \quad (2.91)$$

Rather than a general solution, this is an integral equation which we may solve iteratively by redefining the source order by order as a function of the approximate solution at lower orders. In particular, each insertion of a term in $V'(\phi)$ represents a ‘kick’ of the free waves that transport the interaction at the speed of light. In this way, we can understand how successive kicks due to a mass term $V'(\phi) \sim m^2 \phi$ have the collective effect of reducing the speed of propagation from that of light to a superposition of various speeds below light, as if the waves were a superposition of particles with dispersion $E = \sqrt{\vec{p}^2 + m^2}$, a picture that is ultimately vindicated in the quantum version of these field theories.

A nonlinear generalization of (2.84) involving several scalar fields ϕ^I but still two-derivatives is the so-called nonlinear sigma-model, with many applications in particle physics and condensed matter physics

$$\mathcal{L}_{\sigma\text{-model}} = -\frac{1}{2} \sum_{IJ} G_{IJ}(\phi) \partial_a \phi^I \partial^a \phi^J - V(\phi^I). \quad (2.92)$$

¹⁰It should be noted that one can use *advanced* instead of retarded solutions (or a linear combination thereof). The difference between the two is an adjustment of the ϕ_{wave} term in the general solution (2.89). One usually works with the retarded potentials because they are more intuitive in representing causality.

2.3.2 Spinor Fields

Fields in spinor representations of the Lorentz group are also very important in nature (all fermions) but the appropriate Lagrangians require the Dirac formalism and are, strictly speaking, part of the quantum version of the theory. Here, we shall simply quote for completeness the case of the basic free Dirac field. A correct relativistic propagation is ensured if we write

$$(\partial^2 - m^2)\psi = 0$$

as the field equation. However, using the properties of Dirac matrices, one can see that $\partial^2 - m^2 = (i\gamma^a \partial_a + m)(i\gamma^a \partial_a - m)$. Hence, the relativistic equation on spinors with the most general set of solutions is the so-called Dirac equation

$$(i\gamma^a \partial_a - m)\psi = 0,$$

which follows from the Lagrangian

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}(i\gamma^a \partial_a - m)\psi ,$$

where $\bar{\psi} = \psi^\dagger \gamma^0$ is the Lorentz-conjugated representation. The most salient feature of this Lagrangian is its first order character in derivatives, as opposed to the bosonic counterparts.

2.3.3 Vector Fields

Higher-spin tensor fields always lead to subtleties. Consider, for example, a free vector field ϕ^a of mass m , with equation of motion

$$(\partial^2 - m^2)\phi^a = 0 , \tag{2.93}$$

consisting of four copies of the Klein–Gordon equation. A candidate Lorentz-invariant Lagrangian with the correct equations of motion would be

$$\mathcal{L}[\phi^a]_{\text{KG}} = \frac{1}{2} \eta_{ab} \phi^a \partial^2 \phi^b - \frac{1}{2} m^2 \eta_{ab} \phi^a \phi^b \tag{2.94}$$

as a direct generalization of the scalar Klein–Gordon Lagrangian. However, the fact that η_{ab} is *not* positive definite leads to problems, since the whole Lagrangian of the ϕ^0 field has the wrong sign. This field may then store arbitrarily large amounts of negative energy, and any interaction term would render the dynamics unstable. So, we face an interesting dilemma: the field equation following from this Lagrangian seems to be correct and yet the energy density appears to be ill-defined.

We cannot simply drop the offending component ϕ^0 , because that would spoil the Lorentz symmetry. However, for a plane-wave solution in momentum space, $\phi^a = \tilde{\phi}^a e^{ipx}$,

$$(p^2 + m^2)\tilde{\phi}^a = 0 ,$$

we may impose the Lorentz-invariant condition of transversality between the polarization vector $\tilde{\phi}^a$ and the propagation vector p^a , i.e.

$$p_a \tilde{\phi}^a = 0 . \tag{2.95}$$

This condition forces $\tilde{\phi}^0$ to vanish in the rest frame $(p^a) = (m, 0, 0, 0)$. Therefore, we conclude that such ‘transverse waves’ satisfying (2.95) will not have energy stability problems.

This suggests that we add the transversality condition $\partial_a \phi^a = 0$ as an extra equation, together with the Klein–Gordon equation (2.93). In fact, in this case we may redefine (2.93) by any multiple of $\partial_b \partial_a \phi^a$, and the particular choice (known as the Proca Equation)

$$(\eta_{ab}(\partial^2 - m^2) - \partial_a \partial_b) \phi^b = 0 \quad (2.96)$$

does enforce the transversality condition automatically, provided $m \neq 0$. To see this, we take its divergence to find $m^2 \partial_a \phi^a = 0$. The Lagrangian associated to the equation of motion (2.96) is

$$\mathcal{L}_{\text{Proca}} = -\frac{1}{4} \Phi_{ab} \Phi^{ab} - \frac{1}{2} m^2 \phi_a \phi^a, \quad (2.97)$$

where we have defined $\Phi_{ab} = \partial_a \phi_b - \partial_b \phi_a$. We conclude that (2.97) is the consistent Lagrangian of a free vector field, describing the propagation of *three* polarization degrees of freedom.

More subtleties lurk in the zero-mass limit $m \rightarrow 0$. In this case, the ‘rest frame’ of a plane wave does not exist, as it propagates at the speed of light. It is possible to reach the standard null frame $(p^a) = (\omega, 0, 0, \omega)$, but then the transversality condition only enforces $\tilde{\phi}^0 = \tilde{\phi}^3$, which is not enough to cancel out the offending time-like component ϕ^0 . On the other hand, if (2.97) has no energy balance problems, the same should be true of the $m^2 \rightarrow 0$ limit.

There is an interesting escape out of this paradox. Suppose we *declare* by hand that $\tilde{\phi}^0 = 0$ on the standard null frame. This means that, because of the transversality constraint, we are actually demanding the wave polarization to be transverse in the spatial sense, $\sum_i \tilde{\phi}_i p^i = \omega \tilde{\phi}^0 = 0$, with only two polarization degrees of freedom remaining. The problem with this prescription is of course the violence to Lorentz invariance, since the Lorentz transformation of such a polarization vector will in general produce longitudinal and time-like components. However, whatever the Lorentz transformation does, it must preserve the transversality constraint $p^a \tilde{\phi}_a = 0$, which precisely for $p^2 = 0$ has a *redundancy*: if $\tilde{\phi}^a$ is transverse, so is $\tilde{\phi}^a + \tilde{f} p^a$, with $\tilde{f}$ an arbitrary constant. This means that $\tilde{\phi}_a$ will shift by a term proportional to p_a under Lorentz transformations.

The way to remove the offending degree of freedom in a Lorentz-invariant fashion is simply to declare *all* configurations related by such shifts as *physically equivalent*, in the sense that all physical observables should be exactly invariant under $\tilde{\phi}^a \rightarrow \tilde{\phi}^a + \tilde{f} p^a$, or its position-space counterpart

$$\phi_a \longrightarrow \phi_a + \partial_a f. \quad (2.98)$$

This is called *gauge invariance*.

We may then use the degree of freedom in $\tilde{f}$ to remove $\tilde{\phi}_0$ in the standard null frame. In doing so, we remove $\tilde{\phi}_3$ too, as they are linked by the transversality constraint. Therefore, we end up with a consistent Lorentz-invariant theory of massless vector fields with only two physical degrees of freedom. We can find the Lagrangian of this theory by working with the explicitly transverse fields obtained by the linear projection

$$\tilde{\phi}_T^a = \tilde{P}^a{}_b \tilde{\phi}^b \equiv \left(\delta^a{}_b - \frac{p^a p_b}{p^2} \right) \tilde{\phi}^b, \quad (2.99)$$

which translates into a *non-local* projector in position space,

$$\phi_T^a = P^a{}_b \phi^b \equiv \left(\delta^a{}_b - \frac{\partial^a \partial_b}{\partial^2} \right) \phi^b. \quad (2.100)$$

Notice that such transverse fields are automatically gauge-invariant since

$$P^a{}_b \phi^b = P^a{}_b (\phi^b + \partial^b f) . \quad (2.101)$$

Evaluating then the massless Klein–Gordon Lagrangian on transverse gauge-invariant fields we find the Lagrangian

$$\frac{1}{2} \eta_{ab} \phi_T^a \partial^2 \phi_T^b = -\frac{1}{4} \Phi_{ab} \Phi^{ab} + \partial(\dots) , \quad (2.102)$$

corresponding to the massless limit of (2.97). It is important to keep in mind, however, that the massless vector model is assumed to be defined with the built-in redundancy under $\phi_a \rightarrow \phi_a + \partial_a f$, for an arbitrary function $f(x)$, and in particular it has one less degree of freedom than the naive massless limit of the massive model. So, even if (2.102) is the smooth $m^2 \rightarrow 0$ limit of (2.97) at the level of Lagrangians, there is a discontinuous jump in degrees of freedom as we go to the massless limit.

The gauge redundancy is very useful in constructing dynamical laws for massless vector fields. For example, a linear coupling of the form

$$\mathcal{L}_J = \phi_a J^a$$

is bound to break gauge invariance unless we demand the conservation of the ‘current’: $\partial_a J^a = 0$. Should the current *not* be conserved, so that gauge invariance is broken, then the theory with this coupling would simply describe three degrees of freedom instead of two.

The gauge redundancy is often called a ‘local symmetry’, a rather misleading name, since its real role is to ensure that the theory describes the minimal number of degrees of freedom (two for a vector field) even after Lorentz-invariant couplings are specified.

Problem: Gauge Is Not a Symmetry

Consider the massive vector field model with Lagrangian

$$\mathcal{L}_{\text{Proca}} = -\frac{1}{4} \Phi_{ab} \Phi^{ab} - \frac{1}{2} m^2 \phi_a \phi^a . \quad (2.103)$$

Show that it is *not* gauge invariant. Define the so-called Stueckelberg model, which couples a massless vector field and a massless scalar field φ with Lagrangian

$$\mathcal{L}[\phi_a, \varphi] = -\frac{1}{4} \Phi_{ab} \Phi^{ab} - \frac{1}{2} \eta^{ab} (m\phi_a + \partial_a \varphi)(m\phi_b + \partial_b \varphi) . \quad (2.104)$$

Show that the Stueckelberg model is gauge-invariant under the transformations

$$\phi_a(x) \rightarrow \phi_a(x) + \partial_a f(x) , \quad \varphi(x) \rightarrow \varphi(x) - m f(x) .$$

Show that the Stueckelberg model is equivalent to (2.103) by an appropriate fixing of the gauge redundancy.

Hence, we find that a model with three degrees of freedom and no gauge invariance is equivalent to a gauge-invariant model with two degrees of freedom, plus one explicit degree of freedom, so that gauge ‘symmetry’ is not real... just a matter of bookkeeping conventions in describing the true degrees of freedom.

‘Integrate out’ the scalar field φ in (2.104), by solving its equation of motion and substituting back into the action. Show that the resulting equivalent Lagrangian is

$$\mathcal{L}_{\text{non local}} = -\frac{1}{4} \Phi_{ab} \Phi^{ab} + \frac{1}{4} m^2 \Phi_{ab} \frac{1}{\partial^2} \Phi^{ab} . \quad (2.105)$$

That is, a massless vector gauge theory, corrected by a non-local term proportional to the mass squared. Notice that *it is* gauge invariant. The non-local term is the result of the existence of more degrees of freedom than just the two massless polarizations. This non-local model is actually the result of imposing the gauge invariance ‘by hand’ in the naive Klein–Gordon Lagrangian, i.e. show that

$$\mathcal{L}_{\text{non local}} = \mathcal{L}[\phi^a]_{\text{TKG}} = \frac{1}{2} \eta_{ab} \phi_{\text{T}}^a \partial^2 \phi_{\text{T}}^b - \frac{1}{2} m^2 \eta_{ab} \phi_{\text{T}}^a \phi_{\text{T}}^b .$$

Hence, insisting on gauge invariance for a *massive* vector theory requires either a non-local model or a local model with an extra degree of freedom. This is natural, since we know that a massive vector field must have three polarization degrees of freedom; forcing a gauge-invariant description with two polarizations requires the introduction of the third polarization as an explicit field degree of freedom, the so-called Stueckelberg field, a precursor of the Higgs field.

Maxwell in the Lorentz-Invariant Way

We now reproduce the Lagrangian structure of massless vector fields from the direct physical construction of Maxwell’s theory. One can exhibit the Lorentz covariance of Maxwell’s equations by defining the electromagnetic tensor F_{ab} , via

$$F^{0i} = E^i , \quad F_{ij} = \sum_k \epsilon_{ijk} B_k$$

in terms of the electric and magnetic field vectors $\vec{E}, \vec{B}$. Defining also the current four-vector $(J^a) = (\rho_e, \vec{J}_e)$, the equations can be cast into the system

$$\partial_a F^{ab} = -J^b , \quad \epsilon^{abcd} \partial_b F_{cd} = 0 , \quad (2.106)$$

which are manifestly Lorentz-covariant. Thus, we confirm Poincaré’s dictum that the Maxwell system remains untouched in the new relativistic formalism, whereas the Newtonian particle dynamics had to yield. Curiously, Maxwell was the first to write down a relativistic equation in the mid XIX century!

In order to derive these equations from a Lagrangian, we introduce the potential A^a by solving the second Maxwell equation (sometimes referred to as the Bianchi identity) via the Poincaré lemma: locally, an antisymmetric tensor with vanishing antisymmetrized derivative, $\partial_{[a} F_{bc]} = 0$, can always be written as ¹¹

$$F_{ab} = \partial_a A_b - \partial_b A_a \quad (2.107)$$

in terms of some vector field A^a , which always comes with a built-in redundancy, i.e. a gauge ambiguity by the redefinition $A^a \rightarrow A^a + \partial^a f$. The rest of Maxwell’s equations then follow upon varying the Maxwell Lagrangian

$$\mathcal{L}_{\text{Maxwell}} = -\frac{1}{4} F_{ab} F^{ab} + J_a A^a \quad (2.108)$$

¹¹ An explicit proof is obtained by the following *ansatz*: $A^a(x) = -\int_0^1 d\lambda \lambda F^{ab}(\lambda x) x_b$. Notice that the λ integral over points of the form λx requires that the region where we define the fields be ‘star shaped’, in the sense that it can be contracted to a point through straight lines, each of which being fully contained in the original region. This is what is meant by saying that the Poincaré lemma applies *locally*.

with respect to A^a . As explained before, the coupling to the electric current does not spoil gauge invariance (up to total derivatives) provided the current is indeed conserved, $\partial_a J^a = 0$.

A convenient Lorentz-invariant convention to fix partially the gauge redundancy is the Lorenz condition $\partial_a A^a = 0$, whose main virtue is the simplification of the Maxwell equations, simply collapsing them to four Klein–Gordon equations

$$\partial^2 A^a = -J^a , \quad (2.109)$$

whose standard solution was discussed before

$$A^a(t, \vec{x})_{\text{retarded}} = -\frac{1}{\partial^2} J^a \Big|_{\text{retarded}} = \int \frac{d^3 y}{4\pi} \frac{J^a(t - |\vec{x} - \vec{y}|, \vec{y})}{|\vec{x} - \vec{y}|} , \quad (2.110)$$

for fields that vanish at infinity and with causal (retarded) boundary conditions. This relation accounts for most of classical electrodynamics.

Free electromagnetic waves can be added to A^a as solutions of the homogeneous equation with $J^a = 0$. By the linearity property of Maxwell's equations, any such solution is a superposition of plane waves of the form

$$A^a(x) = \tilde{A}^a(k) e^{ikx} + \text{c.c.} ,$$

where $\tilde{A}^a(k)$ is now a constant vector and $kx = \eta_{ab} k^a x^b = -\omega t + \vec{k} \cdot \vec{x}$. The four-vector k^a includes the wave vector and the frequency. Maxwell equations and the gauge condition impose $k^a \tilde{A}_a(k) = 0$ and $k^2 = 0$. By further use of the gauge redundancy one can set $\tilde{A}^0(k) = 0 = \sum_i k^i \tilde{A}^i(k)$, leaving fully transverse waves with two polarization degrees of freedom.

The coupling to charges, given by the linear term in the field-theory action $\int d^4x J_a A^a$, with the *conserved* electromagnetic current

$$J^a(x) = \sum_p e_p \frac{dx^a}{d\tau_p} \delta^{(4)}(x - x_p(\tau_p)) ,$$

produces a correction to the free particle Lagrangian of the form

$$S_P = - \sum_p m_p \int d\tau_p + \sum_p e_p \int d\tau_p A_a \frac{dx^a}{d\tau_p} , \quad (2.111)$$

which in turn leads to the so-called Lorentz force in its fully relativistic incarnation

$$m_p \frac{d^2 x_p^a}{d\tau^2} = e_p F^a_b \frac{dx_p^b}{d\tau} . \quad (2.112)$$

The simplicity of (2.106) and (2.112) is a bit deceptive. Together they form a set of non-linear equations, because the fields determining the motion of the particles do depend on the history of the particles themselves. There are thorny issues of principle, related to the existence of pointlike objects with formally infinite energy density (the particles) and the fact that the energy carried by their own electric field also has mechanical properties. This complicates the calculation of how a particle reacts when it radiates.

Problem: Parity Violating Photons?

Consider adding a Lorentz-invariant term to the Lagrangian of the Maxwell theory of the form

$$\mathcal{L}_\vartheta = \vartheta \tilde{F}_{ab} F^{ab}$$

where

$$\tilde{F}_{ab} = \frac{1}{2} \epsilon_{abcd} F^{cd}$$

is the Poincaré dual. Check that the dual tensor has the roles of electric and magnetic fields reversed. Write the Lagrangian $\mathcal{L}_\vartheta$ in terms of electric and magnetic fields and study its behavior under parity and time reversal. Why is this Lagrangian not usually considered as a correction to Maxwell theory?

2.3.4 Higher Rank Tensor Fields

The construction of field theories in higher tensor representations reproduces some of the problems already present in the vector case. Lorentz invariance tends to introduce unstable modes, which have to be dealt with via constraints and gauge redundancies. There are general no-go arguments that limit the possibilities of constructing such field theories if we require long-range interactions and, in any case, they do not appear in our current fundamental description of nature, except for two cases, both of them related to rank-two tensor fields.

We consider here the massless case, which is the most interesting from the physical point of view. A naive action for a massless two-index tensor field ϕ_{ab} of the form

$$\mathcal{L}[\phi_{ab}]_{\text{KG}} = \frac{1}{2} \phi_{ab} \partial^2 \phi^{ab}$$

has the proverbial stability issues, which may be solved by introducing gauge redundancies. In particular, for the antisymmetric case, $b_{cd} = -b_{dc}$, we introduce a redundancy by an antisymmetric derivative $b_{cd} \rightarrow b_{cd} + \partial_{[c} f_{d]}$, whereas the symmetric case, $h_{ab} = h_{(ab)}$, requires a corresponding ‘symmetric’ redundancy $h_{ab} \rightarrow h_{ab} + \partial_{(a} f_{b)}$.

The resulting gauge-invariant action for the antisymmetric case is a simple generalization of the Maxwell construction:

$$\mathcal{L}[b_{cd}] = -\frac{1}{12} H_{abc} H^{abc}, \quad H_{abc} \equiv 3 \partial_{[a} b_{bc]}. \quad (2.113)$$

Here, H_{abc} is a generalization of the Maxwell tensor F_{ab} and b_{ab} is a generalization of the potential four-vector A_a . The equation of motion $\partial_a H^{abc} = 0$ implies that the pseudovector Poincaré-dual field $\tilde{H}_a = -\frac{1}{6} \epsilon_{abcd} H^{bcd}$ has vanishing antisymmetric derivative and, using Poincaré’s lemma, it admits the representation $\tilde{H}_b = \partial_b \tilde{a}$ as the gradient of a pseudoscalar field $\tilde{a}$, usually called the *axion*.¹² Then, the Bianchi identity $\partial_{[a} H_{bcd]} = 0$ is equivalent to the standard wave equation for the axion: $\partial^b \partial_b \tilde{a} = 0$. This chain of arguments shows that the theory (2.113) only has *one* physical polarization degree of freedom, that is to say, it is equivalent to the theory of a massless pseudoscalar field.

If we try the same strategy with the 3-form field C_{abc} with field strength $D_{abcd} = 4\partial_{[a} C_{bcd]}$, the equation of motion is equivalent to requiring $\partial_a \tilde{D} = 0$, where $\tilde{D}$ is the Poincaré dual of

¹²A pseudoscalar field is a scalar under small Lorentz transformations which changes sign under a parity transformation. The axion is a hypothetical particle whose existence is not established, but pseudoscalar fields are routinely used to describe, for instance, the π -mesons.

D_{abcd} . Since the solutions are constant fields, there is no interesting propagating degree of freedom associated to this theory. This pattern repeats in d dimensions: a p -form gauge field is dual to a $(d - p - 2)$ -form, so that $(d - 2)$ -forms describe a single scalar degree of freedom and $(d - 1)$ -forms carry no propagating degrees of freedom.

The symmetric case is more complicated, and more interesting for the theory of gravity. There are four independent Lorentz-invariant structures at the quadratic, two-derivative level:

$$\partial_a h_{bc} \partial^a h^{bc}, \quad \partial_a h^{ab} \partial^c h_{cb}, \quad \partial_a h_b^b \partial^a h_c^c, \quad \partial_a h_c^c \partial_b h^{ab},$$

up to total derivatives. Imposing now gauge invariance under $h_{ab} \rightarrow h_{ab} + \partial_{(a} f_{b)}$ one finds a unique answer up to overall normalization, the so-called Fierz–Pauli Lagrangian

$$\mathcal{L}[h_{ab}]_{\text{FP}} = -\frac{1}{4} \partial_a h_{bc} \partial^a h^{bc} + \frac{1}{4} \partial_a h_c^c \partial^a h_d^d - \frac{1}{2} \partial_a h_c^c \partial_b h^{ab} + \frac{1}{2} \partial_a h^{ab} \partial^c h_{bc}, \quad (2.114)$$

which will turn out to be the linearized approximation to Einstein’s theory! Therefore, it describes linearized gravitational waves. For that reason, we will defer further discussion to the section on gravitational radiation.

2.4 Symmetry and the Field

One of the great advantages of the Lagrangian formalism is the efficient handling of symmetries. This fact is accentuated in the case of field theories. The powerful theorem of Noether which links symmetry and conservation becomes an essential tool in analyzing the general complicated dynamics of field theories. In this section we start with a covariant proof of a general version of Noether’s theorem, and then we focus on the example of major interest: the definition of the energy-momentum tensor for general field theories.

2.4.1 Noether

Let $\mathcal{L}[\Psi]$ be a general functional of fields $\Psi(x)$ in arbitrary tensor representations of the Lorentz group, with polynomial dependence on ∂_a , but otherwise generic. We consider a continuous symmetry whose action on the fields, $\Psi \rightarrow \Psi'$, when expanded close to the identity is

$$\Psi'(x) - \Psi(x) = \delta_s \Psi(x) + O(\epsilon^2) = \epsilon s[\Psi] + O(\epsilon^2), \quad (2.115)$$

where the functional $s[\Psi]$ is also assumed to have polynomial dependence on derivatives. Neglecting boundary terms coming from total derivatives, invariance of the action implies that the Lagrangian density transforms as a total derivative $\delta_s \mathcal{L} = \epsilon \partial_a f_s^a[\Psi]$.

We will now implement a formal trick. Consider a *more general* transformation where the symmetry parameter is given a functional dependence on spacetime coordinates, i.e. $\delta_s \Psi = \epsilon(x) s[\Psi]$. The variation of the Lagrangian density can be expanded in derivatives of $\epsilon(x)$ as

$$\delta_s \mathcal{L} = \epsilon(x) K_s[\Psi] + \partial_a \epsilon(x) K_s^a[\Psi] + \partial_a \partial_b \epsilon(x) K_s^{ab}[\Psi] + \dots, \quad (2.116)$$

for some functionals K_s, K_s^a , etc. In particular, by setting $\epsilon = \text{constant}$ we learn that

$$K_s[\Psi] = \partial_a f_s^a[\Psi]. \quad (2.117)$$

We emphasize that the spacetime dependence of $\epsilon(x)$ is just a mathematical trick, not a generalization from a global to a local symmetry. Indeed, (2.116) shows that the Lagrangian is *not* invariant under the local transformation. The action is only invariant under the global transformation with constant ϵ .

Neglecting total derivatives (by choosing for instance $\epsilon(x)$ with compact support), we may now integrate by parts (2.116) and find

$$\delta_s \int \mathcal{L}[\Psi] = - \int \epsilon(x) \partial_a J_s^a, \quad (2.118)$$

where

$$J_s^a = -f_s^a[\Psi] + K_s^a[\Psi] - \partial_b K_s^{ab}[\Psi] + \dots \quad (2.119)$$

Since the variation $\delta_s \Psi = \epsilon(x)s[\Psi]$ is a particular case of a general variation of the fields, on a configuration that extremizes the action we conclude that this current is conserved, $\partial_a J_s^a = 0$. Hence, in (2.119) we have a constructive definition of the Noether current, valid in an arbitrary derivative expansion. To construct the functionals $K_s^a, K_s^{ab}, \dots$ appearing in (2.116) one can make repeated use of the identity

$$\delta_s(\partial_a \Psi) = \partial_a(\delta_s \Psi) = \partial_a(\epsilon(x)s[\Psi]) = \epsilon(x)\partial_a s[\Psi] + s[\Psi]\partial_a \epsilon(x)$$

in the formal expansion

$$\delta_s \mathcal{L} = \frac{\partial \mathcal{L}}{\partial \Psi} \delta_s \Psi + \frac{\partial \mathcal{L}}{\partial(\partial_a \Psi)} \delta_s(\partial_a \Psi) + \frac{\partial \mathcal{L}}{\partial(\partial_a \partial_b \Psi)} \delta_s(\partial_a \partial_b \Psi) + \dots$$

In the simple case where the Lagrangian depends polynomially on a single derivative, the only non-vanishing functional is

$$K_s^a[\Psi] = \frac{\partial \mathcal{L}}{\partial(\partial_a \Psi)} s[\Psi]. \quad (2.120)$$

The resulting Noether current is then the familiar expression

$$J_s^a = \frac{\partial \mathcal{L}}{\partial(\partial_a \Psi)} s[\Psi] - f_s^a[\Psi]. \quad (2.121)$$

For an illustrative example of how Noether's theorem works in field theory, consider a set of N scalars with Lagrangian

$$\mathcal{L}_{O(N)} = -\frac{1}{2} \sum_{I=1}^N \partial_a \phi_I \partial^a \phi_I, \quad (2.122)$$

which is invariant under the $O(N)$ transformations $\phi_I(x) \rightarrow \sum_J R_{IJ} \phi_J(x)$ for any constant orthogonal matrix R_{IJ} . To obtain the Noether current we work near the identity $R_{IJ} = \delta_{IJ} + \Omega_{IJ} + O(\Omega^2)$ with Ω_{IJ} antisymmetric. Picking a basis of $N(N-1)/2$ matrices T_{IJ}^s for the Lie algebra of $O(N)$ we can write $\Omega_{IJ} = \sum_s \epsilon_s T_{IJ}^s$ and the corresponding $N(N-1)/2$ Noether currents taking the form

$$J_a^s = \sum_{IJ} \phi_I T_{IJ}^s \partial_a \phi_J \quad (2.123)$$

are conserved $\partial^a J_a^s = 0$.

The previous derivation implicitly defines the field variations as *functional* variations, in the sense that a field configuration is compared to the transformed field configuration *at the same point*: $\delta\Psi(x) + O(\epsilon^2) = \Psi'(x) - \Psi(x)$. This is the standard situation for symmetries which do not affect the spacetime coordinates (called *internal symmetries*). On the other hand, spacetime symmetries are often presented in tensorial form, which does involve comparing functions at different values of the variables:

$$\Psi'^{\dots a \dots}_{\dots b \dots}(x') = \dots \frac{\partial x'^a}{\partial x^c} \Big|_x \dots \frac{\partial x^d}{\partial x'^b} \Big|_{x'} \dots \Psi^{\dots c \dots}_{\dots d \dots}(x) , \quad (2.124)$$

where we have been very explicit about the functional dependence of all pieces on the coordinates. This formula gives the transformation law for any tensor field under a Poincaré transformation $x^b \rightarrow x'^b = a^b + L^b_c x^c$. In the *active* view of the transformations, it gives the relation between the tensor *functions* at different points in Minkowski spacetime, labeled x and x' .

Therefore, in order to use (2.124) to derive $\delta\Psi(x)$, we need to expand it near the identity and evaluate the variation at the *same* point. In general, any continuous transformation of the coordinates can be written in infinitesimal form as

$$x'^a = x^a - \epsilon \xi^a(x) + O(\epsilon^2) , \quad (2.125)$$

for some vector field $\xi^a(x)$. Inserting this expression into (2.124) and keeping only terms linear in ϵ we find

$$\delta_\xi \Psi = \epsilon \mathcal{L}_\xi \Psi , \quad (2.126)$$

where $\mathcal{L}_\xi \Psi$ is the so-called *Lie derivative* along the vector field $\xi^a(x)$:

$$\mathcal{L}_\xi \Psi^{\dots a \dots}_{\dots b \dots} = \xi^c \partial_c \Psi^{\dots a \dots}_{\dots b \dots} - \dots - \partial_c \xi^a \Psi^{\dots c \dots}_{\dots b \dots} - \dots + \partial_b \xi^d \Psi^{\dots a \dots}_{\dots d \dots} + \dots . \quad (2.127)$$

The Lie derivative is essentially a directional derivative along the vector field, corrected by terms that depend on the spacetime variation of the vector field itself, with signs that depend on the covariant or contravariant nature of the indices.

With this definition of the field variations, we can write the Noether current for any spacetime symmetry $\delta_\xi x^a = -\epsilon \xi^a(x)$ reinterpreting (2.121) in the form

$$J_\xi^a = \frac{\partial \mathcal{L}}{\partial(\partial_a \Psi)} \mathcal{L}_\xi[\Psi] - f_\xi^a . \quad (2.128)$$

Finally, when a spacetime symmetry has also a simultaneous ‘internal’ action beyond the tensorial form (2.124) we can write the field variation to linear order in ϵ as $\delta_\xi \Psi = \epsilon s_\xi[\Psi] + \epsilon \mathcal{L}_\xi[\Psi]$ and we can write for the corresponding Noether current:

$$J_\xi^a = \frac{\partial \mathcal{L}}{\partial(\partial_a \Psi)} (s_\xi[\Psi] + \mathcal{L}_\xi[\Psi]) - f_\xi^a , \quad (2.129)$$

a formula that one encounters when discussing scale and conformal transformations.

Armed with these expressions for the conserved currents, we could write down all the canonical forms of currents associated to the ten independent symmetries of the Poincaré group. The most important of all is the local energy-momentum tensor, T_{ab} , which is none other than the set of Noether currents for translational symmetry. Once T_{ab} is known, it can be used to build all other currents in a constructive fashion.

Noether and the Gauge

The mathematical trick of using a spacetime-dependent parameter $\epsilon(x)$ brings up the following question. What would it take to make the theory invariant under the *local* transformation? Considering the simple case of Lagrangians with explicit dependence on first derivatives of the field, we saw that the variation of the action can be written as

$$\delta_s \int \mathcal{L} = \int J_s^a \partial_a \epsilon(x) , \quad (2.130)$$

that is to say, the Noether current characterizes directly the lack of invariance when the symmetry parameter ϵ is given a functional dependence on spacetime position. A way to make the theory symmetric is to add a new term to the Lagrangian, depending on an extra vector field A_s^a , through the coupling

$$\mathcal{L} \rightarrow \mathcal{L} - \eta_{ab} J_s^a A_s^b , \quad (2.131)$$

where the vector field A_s is required to transform as

$$\delta_s A_s^a = \partial^a \epsilon(x) + \dots , \quad (2.132)$$

so that the complete Lagrangian is now invariant under the local transformation (the dots stand for extra terms that may be needed but would not depend on derivatives of $\epsilon(x)$). This procedure is called the Noether method of gauging, and it manufactures a gauged version out of any global symmetry. We may leave A_s as a mere background field, or we can postulate new terms in the Lagrangian that provide a kinetic term for A_s , such as a Maxwell term in terms of the field strength $\partial^a A_s^b - \partial^b A_s^a$, giving us a fully fledged gauge theory. Notice that the resulting Lagrangian for the gauge field must respect the local symmetry implying, among other things, that it must be massless.

2.4.2 Energy-Momentum Tensor in Field Theories

Having defined the energy-momentum tensor of a particle system with contact interactions, we can also monitor the local exchanges of energy and momentum in the more physical case where the interactions are mediated by fields. Consider for example a system of charged particles whose electromagnetic interactions are governed by Lorentz forces

$$m_p \ddot{x}_p^a = e_p F^a{}_b \dot{x}_p^b . \quad (2.133)$$

Then, the energy-momentum tensor of the particle system is not conserved alone, as some energy flows into the field degrees of freedom. To see this, we insert (2.133) into (2.69) to find

$$\partial_b T_{\text{particles}}^{ab} = F^a{}_b J^b . \quad (2.134)$$

Thus, local energy conservation requires that we add the intrinsic energy-momentum tensor of the electromagnetic field, canceling the source term in (2.134). We know from elementary discussions of electrodynamics that the energy density of the electromagnetic field is given by $\frac{1}{2}(\vec{E}^2 + \vec{B}^2)$, i.e. it is quadratic in components of the electromagnetic tensor F_{ab} . This follows immediately from a glance at the basic Lagrangian (1.47); since the electric field is proportional to the time derivative of the vector potential, the structure of the Lagrangian suggests the

structure of the energy density. Hence we may try a quadratic *ansatz* for the energy-momentum tensor of the electromagnetic field:

$$c_1 F^{ac} F_c^b + c_2 \eta^{ab} F_{cd} F^{cd}$$

depending on the two independent quadratic combinations of F_{ab} which are allowed by Lorentz symmetry and the antisymmetry of the electromagnetic tensor. Imposing now the condition

$$\partial_b T_{\text{Maxwell}}^{ab} = -F^{ab} J_b$$

given the equations of motion $\partial_a F^{ab} = -J^b$ we fix the constants $c_1 = -1$ and $c_2 = -1/4$ to obtain the Maxwell energy-momentum tensor

$$T_{\text{Maxwell}}^{ab} = -F^{ac} F_c^b - \frac{1}{4} \eta^{ab} F_{cd} F^{cd} . \quad (2.135)$$

which recovers the standard expressions for the energy density, $T^{00} = \frac{1}{2} (\vec{E}^2 + \vec{B}^2)$, and energy flux, $T^{0i} = (\vec{E} \times \vec{B})^i$, the so-called Poynting vector.

This *ansatz* method is efficient in finding the expression for T_{ab} . It is often the case that listing the Lorentz structures with two derivatives and requiring conservation $\partial_a T^{ab} = 0$ fixes the energy-momentum tensor for most field theories of interest. For instance, for the Klein–Gordon scalar model (2.84) we find in this way

$$T_{\text{KG}}^{ab} = \partial^a \phi \partial^b \phi - \eta^{ab} (\frac{1}{2} (\partial \phi)^2 + V(\phi)) . \quad (2.136)$$

As a check, we see that T_{00} indeed gives the energy density of the scalar field

$$T_{00} = \rho_\phi = \frac{1}{2} (\partial_t \phi)^2 + \frac{1}{2} (\vec{\partial} \phi)^2 + V(\phi) .$$

The energy-momentum tensor of the Maxwell field has the important property of being *traceless*, $T^a_a = 0$. For the massive vector theory this trace is instead proportional to the ‘photon mass’:

$$T_{\text{Proca}}^{ab} = T_{\text{Maxwell}}^{ab} + m^2 \left(A^a A^b - \frac{1}{2} \eta^{ab} A_c A^c \right) , \quad \eta_{cd} T_{\text{Proca}}^{cd} = -m^2 A_a A^a .$$

The trace of the energy-momentum tensor is in general sensitive to the mass terms in the problem. For example, for a system of particles, we already noticed that the trace of the energy-momentum tensor is controlled by the scalar density of ‘rest mass’:

$$T^a_a \Big|_{\text{particles}} = -J_m = - \sum_p \int d\tau_p m_p \delta^{(4)}(x - x(\tau_p)) .$$

An important qualification of this rule is the case of massless scalars, for which T^a_a *does not* vanish except in two dimensions.

Problem: Energy-Momentum Conservation and Equations of Motion

Show that the local conservation of the energy-momentum tensor, $\partial_a T^{ab} = 0$, implies the equations of motion for both scalar and electromagnetic fields, provided certain genericity conditions are met. Namely the equations of motion are satisfied except at points where $\partial_a \phi = 0$ in the scalar field case. For Maxwell fields one must require that the matrix F_b^a be invertible.

The Noether Way

On general grounds, energy and momentum are ultimately defined as Noether charges for the symmetries of time and space translations. So we can determine the energy-momentum tensor as a set of Noether currents for Minkowski spacetime translations following the formula (2.128).

Let the translation along the x^b direction be expressed as $\delta_b x^a = -\epsilon \xi_{(b)}^a$, for the vector field $\xi_{(b)}^a = \delta_b^a$. Since any Lagrangian density is a Lorentz scalar, we have

$$\delta_b \mathcal{L} = \epsilon \mathcal{L}_{\xi_{(b)}} \mathcal{L} = \epsilon \xi_{(b)}^a \partial_a \mathcal{L} = \epsilon \partial_b \mathcal{L} = \epsilon \partial_a (\delta_b^a \mathcal{L}) ,$$

so that the Lagrangian varies by a total derivative with $f_{\xi_{(b)}}^a = \delta_b^a$. The variation of any Lorentz-tensor field is given by

$$\delta_b \Psi = \epsilon \mathcal{L}_{\xi_{(b)}} \Psi = \epsilon \xi_{(b)}^c \partial_c \Psi = \epsilon \partial_b \Psi ,$$

where we have suppressed the Lorentz indices of Ψ and used the fact that $\xi_{(b)}^a$ has vanishing derivatives. Applying then the general formula (2.128) we obtain a set of four Noether currents

$$\mathcal{J}^a{}_{(b)} = \frac{\partial \mathcal{L}}{\partial (\partial_a \Psi)} \partial_b \Psi - \delta_b^a \mathcal{L} . \quad (2.137)$$

Evaluating for the scalar case and comparing with (2.136) we find that

$$T_{ab} = -\mathcal{J}_{a(b)} . \quad (2.138)$$

Notice the different status of the indices in this expression. The index a is an ordinary vector index of each current, but the index b is a label of the particular current we are considering. In fact, the so-defined tensor T_{ab} is not symmetric in general. To see the tension more explicitly, we can now apply the formalism to the Maxwell theory, with Lagrangian

$$\mathcal{L}_{\text{Maxwell}} = -\frac{1}{4} F_{ab} F^{ab} .$$

The naive energy-momentum tensor for $\delta_a A_b = \epsilon \partial_a A_b$ turns out to be given by

$$F_a{}^c \partial_b A_c + \eta_{ab} \mathcal{L}_{\text{Maxwell}} , \quad (2.139)$$

which is neither symmetric nor gauge-invariant. The lack of symmetry is inconvenient, but the lack of gauge-invariance is just not acceptable, since the energy momentum tensor gives physical energy and momentum densities, and these cannot depend on our choice of vector potential. We can fix both problems by defining a gauge-invariant field variation:

$$\bar{\delta}_b A_c \equiv \delta_b A_c + \delta'_b A_c$$

with $\delta'_b A_c \equiv \partial_c (-\epsilon A_b)$ an appropriate field-dependent infinitesimal gauge transformation. The new gauge-invariant translation is $\bar{\delta}_b A_c = \epsilon F_{bc}$ which, if used in the Noether procedure, yields the correct gauge-invariant energy-momentum tensor,

$$T_{\text{Maxwell}}^{ab} = -F_c^a F^{cb} + \eta^{ab} \mathcal{L}_{\text{Maxwell}} . \quad (2.140)$$

An interesting comment to make is that the difference between the naive (2.139) and the true (2.140) Maxwell energy-momentum tensors is the term $F_{ca} \partial^c A_b$ which, up to the equation

of motion $\partial^c F_{ca} = 0$, is equivalent to the total derivative $\partial^c(F_{ca}A_b)$. This is an example of a subterfuge called ‘improvement’. Just as Lagrangians are not completely determined by the equations of motion, allowing redefinitions by total derivatives, local currents are not completely determined by the sole requirements of symmetry and conservation. In principle, any additive redefinition by a local functional (depending on derivatives of fields at most polynomially) which is a total derivative and preserves conservation (using the equations of motion), introduces a valid current. Thus for any Noether current J_s^a , the ‘Noether ambiguity’ $J_s^a \rightarrow J_s^a + \partial_c N^{[ca]}$ allows us to redefine the current, provided the tensor $N^{[ca]}$ is antisymmetric to ensure conservation. Being a total derivative, the added term does not affect the global charges obtained by integrating the current over any constant-time hypersurface (assuming as usual that we do not pick boundary terms at infinity).

The Noether ambiguity for the energy-momentum tensor is any additive redefinition of the form

$$T_{ab} \longrightarrow T_{ab} + \partial^c N_{[ca]b} , \quad (2.141)$$

where the tensor N_{cab} is assumed antisymmetric in the first two indices. For the Maxwell case we had $N_{cab} = F_{ca}A_b$. More generally, it was shown by Belinfante and Rosenfeld in 1940 that one can always use the Noether ambiguity (2.141) to *improve* the canonical energy momentum tensor (2.138) into a symmetric one.

Why the insistence on a symmetric energy-momentum tensor? One reason is pure convenience. As indicated in (2.75) in the section on the energy-momentum of particle systems, given a conserved and *symmetric* T_{ab} , the set of currents $M^{abc} \equiv x^a T^{bc} - x^b T^{ac}$ are automatically conserved $\partial_c M^{abc} = 0$ and their associated charges,

$$J^{ab} = \int_{\mathbb{R}^3} M^{ab0} , \quad (2.142)$$

define the relativistic generalization of angular momenta. They give literally the angular momenta for spatial indices J^{ij} , and give the charges associated to the Lorentz boosts in the J^{0i} components. Thus, the set T^{ab}, M^{abc} gives the complete list of currents of the ten-parameter Poincaré isometries, and all of this is still true for any field theory whose energy-momentum tensor is symmetric. Notwithstanding all this, the most important reason for insisting on symmetric T_{ab} is the starring role of the energy-momentum tensor in Einstein’s theory of gravitation. It plays for gravity the same role that the electric current plays for electromagnetism, but the version of the energy-momentum tensor that enters the Einstein equations is *always* a symmetric one. Therefore, it is reassuring that one can always choose a symmetric T_{ab} for any reasonable system of particles and fields.

The art of energy-momentum tensor improvement does not stop at Noether ambiguities though. As we noticed above, the relation between the absence of masses and the vanishing trace $T_a^a = 0$ is broken in the case of massless Klein–Gordon field, which satisfies

$$T_{\text{KG}}{}^a{}_a = -(\partial\phi)^2 .$$

There is, however, an improvement of the energy-momentum tensor that makes it traceless (as always, after using the equations of motion). It is given by ¹³

$$T_{ab} \longrightarrow T_{ab} - \frac{1}{6} (\partial_a \partial_b - \eta_{ab} \partial^2) \phi^2 . \quad (2.143)$$

¹³Introduced in 1970 by C. Callan, S. Coleman and R. Jackiw.

This improvement term is different from the Noether ambiguity types but, like those, it also preserves the local conservation and it is a total derivative, so that it does not contribute to global conserved charges.

Problem: Scale Invariance

Find the transformation rules that make the action of a free massless scalar field invariant under the scale transformations (dilatations) $x^a \rightarrow e^{-\epsilon} x^a$, with real ϵ . Show that a mass term breaks the symmetry but there exists one particular polynomial potential which still preserves scale invariance. Apply Noether's theorem to this theory to define the conserved dilatation current D^a . How is it related to the energy-momentum tensor? Find an improvement that brings the dilatation current to be of the form $D^a = x_b T^{ba}$, which is indeed conserved when $T_a^a = 0$.

Problem: Conformal Invariance

This problem extends the previous one. A *conformal transformation* of Minkowski spacetime is a coordinate transformation that preserves the metric up to a local rescaling, $\eta_{ab} dx'^a dx'^b = \Omega^2(x) \eta_{ab} dx^a dx^b$.

- (a) Show that the infinitesimal transformations $x'^a = x^a - \epsilon \xi^a(x)$ with this property are generated by vector fields *called conformal Killing vectors*, satisfying $\partial_a \xi_b + \partial_b \xi_a = \frac{1}{2} \eta_{ab} \partial_c \xi^c$. Find the general solution and show that, besides Poincaré transformations and dilatations, it contains four *special conformal transformations* that can be written as ITI where T is an ordinary translation and I is the inversion $x^a \rightarrow x^a/x^2$.
 - (b) Generalize to d spacetime dimensions. What happens for $d = 2$?
 - (c) Consider a scale-invariant field theory with a symmetric, conserved energy-momentum tensor T^{ab} , whose dilatation current has the form $D^a = x_b T^{ba} + V^a$, with V^a a local operator (the *virial current*). Show that $T^a_a = -\partial_a V^a$. Show that if $V^a = \partial_b L^{ba}$ for some local L^{ab} , then T^{ab} can be improved into a conserved and traceless $\tilde{T}^{ab}$. Show that in this case $J_\xi^a = \tilde{T}^{ab} \xi_b$ is a conserved current for *every* conformal Killing vector ξ , and write down the currents for dilatations and special conformal transformations.
 - (d) Apply these results to the massless scalar field in d dimensions and to the Maxwell theory.
-

Chapter 3

The Principle of Equivalence

It is certainly tempting to apply the previous machinery of relativistic field theory to the problem of gravitation. That is to say, to postulate some tensorial potential that generalizes the Newtonian gravitational potential and just search for the right field theory in Minkowski spacetime that describes the phenomenology of gravitation. This straightforward route, which gives preference to building Lorentz-invariant field theories, can be followed to finally land in General Relativity, but it soon becomes a winding path full of technicalities. History took a different route, when Einstein found a famous shortcut which took him straight into the answer within a decade.¹ The key to the shortcut was the Principle of Equivalence, a fact about gravity that was known to Galileo and Newton, and its connection with the formalism of Riemannian geometry. In this chapter we discuss this crucial chain of ideas, but first we outline the first steps of the longer ‘Lorentz path’.

Given the formal similarity between the Newton potential and the Coulomb potential, it looks natural to follow the blueprint of Maxwell’s theory to build a theory of gravity, since we know that Maxwell’s theory is ultimately ‘relativistic’. We may begin by postulating a vector-like ‘gravitational potential’ g_a with field strength $G_{ab} = \partial_a g_b - \partial_b g_a$ and a time-like component proportional to the Newtonian potential $g^0 \sim \phi_N$. Actually, such a theory was contemplated by Maxwell himself in the XIX century, although of course he did not know anything about Lorentz invariance. An obvious problem is that the analog of electric charges must be the masses, but they are all positive, so that the ‘gravitoelectric’ static force should be repulsive rather than attractive. Maxwell toyed with the idea that one could turn the static force into an attractive one by inverting the sign of Newton’s constant, $G \rightarrow -G$, but he noticed that this would assign large amounts of negative energy to the gravitational field itself, and finally abandoned these ideas.

A much better attempt at a relativistic generalization of Newton’s theory would promote ϕ_N to a fully fledged Lorentz *scalar* field $\varphi(x)$ with Lagrangian

$$\mathcal{L}_{G\text{-scalar}} = -\frac{1}{8\pi G}(\partial\varphi)^2 - \varphi J_m , \quad (3.1)$$

where now J_m is the Lorentz-invariant mass ‘density’. For a system of free particles we have

$$J_m = \sum_p \int d\tau_p m_p \delta^{(4)}(x - x_p(\tau_p)) . \quad (3.2)$$

¹In fact, it can be argued that Einstein and his friend Marcel Grossmann were agonizingly close to the answer as early as 1912.

In fact, as shown in the discussion after (2.61), such density is essentially the trace of the energy-momentum tensor: $J_m = -T_a^a$. This means that such scalar gravity field, while leading to attractive forces, fails to couple to electromagnetic fields and for a fluid of density ρ and pressure p , it couples to the net combination $\rho - 3p$, which vanishes for ultrarelativistic fluids (radiation). It is also in disagreement with experiment, since it predicts no bending of light in a gravitational field and a wrong precession effect for planetary orbits. Nevertheless, the scalar gravity theory is logically consistent, and was indeed seriously considered by many authors, including the likes of Abraham, Nordström and Einstein. Today, it survives in the so-called Jordan–Brans–Dicke form as a possible extension of General Relativity, albeit significantly constrained by precision experimental tests.

If we insist that gravity should couple to energy in the most natural way, we should use the energy-momentum tensor in the role of ‘matter current’, since it is T_{ab} the object that encapsulates the local conservation of energy in SR. In this way, one is naturally postulating a gravitational potential of tensor nature, i.e. a minimal coupling of the form

$$S_{\text{int}} \sim \int d^4x \, h_{ab} T^{ab} , \quad (3.3)$$

between T_{ab} and a symmetric rank-two gravitational potential h_{ab} . The local conservation of the energy-momentum tensor, $\partial_a T^{ab} = 0$, implies that this Lagrangian interaction is ‘gauge invariant’ under the transformations

$$h_{ab} \rightarrow h_{ab} + \partial_a f_b + \partial_b f_a , \quad (3.4)$$

with $f_a(x)$ an arbitrary four-vector function. Hence, we are led to the development of a symmetric two-index generalization of Maxwell’s theory, the Fierz–Pauli theory (2.114), coupled to the matter energy-momentum tensor through (3.3).

However, such a program quickly becomes treacherous. The reason is that any field coupling to energy must necessarily couple to itself, resulting in a nonlinear theory. The gist of the argument is as follows. The Fierz–Pauli Lagrangian contributes a term to the total energy-momentum tensor quadratic in the h -field, i.e. $T_{(h)} \sim (\partial h)(\partial h)$, suppressing the detailed index structure. Since h_{ab} must couple to all forms of energy, including the energy contained in the h -field itself, we must write (3.3) in the ‘improved’ form

$$S_{\text{int}} \sim \int h_{ab} \left(T_{\text{matter}}^{ab} + T_{(h)}^{ab} \right) ,$$

which induces a cubic interaction term in the h_{ab} field with the structure $h \, \partial h \, \partial h$. In turn, taking this term into account in the computation of the h -field energy momentum tensor, we will obtain a similar cubic term in $T_{(h)}^{ab}$, which induces a quartic term when fed into the interaction term, and so on... Hence, we are led to an iterative construction of a full non-linear theory, with arbitrarily large powers of h_{ab} , albeit with only two derivatives. Feynman and others were able to show that this iterative construction indeed converges to Einstein’s theory of General Relativity. We shall return to this issue later on, but for the time being we will be following another route.

3.1 From Galileo to Einstein

Einstein chose to develop the theory of gravity on top of the Principle of Equivalence between gravitation and inertia. A principle (going back to the tales of Galileo at the Pisa tower) stating,

in its weakest version, that the gravitational ‘charge’ of particles is universally proportional to their inertial mass.

When a Lagrangian shows some constraint in its parameters to a great accuracy (recall Eötvös measurements), it is natural to try representing this constraint as the result of some deeper fact. In the case at hand, this is possible in a rather interesting way. The Newtonian equation of motion for a particle of mass m on some gravitational potential is

$$m \frac{d\vec{v}}{dt} = -m \vec{\partial} \phi_N , \quad (3.5)$$

and the Principle of Equivalence simply means that m drops from this equation, so that trajectories are intrinsically determined by initial conditions, independently of the particular mass considered. This suggests that these trajectories have an *intrinsic* geometrical meaning. Consider, for example, a constant field with uniform gravity acceleration $\vec{g}$. The equation of motion of a falling particle is

$$\frac{d^2 \vec{x}}{dt^2} = \vec{g} . \quad (3.6)$$

Now, in a reference frame in free fall,

$$\vec{x}_{\text{ff}} = \vec{x} - \frac{1}{2} \vec{g} t^2 \quad (3.7)$$

we have $d^2 \vec{x}_{\text{ff}} / dt^2 = 0$, recovering the familiar feature that the gravitational field disappears in free fall. However, this was ultimately possible due to the particle mass dropping from the equations. Hence, we can represent the Principle of Equivalence as a statement about the change of reference frames, by saying that all systems in free fall are inertial, in the sense that they feel no gravitational forces.

Conversely, a frame of reference under constant acceleration of magnitude $\vec{g}$ is indistinguishable from a rest frame immersed in a constant gravitational field of the same intensity. In general, this simulation of gravitational fields by accelerations is only possible locally. We shall refer to such gravitational fields that can disappear globally by entering free fall as ‘fictitious’ gravitational fields. In ‘true’ ones, one can still eliminate the gravitational field, but only locally. If the gravitational acceleration at the spacetime point P is given by $\vec{g}_P = -\vec{\partial} \phi_N(P)$, the frame transformation $\vec{x}_{\text{ff}} = \vec{x} - \frac{1}{2} \vec{g}_P t^2$ gives

$$\frac{d^2 \vec{x}_{\text{ff}}}{dt^2} = \vec{g}(\vec{x}, t) - \vec{g}_P , \quad (3.8)$$

so that the acceleration vanishes at the point P . Away from the free-fall point P , there is a residual ‘tidal’ acceleration, proportional to the first derivatives of the gravitational force, or the second derivative of the potential. We can define the ‘tidal tensor’ for static fields as

$$\mathcal{R}_{ij} \equiv -\partial_i \partial_j \phi_N , \quad (3.9)$$

which provides a local diagnostic of whether a gravitational field is ‘true’ or ‘fictitious’. In particular, when the tidal tensor of a *static* field vanishes everywhere, the potential is linear in the spatial coordinates, $\phi_N(\vec{x}) = a + \vec{b} \cdot \vec{x}$. In this case the gravitational force is $\vec{g}(\vec{x}) = -\vec{b}$, a uniform field, which may be removed by the free-fall frame $\vec{x}_{\text{ff}} = \vec{x} + \frac{1}{2} \vec{b} t^2$.

In the relativistic theory, the Principle of Equivalence admits the same formulation: *gravitational fields can be locally eliminated by entering free fall*, except that this time it is Special

Relativity (SR), rather than Newtonian particle mechanics, that is assumed to be valid at the freely falling frames. The Principle of Equivalence is sometimes qualified as ‘weak’ or ‘strong’ depending on the scope of its application. The weak version only refers to the mass of a probe particle dropping from the equation of motion. Stronger versions differ by the generality of phenomena becoming ‘blind’ to gravitation in free fall, from purely mechanical to electromagnetic, gravitational self-energy effects or even quantum effects. Experimental tests of the Principle of Equivalence show no deviation from the strongest versions, so we will not make distinctions unless otherwise stated and speak of *the* Principle of Equivalence.

Since the Principle of Equivalence must be applied locally in generic gravitational fields, there is a degree of vagueness as to the size of the region where free-fall conditions apply. We shall give a more mathematical definition of the principle in coming sections, but for the time being we will continue exploring fictitious gravitational fields in order to gain some intuition in the relativistic regime.

3.1.1 Relativistic Inertial Fields

There are at least two novel relativistic effects with far-reaching consequences in relation to the Principle of Equivalence. The first is the existence of a novel kind of time dilation effects, associated to gravitational fields, and second is the occurrence of non-Euclidean geometry. Both effects can be illustrated in a simple heuristic way by studying appropriate fictitious gravitational fields.

The Relativistic Elevator and Rindler Spacetime

We saw that, according to the Principle of Equivalence, uniform and constant gravitational fields can be manufactured by constructing an elevator undergoing constant acceleration. In the relativistic theory, such an elevator would reach eventually the speed of light, so it is not obvious what is the relativistic concept associated to ‘constant acceleration’. Given a timelike trajectory $x(\tau)$, with four-velocity

$$u^a = \frac{dx^a}{d\tau} , \quad u^2 = -1 , \quad (3.10)$$

we define the acceleration four-vector by

$$a^b = \frac{du^b}{d\tau} , \quad a^2 = a_b a^b . \quad (3.11)$$

Taking the derivative of $u^2 = -1$ we find that acceleration and velocity are orthogonal $a^b u_b = 0$. In an inertial frame x'^a momentarily at rest with respect to the accelerated trajectory we have $u = (1, \vec{0})$ and the orthogonality implies $a = (0, \vec{a}')$, $\vec{a}' \cdot \vec{a}' = a^2$. Hence, we shall define constant acceleration as $a^2 = g^2$ with g a pure number, constant in time.

Now we go back to the ‘laboratory’ frame and take an accelerated rectilinear motion in the z -direction, $\vec{a} = d^2\vec{x}/d\tau^2 = (0, 0, a^z)$. We take $a^z > 0$ by convention. From the three algebraic equations

$$u^2 = -1 , \quad u \cdot a = 0 , \quad a^2 = g^2$$

we derive

$$a^0 = \frac{du^0}{d\tau} = g u^z , \quad a^z = \frac{du^z}{d\tau} = g u^0 . \quad (3.12)$$




Figure 3.1: The so-called Rindler wedge is the region $|t| < z$. Uniformly accelerated observers follow the $\xi = \text{const}$ hyperbolic trajectories, which degenerate into the past and future particle horizons $\mathcal{H}^\pm$. We also show the timelike trajectory of another observer. The part of her history to the future of $\mathcal{H}^+$ is completely inaccessible to the accelerated observer, whereas her crossing of the horizon takes an infinite subjective time for him.

The solution of these ordinary differential equations is

$$(u^0 \pm u^z) = (u^0 \pm u^z)(0) e^{\pm g\tau} .$$

Choosing now $u^0(0) = 1$ and $u^z(0) = 0$ (zero velocity at $\tau = 0$) we obtain the parametric form of the trajectory,

$$t - t(0) = \frac{1}{g} \sinh(g\tau) , \quad z - z(0) = \frac{1}{g} (\cosh(g\tau) - 1) , \quad (3.13)$$

and choosing the origin of coordinates so that $t(0) = 0, z(0) = 1/g$ we see that ‘constantly accelerated’ trajectories lie on the hyperbolae

$$z^2 - t^2 = \frac{1}{g^2} . \quad (3.14)$$

Now consider a family of accelerated observers running on hyperbolae $\xi = \text{constant}$, according to the change of coordinates

$$z \pm t = \frac{1}{g} e^{g\xi} e^{\pm g\eta} . \quad (3.15)$$

Each such observer feels a constant acceleration $g_\xi = g e^{-g\xi}$. If we arrange all of them to pass through the $t = \eta = 0$ line simultaneously, then they will always share the same value of the η coordinate at all times. Any such line is related to the $t = 0$ line by a Lorentz transformation, so that the relative rest condition of the accelerated observers is maintained in time. Hence, the constructed family of accelerated observers furnishes a ‘relativistic elevator’.

The space (η, ξ) with arbitrary real values of the coordinates is called *Rindler space*. It covers the wedge $|t| < z$ of Minkowski space, and represents the part of it accessible to accelerated

observers moving with positive uniform acceleration in the z -direction. Locally, Rindler space has identical geometry to Minkowski space, although the metric in Rindler coordinates reads

$$ds^2 = -dt^2 + dz^2 + dx^2 + dy^2 = e^{2g\xi}(-d\eta^2 + d\xi^2) + dx^2 + dy^2. \quad (3.16)$$

The future boundary of the Rindler wedge $\mathcal{H}^+$ is called the future *horizon*. Beyond this horizon, we find all the events that are causally disconnected from any accelerated observer in Rindler space. An observer at rest at $z > 1/g$ will eventually be passed by our accelerated elevator and by the time her worldline passes through the curve $z = t$, any subsequent signal that she might emit will never reach the elevator. At the same time, the observer on the elevator will never see her cross the horizon, her motions being constantly elongated in time by an infinite time dilation factor.

There is an analogous notion of ‘past horizon’, $\mathcal{H}^-$, which is defined by the time-reversal of the future horizon. Such notions of causal horizons feature prominently in the theory of black holes and also on the global properties of cosmological spacetimes.

The proper time of each observer at fixed ξ can be computed from the relation (3.13)

$$t(\tau_\xi) = \frac{1}{g} e^{g\xi} \sinh(g\eta(\tau_\xi)) = \frac{1}{g_\xi} \sinh(g_\xi \tau_\xi)$$

which results in $g\eta(\tau_\xi) = g_\xi \tau_\xi$, or

$$\tau_\xi = \eta e^{g\xi}. \quad (3.17)$$

Using this result, we find the ratio of proper times for observers at different heights in the elevator to be

$$\frac{\Delta\tau(\xi)}{\Delta\tau(\xi')} = e^{g(\xi-\xi')} = \frac{g_{\xi'}}{g_\xi}. \quad (3.18)$$

According to the Principle of Equivalence, this means that clocks run slower the more intense the gravitational field. This effect is one of the classic predictions of GR, called ‘gravitational red shift’. We can derive a local version of this important fact by considering two clocks ‘at rest’ in the elevator frame (that is to say, at equal values of η), and small proper distance apart in ‘elevator height’. Putting them at the same location in the transverse (x, y) plane, their height positions are ξ and $\xi' = \xi - \Delta\xi$. To leading order in $\Delta\xi$ their proper distance is then $h = \Delta s \approx e^{g\xi} \Delta\xi$. Applying (3.18) to leading order in the height, h , we find for the relative period shift of clocks

$$\frac{\Delta T}{T} = \frac{\Delta\tau_\xi - \Delta\tau_{\xi-\Delta\xi}}{\Delta\tau_\xi} \approx g_\xi h. \quad (3.19)$$

A local observer sitting in the elevator interprets g_ξ as the intensity (acceleration) of the gravitational field at point ξ .

Problem: More about the Twins

Consider the proverbial Langevin’s twins. They are separated at birth, one remaining on Earth and the other one embarking on a spaceship to planet X. The spaceship accelerates all the time at the Earth-friendly rate of one g . The trip consists of an accelerating phase, a decelerating phase and a similar procedure on the way back from X to Earth. The traveling twin arrives on Earth right on time to his

brother's 90th birthday party. What is the distance to planet X and how old is the traveling twin at the party?

Rotation and Non-Euclidean Geometry

Consider two frames of reference in SR. K is inertial and K' rotates at angular velocity Ω . Imagine setting up an operational measure of π , i.e. we draw a circumference of radius R and measure its length. The ratio being 2π in the inertial system K . An observer on K' would use measuring rods that, according to K , suffer Lorentz contraction when laid along the circumference, but remain intact when measuring the radius. Hence, the observer on K' will measure a larger ratio of circumference to diameter, by a factor of the Lorentz contraction $\sqrt{1-v^2} = \sqrt{1-\Omega^2 R^2}$. This leads to a 'measurement' of π as

$$\pi_{K'} = \frac{\pi}{\sqrt{1-\Omega^2 R^2}} > \pi. \quad (3.20)$$

This is characteristic of spaces of *negative* curvature. According to the Principle of Equivalence, there is no local difference between accelerated frames and true gravitational fields, the two differing only at the level of global properties. Hence we should expect non-Euclidean geometry to arise in true gravitational fields as a general rule. This simple *gedanken experiment* was pivotal in Einstein's own heuristic path to GR.

Incidentally, we can also use this example of the rotating frame to give a derivation of the gravitational time delay. Distributing clocks at different radial positions, their rhythm will be proportional to $\sqrt{1-v^2} = \sqrt{1-\Omega^2 R^2}$ as witnessed by the observers on K . Observers on K' will notice this delay and, seeing the clocks at rest, will interpret it as a consequence of the local centrifugal 'gravitational' field. The larger the centrifugal field, the slower the clock rate.

Problem: Rotating Clock Delay

Given two clocks in the rotating frame at radii r and $r-h$, with small h , show that the relative period shift to leading order in h is given by

$$|\Delta T/T| = g(r) h$$

where $g(r)$ is the local centrifugal acceleration measured at radius r . Check that $g(r)$ is given by the Newtonian formula $g(r) = \Omega^2 r$ in the non-relativistic approximation.

3.1.2 A Tale of Two Frames: a First Look at General Covariance

We have seen that, according to the Principle of Equivalence, we can regard any field of inertial forces, induced by a change of frame, as a 'fictitious' gravitational field. It is fictitious in the sense that we can make it disappear by simply changing the frame back to the inertial one, which looks to be in free fall from the point of view of the non-inertial frame. In other words, we are just describing Lorentz-invariant physics of Minkowski spacetime, in some general coordinate frame, a spacetime generalization of the well-known fact that we may describe Newtonian physics in special 'curvilinear' coordinates such as spherical, accelerated or rotating coordinates.

It will be useful to develop some formalism about such fictitious fields, in a manner that is as coordinate-independent as possible. The idea is to generalize the two cases of accelerated/rotating observers studied in the last two sections. We will refer to the inertial frame with Minkowskian coordinates x^a as the *frefo* frame, indicating that it is in ‘free fall’, in the sense of the Principle of Equivalence. We will also consider a frame y^α of curvilinear and/or accelerated coordinates, called the *fido* frame, for ‘fiducial observer’, referring to the general situation exemplified by the families of observers in uniform acceleration or rotation in the previous subsections. It could be that the *fido* frame has a physical interpretation in terms of standard observers, like the cases above, but in general we will interpret the y^α frame in purely mathematical terms, just as we do for polar spherical coordinates for instance. We will only ask for smoothness of the transformation $x^a \rightarrow y^\alpha$.

Differentiating the functions $x^a(y)$ we obtain the expression of the interval in arbitrary coordinates:

$$ds^2 = \eta_{ab} dx^a dx^b = g_{\alpha\beta} dy^\alpha dy^\beta, \quad (3.21)$$

where the 4×4 symmetric matrix of functions

$$g_{\alpha\beta}(y) \equiv \eta_{ab} \frac{\partial x^a}{\partial y^\alpha} \frac{\partial x^b}{\partial y^\beta} \quad (3.22)$$

is the metric transformed to the new coordinates. If we want to relate the metric in different choices of generalized coordinates we just apply the chain rule of differentiation to obtain

$${}^{(z)}g_{\alpha\beta}(z) = {}^{(y)}g_{\gamma\delta}(y) \frac{\partial y^\gamma}{\partial z^\alpha} \frac{\partial y^\delta}{\partial z^\beta}. \quad (3.23)$$

The transformation matrices $\partial y^\alpha / \partial z^\beta$ are general invertible matrices in $GL(4, \mathbb{R})$, with smooth dependence on the points of spacetime, for any smooth invertible map $z \rightarrow y(z)$, a so-called Diffeomorphism (Diff).² We see that the metric functions $g_{\alpha\beta}$ transform like a covariant tensor of rank two under this Diff group. The transformation rules are formally the same that we had for Lorentz tensors, with the $O(1, 3)$ matrices replaced with the $GL(4)$ matrices, point by point. Notice however that, while the η_{ab} were Lorentz-invariant, the $g_{\alpha\beta}$ are only Diff-covariant, but *both* are valid representations of the Minkowski metric, because they compute the same interval (3.21).

Diff Tensor Calculus

Any Lorentz-covariant equation in SR, written in terms of Lorentz tensors, can be written as a Diff-covariant equation by passing to Diff-covariant tensors. To do this, focus on the transformation matrices between a *frefo* and a *fido*:

$$E_\alpha^a(y) \equiv \frac{\partial x^a}{\partial y^\alpha}(y). \quad (3.24)$$

The matrix of functions $E_\alpha^a(y)$ transform as contravariant Lorentz-vectors in the *frefo* index and as covariant Diff-vectors in the *fido* index. One can consider quantities related to these by an operation of raising and lowering of indices, with the convention that *frefo* indices are raised and

²We will always consider in these notes the ‘small’ Diff group, namely smooth maps continuously deformable to the identity. In particular, all these maps are orientation-preserving.

lowered with the Lorentz metric and the *fido* indices are lowered with the $g_{\alpha\beta}$ metric and raised with its inverse $g^{\alpha\beta} \equiv (g^{-1})^{\alpha\beta}$:

$$E_{a\alpha} = \eta_{ab} E_{\alpha}^b, \quad E^{a\alpha} = g^{\alpha\beta} E_{\beta}^a, \quad E_a^{\alpha} = \eta_{ab} g^{\alpha\beta} E_{\beta}^b, \quad (3.25)$$

transform as indicated by the position of the indices. Our definition of covariant vs contravariant transformations for Diff are the same we adopted for Lorentz, namely we say that $\partial_{\alpha} = \partial/\partial y^{\alpha}$ transforms *covariantly* in the index α , whereas dy^{α} transforms *contravariantly* in the index α , when these transformations are effected by applying the chain rule of differentiation:

$$\frac{\partial}{\partial y^{\alpha}} \rightarrow \frac{\partial z^{\beta}}{\partial y^{\alpha}} \frac{\partial}{\partial z^{\beta}}, \quad dy^{\alpha} \rightarrow \frac{\partial y^{\alpha}}{\partial z^{\beta}} dz^{\beta}. \quad (3.26)$$

Then, given any Lorentz tensor of components $T_{\dots b \dots}^{\dots a \dots}(x)$, the related set of quantities

$$T_{\dots \beta \dots}^{\dots \alpha \dots}(y) \equiv T_{\dots b \dots}^{\dots a \dots}(x) \cdots E_a^{\alpha}(y) \cdots E_{\beta}^b(y) \cdots, \quad (3.27)$$

contains the same information as the $T_{\dots b \dots}^{\dots a \dots}(x)$, once the diffeomorphism $x \rightarrow y(x)$ is known. This expression defines the Diff-tensor version of the formal object

$$\mathbf{T} = T_{\dots \beta \dots}^{\dots \alpha \dots}(y) \cdots \otimes \frac{\partial}{\partial y^{\alpha}} \otimes \cdots \otimes dy^{\beta} \otimes \cdots \quad (3.28)$$

which is Diff-invariant.

The definition of the metric itself (3.22) is a particular case of the general rule (3.27) to construct Diff tensors out of Lorentz tensors. As we mentioned above, the metric loses its ‘invariant-tensor’ character when generalized to the Diff group. On the other hand, the Kronecker delta δ^a_b retains its invariance property, since its Diff version $\delta^{\alpha}_{\beta} = E_a^{\alpha} E_{\beta}^b \delta_b^a$ remains a Kronecker delta. The situation with the other classic invariant tensor, the Levi-Civita tensor ϵ_{abcd} is slightly different. Recall our conventions: $\epsilon_{abcd} = [abcd]$, the permutation symbol with $[0123] = 1$. Then, we notice that its Diff-covariantization

$$\epsilon_{abcd} E_{\alpha}^a E_{\beta}^b E_{\gamma}^c E_{\delta}^d \quad (3.29)$$

is completely antisymmetric in the indices $\alpha, \beta, \gamma, \delta$. We may then write

$$\epsilon_{abcd} E_{\alpha}^a E_{\beta}^b E_{\gamma}^c E_{\delta}^d = [\alpha\beta\gamma\delta] \epsilon_{abcd} E_0^a E_1^b E_2^c E_3^d = [\alpha\beta\gamma\delta] \det(E),$$

where $\det(E)$ is a shorthand for the determinant of the matrix of functions E^a_{β} . Since the transformation $x^a \rightarrow y^{\alpha}$ is invertible, $\det(E)$ cannot vanish. In fact, its sign is positive if the *frefo* $\rightarrow$ *fido* map preserves orientation, and negative otherwise. Assuming that orientation is indeed preserved, we have $\det(E) > 0$. We will also adopt the following shorthand notation for the determinant of the metric matrix of functions:

$$g \equiv \det(g_{\alpha\beta}). \quad (3.30)$$

Then (3.22) implies that $g = \det(E)^2 \det(\eta) = -|\det(E)|^2$, so that we can write $\det(E) = (-g)^{1/2}$ and

$$\epsilon_{\alpha\beta\gamma\delta} \equiv \epsilon_{abcd} E_{\alpha}^a E_{\beta}^b E_{\gamma}^c E_{\delta}^d = \sqrt{-g} [\alpha\beta\gamma\delta] \quad (3.31)$$

defines a rank-four covariant Diff-tensor. Analogously, we can define the contravariant Levi-Civita tensor $\epsilon^{\alpha\beta\gamma\delta}$ by the usual raising of indices with the inverse metric $g^{\alpha\beta}$, and obtain an associated rank-four contravariant Diff-tensor

$$\epsilon^{\alpha\beta\gamma\delta} = -\frac{1}{\sqrt{-g}} [\alpha\beta\gamma\delta] . \quad (3.32)$$

In general, quantities that are tensors up to a power of the Jacobian of the transformation are called tensor densities. The object $\mathsf{T}^{(w) \dots \alpha \dots}_{\dots \beta \dots}$ is called *tensor density* of weight w when it transforms as

$$\mathsf{T}'^{(w) \dots \mu \dots}_{\dots \nu \dots}(x') = \left| \frac{\partial x}{\partial x'} \right|^w \dots \frac{\partial x'^\mu}{\partial x^\alpha} \dots \frac{\partial x^\beta}{\partial x'^\nu} \dots \mathsf{T}^{(w) \dots \alpha \dots}_{\dots \beta \dots}(x) . \quad (3.33)$$

In addition to the permutation symbol, one notable example is the delta function $\delta^{(4)}(x - x_0)$, which transforms as a scalar density of weight one. In particular $(-g)^{-1/2} \delta^{(4)}(x - x_0)$ is a true scalar.

The Covariant Derivative

So far the construction of Diff tensors out of Lorentz tensors was quite straightforward. The next slightly non-trivial step involves defining a Diff-tensor version of the derivative operator. When acting on Diff-scalar functions, the ordinary *fido* derivative $\partial_\alpha \phi$ is related to the *frefo* derivative $\partial_a \phi$ by the standard formula (3.27), namely

$$\partial_\alpha \phi = E^a{}_\alpha \partial_a \phi .$$

It is interesting that this *does not* generalize to derivative operators acting on higher-rank Diff-tensors. In particular, the object $\partial_\alpha V_\beta$ is not a Diff-tensor, even if we define V_β as a covariant Diff-vector. The trouble arises when the derivative acts on the transformation matrices, which now are general functions of position. On the other hand, the general rule (3.27) does produce a Diff-tensor out of the Lorentz tensor $\partial_a V_b$, namely the object

$$\nabla_\alpha V_\beta \equiv E^a{}_\alpha E^b{}_\beta \partial_a V_b \quad (3.34)$$

is a rank-two covariant Diff-tensor by construction. By direct computation, we can easily find that

$$\partial_\alpha V_\beta = \partial_\alpha (E^a{}_\beta V_a) = (\partial_\alpha E^a{}_\beta) V_a + E^b{}_\beta \partial_\alpha V_b = E_a{}^\delta \partial_\alpha E^a{}_\beta V_\delta + E^a{}_\alpha E^b{}_\beta \partial_a V_b$$

Therefore, we can write the so-called *covariant derivative* acting on covariant Diff-vectors as

$$\nabla_\alpha V_\beta = \partial_\alpha V_\beta - \Gamma_{\alpha\beta}^\delta V_\delta , \quad (3.35)$$

where the coefficients

$$\Gamma_{\alpha\beta}^\delta \equiv (\partial_\alpha E^a{}_\beta) E_a{}^\delta = \frac{\partial^2 x^a}{\partial y^\alpha \partial y^\beta} E_a{}^\delta \quad (3.36)$$

depending on second derivatives of the *frefo-fido* mapping, are usually referred to as Christoffel symbols, or affine connection coefficients. They provide the correction term that is needed to write a covariant version of the derivative operator.

Straight from the definition (3.36) one can prove that the Christoffel symbols can be entirely expressed as linear combinations of the first derivatives of the metric:

$$\Gamma_{\alpha\beta}^{\delta} = \frac{1}{2} g^{\delta\gamma} (\partial_{\alpha} g_{\gamma\beta} + \partial_{\beta} g_{\alpha\gamma} - \partial_{\gamma} g_{\alpha\beta}) . \quad (3.37)$$

It is very important to remember that, despite the notation, the Christoffel symbols *do not* define a Diff tensor. This is just natural, since their purpose is to cancel the non-tensor components of the ordinary derivative when acting on Diff-tensors. Indeed, a direct calculation from either (3.36) or (3.37) shows that, under a diffeomorphism $y \rightarrow z$, the connection coefficients transform as

$${}^{(y)}\Gamma_{\alpha\beta}^{\delta} = {}^{(z)}\Gamma_{\gamma\sigma}^{\rho} \frac{\partial z^{\gamma}}{\partial y^{\alpha}} \frac{\partial y^{\delta}}{\partial z^{\rho}} \frac{\partial z^{\sigma}}{\partial y^{\beta}} - \frac{\partial z^{\rho}}{\partial y^{\alpha}} \frac{\partial z^{\sigma}}{\partial y^{\beta}} \frac{\partial^2 y^{\delta}}{\partial z^{\rho} \partial z^{\sigma}} . \quad (3.38)$$

There is a tensorial part and an inhomogeneous part depending on second derivatives of the diffeomorphism. This non-tensorial character is responsible for the fact that the Christoffel symbols can vanish in a particular frame, such as the *frefo*, and not vanish in a generic *fido*.

Entirely similar manipulations allow us to define covariant derivatives acting on contravariant Diff-vectors, the result differing from (3.35) only by a sign in the correction term:

$$\nabla_{\alpha} V^{\beta} \equiv E^a{}_{\alpha} E_b{}^{\beta} \partial_a V^b = \partial_{\alpha} V^{\beta} + \Gamma_{\alpha\delta}^{\beta} V^{\delta} . \quad (3.39)$$

Upon further generalization along the same lines, we obtain the general rule for the covariant derivative of an arbitrary Diff tensor:

$$\nabla_{\delta} T^{\dots\alpha\dots}_{\dots\beta\dots} = \partial_{\delta} T^{\dots\alpha\dots}_{\dots\beta\dots} + \dots + \Gamma_{\delta\gamma}^{\alpha} T^{\dots\gamma\dots}_{\dots\beta\dots} + \dots - \Gamma_{\delta\beta}^{\gamma} T^{\dots\alpha\dots}_{\dots\gamma\dots} - \dots . \quad (3.40)$$

It is not difficult to check that these definitions give a covariant derivative operator ∇_{α} satisfying Leibniz's rule, namely $\nabla(T \otimes S) = (\nabla T) \otimes S + T \otimes (\nabla S)$ for any tensors T and S .

Using these expressions, one can derive a number of useful identities. The first and most important is the so-called *metric compatibility* of the covariant derivative, which simply means that the covariant derivative of the metric vanishes identically.

$$\nabla_{\delta} g_{\alpha\beta} = 0 , \quad (3.41)$$

a property implying that the covariant derivative commutes with the operation of raising and lowering Diff-indices.

A second important property is that the Christoffel connection drops from the antisymmetric derivatives. Given an antisymmetric tensor $A_{\alpha\beta\dots} = A_{[\alpha\beta\dots]}$, the antisymmetric derivative satisfies

$$\nabla_{[\alpha} A_{\beta\dots]} = \partial_{[\alpha} A_{\beta\dots]} . \quad (3.42)$$

This follows technically from the symmetry of the connection in lower indices, $\Gamma_{\alpha}{}^{\gamma}{}_{\beta} = \Gamma_{\beta}{}^{\gamma}{}_{\alpha}$. A related identity applies to the contracted derivative, again acting on antisymmetric tensors:

$$\nabla_{\alpha} A^{[\alpha\dots]} = \frac{1}{\sqrt{-g}} \partial_{\alpha} \left(\sqrt{-g} A^{[\alpha\dots]} \right) . \quad (3.43)$$

These two relations allow us to generalize the exterior calculus on p -forms (the machinery associated to the exterior derivative operator d and its adjoint $d^{\dagger}$) in a Diff-covariant manner.

Together with the Diff versions of the Levi–Civita tensors, we can also define the Diff-covariant version of Poincaré duality and all the theory of integral invariants.

A particular case of (3.43) is the very useful expression for the Laplacian operator acting on a scalar function:

$$\nabla^2 \phi = \nabla_\alpha \nabla^\alpha \phi = \frac{1}{\sqrt{-g}} \partial_\alpha \left(\sqrt{-g} g^{\alpha\beta} \partial_\beta \right) \phi . \quad (3.44)$$

Another important particular case is that of the covariant divergence of a vector field J^μ , which appears in the Diff-covariant version of the continuity equation $\nabla_\mu J^\mu = 0$. In this case we can write

$$\nabla_\mu J^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} J^\mu) . \quad (3.45)$$

This form can be used to prove the Diff-covariant version of Gauss’ integration theorem for a total derivative over a four-dimensional region with boundary:

$$\int_{V^{1+3}} \nabla_\mu J^\mu = \int_{\partial V^{1+3}} n_\mu J^\mu , \quad (3.46)$$

where n^μ with $n_\mu n^\mu = \pm 1$ is a normal unit vector to ∂V^{1+3} with the convention that it is outward pointing when n^μ is spacelike and inward pointing when n^μ is timelike.

Another handy device is the ability to compute a derivative of a tensor along a curve γ , parametrized by functions $x^a(\sigma)$ or $y^\alpha(\sigma)$. In the *frefo* we have

$$\frac{d}{d\sigma} T^{\dots a \dots}_{\dots b \dots} = \frac{dx^c}{d\sigma} \partial_c T^{\dots a \dots}_{\dots b \dots} .$$

Now the expression on the right hand side can be Diff-covariantized into

$$\frac{\nabla}{d\sigma} T^{\dots \alpha \dots}_{\dots \beta \dots} = \frac{dy^\delta}{d\sigma} \nabla_\delta T^{\dots \alpha \dots}_{\dots \beta \dots} \quad (3.47)$$

where we have introduced a new notation, $\nabla/d\sigma$, for the notion of ‘covariant derivative along the parametrized curve’. Since the tangent vector γ_σ to the parametrized curve γ , with components $dy^\delta/d\sigma$, is a Diff-vector, we can also write this as a ‘directional covariant derivative’ in the direction of the tangent vector,

$$\frac{\nabla}{d\sigma} T^{\dots \alpha \dots}_{\dots \beta \dots} = \nabla_{\gamma_\sigma} T^{\dots \alpha \dots}_{\dots \beta \dots} \quad (3.48)$$

where

$$\nabla_V = V^\alpha \nabla_\alpha \quad (3.49)$$

is the directional covariant derivative operator along the vector V .

An example where we can put these definitions to work is the equation of motion for a free massive particle in SR. In a *frefo* frame it reads

$$\frac{d^2 x^a}{d\tau^2} = \frac{du^a}{d\tau} = 0 , \quad (3.50)$$

where τ is the proper time along the parametrized spacetime path $x^a(\tau)$ of the particle and $u^a = dx^a/d\tau$ is the four-velocity. In a *fido* frame the path is described by functions $y^\alpha(\tau)$ and the equation of motion can be written in a variety of forms:

$$\frac{\nabla^2 y^\alpha}{d\tau^2} = \frac{\nabla u^\alpha}{d\tau} = u^\beta \nabla_\beta u^\alpha = \nabla_u u^\alpha = 0 , \quad (3.51)$$

or, in full component glory:

$$\frac{d^2 y^\alpha}{d\tau^2} + \Gamma_{\beta\delta}^\alpha \frac{dy^\beta}{d\tau} \frac{dy^\delta}{d\tau} = 0 . \quad (3.52)$$

Another example is Maxwell's equation, $\partial_a F^{ab} = -J^b$, which can be covariantized into

$$\nabla_\alpha F^{\alpha\beta} = -J^\beta . \quad (3.53)$$

A further useful presentation of the same equation is

$$\frac{1}{\sqrt{-g}} \partial_\alpha (\sqrt{-g} F^{\alpha\beta}) = -J^\beta . \quad (3.54)$$

Physics, what Physics?

We have introduced in the previous sections a large part of the mathematical machinery that is used in the theory of General Relativity, yet one can say that the exercise is purely formal. Indeed, all we were doing is describing good old SR in a Diff-covariant way. We never left Minkowski spacetime, and all the Diff-machinery could evaporate immediately if we decided to use a *frefo* frame throughout $\mathbb{R}^{1+3}$.




Figure 3.2: An arbitrary ‘fictitious’ field is just Minkowski spacetime (left) disguised in a general system of curvilinear and/or accelerated coordinates (right).

The ability to express physical equations in terms of a general coordinate system, the Diff-covariant description outlined above, is called *general covariance* and it is by itself devoid of any physical content. Every equation can be made Diff-covariant without changing its physical content. If the equation is local, it can be written in terms of local quantities over spacetime and its derivatives. Then, the dependence on the arbitrary frame is for the most part encapsulated in the metric matrix $g_{\alpha\beta}$ and the Christoffel symbols $\Gamma_{\alpha\beta}^\delta$.

Let us take up the example of the particle equation of motion (3.52). In the *frefo* we have the standard $d^2 x^a / d\tau^2 = 0$ which is solved by straight lines $x^a(\tau) = x_0^a + u^a \tau$ with sub-luminal velocity. In the *fido*, the histories $y^\alpha(\tau)$ will not be straight lines in general of course, but they will be the images of the straight lines, under the *frefo* $\rightarrow$ *fido* map. Their generalized acceleration will be equal to a non-trivial force quadratic in the four-velocities which is nothing but the relativistic generalization of the good old Newtonian ‘inertial’ forces:

$$\frac{d^2 y^\alpha}{d\tau^2} = \frac{1}{m} F_{\text{inertial}}^\alpha = -\Gamma_{\delta\gamma}^\alpha \frac{dy^\gamma}{d\tau} \frac{dy^\delta}{d\tau} . \quad (3.55)$$

With regard to gravitation, the physical content of this discussion is just the *interpretation* of inertial forces as ‘gravitational’ in the sense that, locally there is no way to tell the difference. The reader may feel that the generally covariant formalism introduced in this section is a bit heavy handed and not quite worth the effort, but we will reap the benefits in the next section.

Problem: Relativistic Rotation

We discuss here relativistic inertial forces under rotation. Let (t, z, r, θ) be standard cylindrical coordinates in Minkowski spacetime, with metric

$$ds^2 = -dt^2 + dz^2 + dr^2 + r^2 d\theta^2 ,$$

and pass to a rotating frame (t, z, r, ϕ) with $\phi = \theta - \Omega t$. Find the metric in the rotating frame, compute the Christoffel symbols and write down the equation of motion for massive particles. Interpreting (3.55) as the relativistic Newton law with inertial forces, take the non-relativistic limit and find the expected centrifugal and Coriolis inertial forces.

3.2 Riemannian Geometry and the Principle of Equivalence

Having exploited the Principle of Equivalence for the particular case of fictitious gravitational fields, those that can be regarded as having purely inertial forces, we are now ready to consider the more general case in which the condition of ‘free fall’ can only be implemented locally around a point in spacetime.

Let us consider a spacetime X^{1+3} with a general gravitational field. We characterize the Principle of Equivalence in a pragmatic way as follows: Around any point $P \in X^{1+3}$, we can erect a *frefo* frame through P , denoted X_P^a , with the property that Special Relativity is locally valid in a small neighborhood of P , when described in this particular frame. The whole spacetime X^{1+3} can be covered by such *refos*. We can also refer all phenomena to an external frame, x^μ , the *fido*, to signify the fact that it may be associated to a fixed set of fiducial observers. Notice that no restrictions are placed on the properties of the *fido*, and no canonical choice of *refos* throughout spacetime exists in general, if only because any *frefo* can be rotated by a Lorentz transformation and remain a valid *frefo*, and this can be done independently at each point in spacetime.

3.2.1 Particle Probes

There is some uneasy vagueness to the previous statements; just how small must the *frefo* region around P be in order to claim that SR applies? In order to make more precise what is meant by X_P^a being ‘in free fall at P ’, we send a particle probe of mass m . Its spacetime path is parametrized in the *fido* by functions $x^\alpha(\sigma)$. We can partition the path as $\gamma = \cup_P \gamma_P$, with γ_P small ‘pieces’ around points P in the path, each of them parametrized in the *refos* by the functions $X_P^a(\sigma)$. According to the Principle of Equivalence, the equation of motion at point P , when referred to the *frefo* at P , takes the special-relativistic form:

$$\left. \frac{d^2 X_P^a}{d\tau^2} \right|_P = 0 . \quad (3.56)$$

But we can now use the Diff-covariant formalism that we developed for fictitious fields in the previous section, to map this equation into the *fido* frame. Defining the matrices of partial derivatives

$$E_P{}^a{}_\alpha(x) \equiv \frac{\partial X_P^a}{\partial x^\alpha}(x) , \quad (3.57)$$

we can carry over the formalism of Christoffel connections to this case, replacing the previous matrices of functions E_α^a by the new ones (3.57), referred to the local *frefo* at P .³ The result is

$$\left(\frac{d^2 x^\mu}{d\tau^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} \right) \Big|_P = 0 . \quad (3.58)$$

In this expression, the Christoffel symbols have the same definition as they had for fictitious fields:

$$\Gamma_{\alpha\beta}^\delta(P) = E_P{}^a{}_\delta(P) (\partial_\alpha E_P{}^a{}_\beta) \Big|_P = E_P{}^a{}_\delta(P) \frac{\partial^2 X_P^a}{\partial x^\alpha \partial x^\beta} \Big|_P , \quad (3.59)$$

and also the same expression (3.37) in terms of first derivatives of the metric. The metric itself in the *fido* frame is given by

$$g_{\alpha\beta}(P) = E_P{}^a{}_\alpha(P) E_P{}^b{}_\beta(P) \eta_{ab} , \quad (3.60)$$

and it computes the same proper time interval as the Minkowski metric, provided $d\tau$ is small enough to be well-approximated by a given *frefo*. The crucial difference with the case of fictitious fields is the absence of a global, unique *frefo*. Rather, in this case the metric can be computed using data from each *frefo* at P , but the *frefo* changes from point to point. In general, there will *not* exist a global *frefo* on which the metric takes the form η_{ab} throughout a finite region.

We conclude that the Principle of Equivalence, together with Special Relativity, just forces upon us the description of gravitation in terms of a Riemannian metric, namely a set of functions $g_{\mu\nu}$ which compute the proper time as

$$d\tau^2 = -g_{\mu\nu} dx^\mu dx^\nu , \quad (3.61)$$

and that transform as a rank-two covariant Diff-tensor under a smooth change of coordinates:

$$g_{\mu\nu}(x) \longrightarrow g'_{\mu\nu}(x') = \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu} g_{\alpha\beta}(x) . \quad (3.62)$$

All the formulas involving Diff-covariant quantities which we derived for fictitious fields, including the rules for covariant differentiation (3.40) and the metric compatibility property (3.41), *do* apply in a general gravitational field, provided we use as fundamental building block the general Riemannian metric (3.61).

We are now in position to reduce the vagueness of the previous discussion. The Equivalence Principle states that, through any point P in spacetime, there exists a *frefo* X_P^a , with the property that, in this frame, the metric at P reduces to the Minkowski form and the Christoffel symbols vanish at P . Equivalently, all first derivatives of the metric vanish at P :

$$g_{ab}(P) = \eta_{ab} , \quad \partial_c g_{ab}(P) = 0 = \Gamma_{bc}^a(P) . \quad (3.63)$$

³We emphasize that the functions (3.57) do extend throughout a neighborhood of P , but they are *different* functions for different anchoring points P .

Therefore, the ‘size’ of the *frefo* at P just depends on how far from P we may want to go, while ignoring the second order term in the Taylor expansion of the metric around P (controlled by the second derivatives of the metric). Fictitious fields are those for which a *frefo* exists such that these equations hold exactly in a finite region of spacetime, so that the metric can be exactly put in Minkowskian form throughout that region.




Figure 3.3: A particle path γ through X^{1+3} with two *frefos* at points P and Q and an external *fido* frame x^μ . Notice that the *frefos* may extend beyond the region where they are approximately in free fall. Beyond this region, they will lose the *frefo* character except for fictitious fields.

Covariant Particle Action

We can obtain the same results by focusing on the action of the probe particle, rather than its equations of motion. We shall reproduce this reasoning here because it gives an interesting perspective on the formalism and it serves as a blueprint for the ‘minimal coupling’ prescriptions that will be introduced later. Let us split the path γ in small segments, each of them passing through *frefos* X_P . On each segment, the Principle of Equivalence dictates the form of the action to be the same as in SR. Piecing together all such segments and taking the limit of a fine partition we obtain

$$S[\gamma] = \lim_{\{P\} \rightarrow \gamma} \sum_P \Delta S[\gamma_P], \quad (3.64)$$

where

$$\Delta S[\gamma_P] \approx -m \Delta \tau_P = -m \Delta \sigma_P \sqrt{-\eta_{ab} \frac{dX_P^a}{d\sigma} \frac{dX_P^b}{d\sigma}}, \quad (3.65)$$

which gives in the limit

$$S[\gamma] = -m \int_\gamma d\tau = -m \int_\gamma d\sigma \sqrt{-\eta_{ab} \frac{dX_{P(\sigma)}^a}{d\sigma} \frac{dX_{P(\sigma)}^b}{d\sigma}}. \quad (3.66)$$

Hence the Principle of Equivalence establishes the particle action as proportional to the elapsed proper time, just as in ordinary SR, and provides a way to compute it, given a family of *frefos* along γ . This parametrization serves the purpose of *defining* the action through the Principle of Equivalence, but it is not particularly ‘user-friendly’, since it contains a large amount of redundant information, related to the arbitrary choice of *frefos* at each point in γ . All this is simplified by passing to the *fido* description using

$$\frac{dX_P^a}{d\sigma} = \frac{\partial X_P^a}{\partial x^\alpha} \frac{dx^\alpha}{d\sigma} = E_P^a{}_\alpha \frac{dx^\alpha}{d\sigma} , \quad (3.67)$$

which allows us to write

$$\eta_{ab} \frac{dX_{P(\sigma)}^a}{d\sigma} \frac{dX_{P(\sigma)}^b}{d\sigma} = g_{\alpha\beta}(x(\sigma)) \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma} ,$$

and finally obtain the particle action in *fido* parametrization:

$$S[\gamma] = -m \int_\gamma d\tau = -m \int_\gamma d\sigma \sqrt{-g_{\alpha\beta}(x(\sigma)) \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma}} . \quad (3.68)$$

All the redundancy implicit in the choice of *frefo* at each point has been removed and the physical description of the particle dynamics is completely specified once the Riemannian metric $g_{\mu\nu}$ is known.

We can actually go further and derive the equation of motion stemming from (3.68) by demanding the action to be an extremum with respect to variations of the trajectory $x^\alpha(\sigma) \rightarrow x^\alpha(\sigma) + \delta x^\alpha(\sigma)$. We find

$$0 = \delta S_P = \int \frac{d\sigma}{2\sqrt{-g_{\alpha\beta} \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma}}} \delta \left(g_{\alpha\beta} \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma} \right) .$$

We may now take advantage of the reparametrization invariance to restore the proper time parameter $\sigma \rightarrow \tau$ to remove the square root in the denominator and write

$$0 = \int d\tau \left(\frac{1}{2} \partial_\mu g_{\alpha\beta} \dot{x}^\alpha \dot{x}^\beta \delta x^\mu + g_{\alpha\beta} \dot{x}^\alpha \frac{d}{d\tau} \delta x^\beta \right) .$$

Reshuffling the indices, integrating by parts and multiplying the resulting equation of motion by the inverse of the metric matrix we obtain the, by now familiar, equation of motion

$$0 = \frac{d^2 x^\alpha}{d\tau^2} + \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} , \quad (3.69)$$

where the Christoffel symbols appear directly in their expression in terms of the first derivatives of the metric:

$$\Gamma_{\mu\nu}^\alpha = \frac{1}{2} g^{\alpha\beta} (\partial_\mu g_{\beta\nu} + \partial_\nu g_{\beta\mu} - \partial_\beta g_{\mu\nu}) . \quad (3.70)$$

3.2.2 Gravitational Forces

Writing the free fall equation of motion (3.69) in morally Newtonian form

$$m \frac{d^2 x^\alpha}{d\tau^2} = F^\alpha = -m \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} , \quad (3.71)$$

we see that relativistic gravitational forces are velocity-dependent. We already noticed this fact when considering fictitious gravitational fields, otherwise known as ‘inertial forces’, as in (3.55). It is certainly not surprising that inertial forces should depend on velocities, because they certainly do in the Newtonian case (think of the formula for Coriolis forces). The remarkable fact is that the full equation of motion in an arbitrary gravitational field, does have exactly the same structure as the relativistic generalization of inertial forces. In hindsight, this is just natural, because the Principle of Equivalence is predicated on the identification of gravitational and inertial forces at a local level. Since the equation of motion is certainly local, this is the expected result.

Structurally, the forces are also proportional to the connection coefficients Γ , so that the metric functions acquire the interpretation of ‘gravitational potentials’, since their first derivatives determine the ‘forces’. We can make these intuitions more precise by taking the weak-field limit and further the Newtonian limit.

Weak Fields

In keeping with the idea that gravitational physics amounts to replacing the Minkowski metric η_{ab} by a general Riemannian metric $g_{\alpha\beta}$ we may study the case where gravitational effects are small, so that we write $g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta}$, with $|h_{\alpha\beta}| \ll 1$, a condition that requires restricting the reference frames, since we may induce large components of the metric by simply using a frame with large accelerations. Hence, we assume that an ‘almost’ *frefo* frame extends over the region of interest and we only allow further coordinate transformations that maintain the ‘small field’ property. Writing those transformations as $x^\alpha \rightarrow x'^\alpha = x^\alpha - \varepsilon^\alpha(x)$, the metric transformations (3.62) imply

$$\eta_{\alpha\beta} + h_{\alpha\beta}(x) \rightarrow \eta_{\alpha\beta} + h'_{\alpha\beta}(x') = \frac{\partial x^\gamma}{\partial x'^\alpha} \frac{\partial x^\delta}{\partial x'^\beta} (\eta_{\gamma\delta} + h_{\gamma\delta}(x)) .$$

Upon explicit computation to leading order in ε^α one finds that the *functional* shift in $h_{\alpha\beta}$ is given by

$$h'_{\alpha\beta}(x) - h_{\alpha\beta}(x) = \partial_\alpha \varepsilon_\beta(x) + \partial_\beta \varepsilon_\alpha(x) + O(\varepsilon^2) . \quad (3.72)$$

Therefore, the condition on the allowed ‘weak-diffeomorphisms’ is $|\partial_\alpha \varepsilon_\beta| \ll 1$. Incidentally, the weak diffeomorphisms act on the weak gravitational field $h_{\alpha\beta}$ as a gauge symmetry for a Lorentz-covariant symmetric tensor field. Expanding now the particle action to leading order in $h_{\alpha\beta}$ one finds

$$-m \int_\gamma d\sigma \sqrt{-(\eta_{\alpha\beta} + h_{\alpha\beta}) \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma}} = S[\gamma]_{\text{Minkowski}} + \frac{1}{2} m \int d\tau h_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} + O(h^2) , \quad (3.73)$$

where τ in the last term is defined as the proper time with respect to the Minkowski metric. The term linear in $h_{\alpha\beta}$ can be rewritten as

$$\frac{1}{2} \int d^4x h_{\alpha\beta} T^{\alpha\beta} ,$$

where $T^{\alpha\beta} = m \int d\tau \dot{x}^\alpha \dot{x}^\beta \delta^{(4)}(x - x(\tau))$ is the standard definition of the particle's energy momentum tensor in SR (cf. (3.3)). Hence, we conclude that the leading perturbative coupling of gravity, as dictated by the Principle of Equivalence, conforms to the general arguments based on SR. However, the Principle of Equivalence is much more powerful, giving a precise resummation of powers of $h_{\alpha\beta}$ in the matter-gravity coupling.

Newtonian Limit

We now sharpen the interpretation of Γ as gravitational forces by taking the non-relativistic limit of the equation of motion (3.69). It is useful to transform it from the proper-time parametrization to coordinate time parametrization. Using the chain rule one finds

$$\frac{d^2 x^\alpha}{dt^2} + \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{dt} \frac{dx^\nu}{dt} = f(t) \frac{dx^\alpha}{dt} , \quad (3.74)$$

with

$$f(t) = -\frac{d^2 t/d\tau^2}{(dt/d\tau)^2} .$$

We may consider now the spatial components of (3.74) in the weak-field regime, for *stationary* fields, $\partial_t h_{\alpha\beta} \ll \partial_i h_{\alpha\beta}$, and in the non-relativistic limit, $v^i = dx^i/dt \ll 1$. Keeping terms up to first order in velocities we obtain

$$\frac{dv_i}{dt} \approx \frac{1}{2} \partial_i h_{00} - \sum_j v_j (\partial_j h_{0i} - \partial_i h_{0j}) . \quad (3.75)$$

The first term is the standard gradient gravitational force for a field with potential

$$\phi_N \approx -\frac{1}{2} h_{00} , \quad (3.76)$$

whereas the second term has the form of a Coriolis force. To see this, notice that we may define the effective ‘angular velocity’ vector

$$\vec{\Omega} = \frac{1}{2} \vec{\partial} \times \vec{h} , \quad (3.77)$$

where $\vec{h} = (h_{0i})$. The effective Newton equation takes then the form

$$\frac{d\vec{v}}{dt} \approx -\vec{\partial} \phi_N + 2 \vec{v} \times \vec{\Omega} , \quad (3.78)$$

which contains the potential and Coriolis terms.⁴ Hence, we confirm that the Principle of Equivalence *unifies* what in Newtonian theory is regarded as separate ‘inertial’ and ‘gravitational’ forces.

In practice this analysis shows that, once we pick a non-rotating frame, the leading Newtonian approximation to the motion of test particles is characterized by an effective metric whose time-time component is fixed by the Newtonian potential:

$$ds_{\text{Newtonian}}^2 \approx -(1 + 2\phi_N) dt^2 + \dots . \quad (3.79)$$

⁴If $\vec{\Omega}$ really represents the angular velocity of a rotating frame, there should be a centrifugal force proportional to Ω^2 in (3.78). This term is indeed found by going to second order in the weak field expansion (see the Problem on Relativistic Rotation).

3.3 Minimal Coupling

We are ready to use the formal machinery introduced so far to couple a generic Lorentz-invariant Lagrangian $\mathcal{L}(\Psi \dots_{b\dots}^{a\dots}, \partial_a, \dots)$ with arbitrary ‘matter’ fields in tensor representations of the Lorentz group, to an external gravitational field characterized by a metric field $g_{\alpha\beta}$. The strategy is identical to the one that worked before for free particles. We start by covering the spacetime X^{1+3} with *frefos* and specify the action in terms of the limit of a fine partition:

$$S = \lim_{\{P\} \rightarrow X^{1+3}} \sum_P \Delta_P S , \quad (3.80)$$

with

$$\Delta_P S \approx (\Delta^4 X_P) \mathcal{L}(\Psi \dots_{b\dots}^{a\dots}, \partial/\partial X_P^a, \dots)|_P \quad (3.81)$$

the element of action in a *frefo*. We now convert to the reference *fido* with the translation of Lorentz tensorial structures to Diff tensorial structures. As for the volume measure we have

$$(\Delta^4 X_P) \approx (\Delta^4 x) \left| \det \left(\frac{\partial X_P}{\partial x} \right) \right| = \Delta^4 x |\det(E_P)| = \Delta^4 x \sqrt{-\det(g_{\alpha\beta})} ,$$

or, in the more familiar differential form $d^4 X = d^4 x \sqrt{-g}$. Hence, the coupling to gravity is reduced to a mere exercise in notation:

$$S[\Psi, g_{\alpha\beta}] = \int_{X^{1+3}} d^4 x \sqrt{-g} \mathcal{L}(\Psi \dots_{\beta\dots}^{\alpha\dots}, \nabla_\alpha, \dots) . \quad (3.82)$$

Namely, we replace Lorentz tensors $T^{a\dots}$ by their Diff-tensor forms $T^{\alpha\dots}$ using the map in (3.27), we replace the ordinary derivatives ∂_a by the covariant derivatives ∇_α , and we integrate the Lagrangian over spacetime with a Diff-invariant measure $d^4 x \sqrt{-g}$. If we want to include parity-violating interactions, we use the Diff-tensor Levi-Civita, $\epsilon_{\alpha\beta\gamma\delta}$, in place of the Minkowski one, ϵ_{abcd} . This algorithmic procedure is known as the ‘minimal coupling prescription’.

Examples of the procedure include the Lagrangian of a scalar field,

$$S_{\text{KG}} = - \int d^4 x \sqrt{-g} \left(\frac{1}{2} g^{\mu\nu} \nabla_\mu \phi \nabla_\nu \phi + V(\phi) \right) , \quad (3.83)$$

with field equation

$$\nabla^2 \phi = V'(\phi) ,$$

or the Maxwell theory in a gravitational field,

$$S_{\text{Maxwell}} = - \int d^4 x \sqrt{-g} \left(\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - J_\mu A^\mu \right) , \quad (3.84)$$

where $A_\mu = E_\mu^a A_a$, $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu = \partial_\mu A_\nu - \partial_\nu A_\mu$, $F^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta}$. The field equations

$$\nabla_\mu F^{\mu\nu} = -J^\nu$$

may also be written as

$$\partial_\mu (\sqrt{-g} F^{\mu\nu}) = -\sqrt{-g} J^\nu .$$

We have already seen the minimal coupling algorithm in action in deriving the point particle dynamics, which we quote here again in its more general version (which allows the study of the massless limit) as a covariant form of (2.46):

$$S_P[x^\alpha, e] = \frac{1}{2} \int d\sigma \left(\frac{1}{e(\sigma)} g_{\alpha\beta} \frac{dx^\alpha}{d\sigma} \frac{dx^\beta}{d\sigma} - m^2 e(\sigma) \right) \quad (3.85)$$

with the Minkowski equation of motion $du^a/d\tau = 0$ transformed into $u^\beta \nabla_\beta u^\alpha = 0$. In an analogous fashion, the free equations for the spin degree of freedom of a particle, $dS^a/d\tau = 0$ and $S_a u^a = 0$, generalize to

$$\frac{\nabla S^\alpha}{d\tau} = u^\beta \nabla_\beta S^\alpha = 0, \quad S_\alpha u^\alpha = 0 \quad (3.86)$$

in a general gravitational field. In components, the first of these equations reads

$$\frac{dS^\alpha}{d\tau} + \Gamma_{\beta\gamma}^\alpha u^\beta S^\gamma = 0. \quad (3.87)$$

These equations can be used to describe the precession of gyroscopes in a gravitational field.

Energy-Momentum Tensor

We can now repeat the weak-gravity analysis which was already done for the case of point particles. Setting $g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta}$ and keeping the linear term in $h_{\alpha\beta}$ one finds

$$S[\Psi, \eta + h] = S[\Psi, \eta] + \frac{1}{2} \int d^4x h_{\alpha\beta} \left(2 \frac{\partial \mathcal{L}}{\partial h_{\alpha\beta}} + \eta^{\alpha\beta} \mathcal{L} \right) \Big|_{h=0} + O(h^2), \quad (3.88)$$

where we have used $\sqrt{-\det(\eta + h)} = 1 + \frac{1}{2} \eta^{\alpha\beta} h_{\alpha\beta} + O(h^2)$, easily proved in the eigenvalue basis. The quantity in parenthesis in (3.88) measures the *linear response* of the matter system to the perturbation by a gravitational field. By explicit computation we see that it coincides with the energy-momentum tensor of the matter system in all the usual cases involving point particles, scalars and electromagnetic fields. Rather than quoting these results for the case of weak fields, we can actually generalize the result for the case of linear response in an arbitrary gravitational field, i.e. the behavior under a perturbation $g_{\alpha\beta} \rightarrow g_{\alpha\beta} + \delta g_{\alpha\beta}$. In this case we have

$$S[\Psi, g + \delta g] = S[\Psi, g] + \frac{1}{2} \int d^4x \sqrt{-g} \delta g_{\alpha\beta} T^{\alpha\beta} + O(\delta g^2), \quad (3.89)$$

where the symmetric tensor $T^{\alpha\beta}$ is given by

$$T^{\alpha\beta} = 2 \frac{\partial \mathcal{L}}{\partial g_{\alpha\beta}} + g^{\alpha\beta} \mathcal{L}. \quad (3.90)$$

In obtaining this equation we used $\sqrt{-(g + \delta g)} = \sqrt{-g} + \delta \sqrt{-g} = \sqrt{-g} + \frac{1}{2} \sqrt{-g} g^{\alpha\beta} \delta g_{\alpha\beta} + O(\delta g^2)$, which is again easily proved in the eigenvalue basis. Another useful identity which is proved in a similar way is $\delta g^{\alpha\beta} = -g^{\alpha\mu} g^{\beta\nu} \delta g_{\mu\nu}$. Using all these results, it is easy to show that, for a system of particles, scalars and electromagnetic fields, we get a total linear-response function

$$T^{\alpha\beta} = T_{\text{particles}}^{\alpha\beta} + T_{\text{scalar}}^{\alpha\beta} + T_{\text{Maxwell}}^{\alpha\beta},$$

where

$$\begin{aligned}
T_{\text{particles}}^{\alpha\beta} &= \sum_p m_p \int d\tau_p \frac{dx_p^\alpha}{d\tau_p} \frac{dx_p^\beta}{d\tau_p} \frac{\delta^{(4)}(x - x_p(\tau_p))}{\sqrt{-g}} \\
T_{\text{scalar}}^{\alpha\beta} &= \partial^\alpha \phi \partial^\beta \phi - g^{\alpha\beta} \left(\frac{1}{2} (\partial\phi)^2 + V(\phi) \right) \\
T_{\text{Maxwell}}^{\alpha\beta} &= -F^{\alpha\gamma} F_\gamma^\beta - \frac{1}{4} g^{\alpha\beta} F^{\mu\nu} F_{\mu\nu} .
\end{aligned} \tag{3.91}$$

The answer is just the Diff-tensor forms of the Noetherian energy-momentum tensors for each of these systems. Hence, we conclude that the energy-momentum tensor measures the response of the matter system to a general gravitational perturbation in an arbitrary gravitational field. In fact, the simplicity of the minimal coupling prescription implies that it is often better to compute the energy momentum tensor through the formulas (3.89) and (3.90) than using the canonical methods based on Noether's theorem.

Limitations of Minimal Coupling and General Covariance

The minimal coupling algorithm described in the previous sections follows from the Principle of Equivalence and allows us to write down the effects of gravitation for an arbitrary system. The question thus arises, why is it called *minimal* ?

The reason is that there could be higher order terms in the Lagrangian, depending on second derivatives of the metric tensor or higher, that vanish in Minkowski space (hence are invisible in SR) but do not vanish locally in a *frefo* (recall that the Principle of Equivalence only requires the first derivatives of the metric to vanish in free fall). In other words, there could be different scalar Lagrangians with the same Minkowskian limit. This phenomenon occurs for example when considering powers of covariant derivatives of tensors.

Let V_ρ be a covariant vector field and consider the double covariant derivative $\nabla_\mu \nabla_\nu V_\rho$. This is a rank-three covariant tensor by construction. Its flat-space limit, obtained setting $g = \eta$, is the Lorentz tensor $\partial_a \partial_b V_c$. This Lorentz tensor is obviously symmetric in the indices a, b . However, the original general tensor is not always symmetric in the indices μ, ν . Noting that $\nabla_\mu \sim \partial_\mu + \Gamma_\mu$, we see that this tensor contains terms of the form $\partial_\mu \partial_\nu V$ that are indeed symmetric in μ, ν , and terms of the form $\Gamma_\mu \partial_\nu V$ and $\Gamma_\mu \Gamma_\nu V$, that vanish in a local *frefo*, but there are also terms of the form $(\partial_\mu \Gamma_\nu) V$ that *do not* vanish in a local *frefo*, and are not necessarily symmetric in μ, ν . These terms do vanish in a global *frefo* for a fictitious gravitational field, but not in a general one.

Hence, any term in the Lagrangian involving more than one covariant derivative of tensors, with some antisymmetry property, will not be generated by the minimal coupling algorithm. Such terms are perfectly possible and do not violate the strictly formulated Principle of Equivalence, because the latter specifies that free fall removes gravitational effects only at the level of the metric and its first derivatives, leaving the second derivatives (curvature) untouched (except in fictitious fields, where we remove the gravitational effects to all orders). All these terms involve higher powers of derivatives in the Lagrangian, and thus are suppressed by inverse powers of the spatial range of variation of the gravitational field.

In order to cope with these subtleties, one usually extends the method of minimal coupling to the so-called 'Principle of General Covariance', which states that one should write down all possible terms in the Lagrangian that are Diff scalars and that reduce to Lorentz scalars in the flat space limit $g \rightarrow \eta$. This may include terms that effectively reduce to vanishing Lorentz

tensors, such as the antisymmetric second covariant derivative of tensor fields. In keeping with the strategy of the ‘locality principle’, all such terms should be organized in a long-distance expansion and their effects only show up at sufficiently small scales.

Spin Connection

There is an alternative view of geometry, due essentially to E. Cartan, in which one shifts focus from the metric to its formal ‘square root’, namely the vector fields that specify the local *frefos* at each point. Recall the basic definition of the Riemannian metric out of the Principle of Equivalence:

$$g_{\mu\nu}(P) = E_P^a{}_\mu(P) E_P^b{}_\nu(P) \eta_{ab}, \quad E_P^a{}_\alpha(P) \equiv \left. \frac{\partial X_P^a}{\partial x^\alpha} \right|_P. \quad (3.92)$$

The four covariant Diff-vector fields

$$e_\alpha^a(P) = E_P^a{}_\alpha(P) \equiv \left. \frac{\partial X_P^a}{\partial x^\alpha} \right|_P, \quad (3.93)$$

labeled by $a = 0, 1, 2, 3$ are called the *tetrads* or ‘moving frames’ (*repère mobile*), and they can be regarded as a ‘Lorentzian’ square-root of the metric. They contain the same information as the metric, but with larger redundancy, since any choice of tetrads can be rotated by a different Lorentz transformation at each point and still produce the same metric.

Notice the subtle difference between matrix of functions $E_P^a{}_\alpha(x)$ and $e_\alpha^a(P)$. The first is a different function for each P , but each such function extends beyond P . The second is just the value of these functions at P . In practice, anytime we evaluate $E_P^a{}_\alpha(x)$ at P , we can use instead $e_\alpha^a(P)$, like in the definition of Diff tensors out of Lorentz tensors (3.27). On the other hand, derivatives of $E_P^a{}_\alpha(x)$ *do not* coincide with derivatives of $e_\alpha^a(x)$, since the first type of derivative gives the shape around P of a function which is different for different choices of P , whereas the second type of derivative just monitors how the P -values change as we change P . The main situation where the subtlety arises is in the definition of the Christoffel symbols (3.59), where the expression

$$\partial_\mu E_P^a{}_\alpha|_P = \left. \frac{\partial^2 X_P^a}{\partial x^\mu \partial x^\alpha} \right|_P$$

appears, in which one evaluates the functions at the point P only *after* the second derivative of $X_P^a(x)$ has been taken. In particular, this expression is different from $\partial_\mu e_\alpha^a$. It turns out that the parametrization of this subtle difference is a useful object to carry around, called the *spin connection*:

$$\omega_\mu{}^a{}_b \equiv -e_b^\nu (\partial_\mu e_\nu^a - \partial_\mu E_P^a{}_\nu) . \quad (3.94)$$

We may remove all reference to local *frefos* by expressing the E_P part in terms of the Christoffel symbols to get

$$\omega_\mu{}^a{}_b = -e_b^\nu (\partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\alpha e_\alpha^a) \equiv -e_b^\nu \nabla_\mu e_\nu^a = e_\nu^a \nabla_\mu e_b^\nu . \quad (3.95)$$

The last form, in terms of the covariant derivative of the tetrads, shows that, unlike the Christoffel symbol, the spin connection does transform as a covariant vector in the Diff index μ . It also shows that the Lorentz covariant form $\omega_{\mu ab}$ is antisymmetric in the ab Lorentz indices, as a result of the metric compatibility of the Christoffel connection.

On the other hand, the spin connection does not have a homogeneous transformation rule with respect to the local Lorentz rotations, due to the derivatives of the tetrads:

$$\partial_\mu e_\alpha^a \rightarrow \partial_\mu (L^a{}_b e_\alpha^b) = L^a{}_b \partial_\mu e_\alpha^b + e_\alpha^b \partial_\mu L^a{}_b .$$

In particular, under $e_\mu^a \rightarrow L^a{}_b e_\mu^b$, the spin connection transforms as

$$\omega_\mu \rightarrow L (\omega_\mu + \partial_\mu) L^{-1} , \quad (3.96)$$

in matrix notation for the Lorentz indices. Therefore, the operator $D_\mu = \partial_\mu + \omega_\mu$ is a covariant derivative with respect to the local (gauge) Lorentz symmetry! Namely, given a vector field $V^a = e_\mu^a V^\mu$, the derivative

$$D_\mu V^a = \partial_\mu V^a + \omega_\mu^a{}_b V^b \quad (3.97)$$

is Lorentz-covariant, $D_\mu V^a \rightarrow L^a{}_b D_\mu V^b$.

These definitions imply that the Christoffel symbols can also be calculated in terms of the Lorentz-covariant derivatives of the tetrads:

$$\Gamma_{\mu\nu}^\alpha = e_\alpha^a D_\mu e_\nu^a \equiv e_\alpha^a \left(\partial_\mu e_\nu^a + \omega_\mu^a{}_b e_\nu^b \right) \quad (3.98)$$

in a general gravitational field. In this way, the two representations of a tensor in terms of ‘curved’ or ‘flat’ indices have ∇_μ and D_μ as the respective covariant derivative operators:

$$D_\mu V^a = e_\alpha^a \nabla_\mu V^\alpha . \quad (3.99)$$

This relation expresses the fact that the spin connection does not introduce new degrees of freedom and could be used to ‘define’ ω_μ .

So, the subtlety in handling the derivatives of the tetrads turned out to hold the solution of the problem of implementing the local Lorentz rotations as a conventional gauge redundancy, the spin connection being the corresponding gauge field. All this is very nice, but a bit formal. What is the use of the Cartan formalism in physics? While it offers some simplifications in the computation of curvature tensors, by far the main application is to the extension of the minimal coupling procedure to spinor fields.

The simplicity of the minimal coupling prescription is based on the deceptively ‘trivial’ replacement of latin (Lorentz) indices by greek (Diff) indices. This is ultimately possible because the point-by-point transformation matrices of Diff tensors belong to the $GL(4, \mathbb{R})$ group and, the Lorentz group being a subgroup, we can always get a representation of the Lorentz group by restricting a representation of the general linear group. Alas, the converse is not true, as there are representations of $O(1, 3)$ that *cannot* be extended into representations of $GL(4, \mathbb{R})$. These are precisely the spinorial representations, used to describe fermions in nature. For this reason, to couple spinors to gravity one must use the spin connection, which emphasizes the Lorentz indices, instead of the Christoffel one.

In particular, the Dirac operator in the presence of gravitation is defined as

$$\not{D} \equiv \gamma^\mu D_\mu = \gamma^\mu \left(\partial_\mu + \frac{i}{2} \omega_\mu^{ab} \sigma_{ab} \right) , \quad (3.100)$$

where $\gamma^\mu = e_\alpha^\mu \gamma^\alpha$ is a basis of curved-space Dirac matrices satisfying the generalized Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = -2g^{\mu\nu}$.

3.4 An Interlude: Vernacular Riemannian Geometry

We have argued on physical terms that the Principle of Equivalence, together with Special Relativity, sets the spacetime arena to be given by Riemannian geometry. In this section we offer some contact with the traditional geometrical language in which Riemannian geometry is discussed.

In the mathematical literature, spaces with a positive-definite metric determining local distances

$$d\ell^2 = \sum_{ij=1}^d g_{ij} dy^i dy^j \quad (3.101)$$

are called d -dimensional Riemannian manifolds.⁵ The meaning of this expression is that any curve γ , parametrized by $y^i(\sigma)$, has length

$$\ell(\gamma) = \int_{\gamma} d\sigma \sqrt{\sum_{ij} g_{ij}(y(\sigma)) \frac{dy^i}{d\sigma} \frac{dy^j}{d\sigma}}. \quad (3.102)$$

An intuitive interpretation of a Riemannian metric can be given by visualizing the manifold X^d as a curved submanifold embedded⁶ in a higher-dimensional flat space $\mathbb{R}^D$, with a sufficiently large $D > d$. Let X^I denote Cartesian coordinates on $\mathbb{R}^D$ and let the submanifold X^d of interest be parametrized by functions $X^J(y^j)$. At a given point on X^d we can consider the tangent hyperplane, generated by the vectors of components

$$E_i^I = \frac{\partial X^I}{\partial y^i},$$

with scalar product

$$E_i \cdot E_j = \sum_{I=1}^D E_i^I E_j^I.$$

The length of a curve $X^I(\sigma)$ in $\mathbb{R}^D$ is given by

$$\ell(\gamma) = \int_{\gamma} d\sigma \sqrt{\sum_I \frac{dX^I}{d\sigma} \frac{dX^I}{d\sigma}}. \quad (3.103)$$

However, when the curve lies inside X^d we can also describe it by functions $y^i(\sigma)$, then

$$\frac{dX^I}{d\sigma} = \sum_j \frac{dy^j}{d\sigma} \frac{\partial X^I}{\partial y^j} = \sum_j E_j^I \frac{dy^j}{d\sigma}$$

and the formula (3.102) follows with the metric “induced” from the embedding:

$$g_{ij}^{(\text{induced})} = \sum_{I=1}^D E_i^I E_j^I. \quad (3.104)$$

⁵Technically, the notion of manifold requires that the space be locally like a subset of $\mathbb{R}^d$, in the topological and differential sense. In practice, this means that it can be parametrized by well-behaved coordinate systems.

⁶We use the word ‘embedded’ as opposed to ‘immersed’ to emphasize that X^d is required to have no self-intersections when viewed from $\mathbb{R}^D$, so that it will look locally like $\mathbb{R}^d$ and thus be a manifold.

The starting point of Riemannian geometry is the abstraction of the embedding space. Once the intrinsic metric g_{ij} is defined, Riemann concentrates on those properties that follow from the metric alone.⁷ The vectors E_i at each point P generate the so-called tangent space of the manifold at P denoted $T_P X^d$. Smoothness of X^d means that $T_P X^d$ is a good linear approximation to X^d around any point P .⁸




Figure 3.4: A two-dimensional surface X^2 embedded in $\mathbb{R}^3$. The vectors $\vec{E}_1$ and $\vec{E}_2$ give a basis of a local tangent space. The induced metric is given by $g_{ij} = \vec{E}_i \cdot \vec{E}_j$.

The appropriate generalization for GR is that of a Riemannian manifold with Lorentzian signature, meaning that the eigenvalues of the metric in a special (free fall) frame, have signs $(-+++)$ (these are sometimes called Pseudo-Riemannian manifolds). Instead of a Euclidean tangent space at each point, the Lorentzian manifolds have tangent spaces isomorphic to Minkowski spacetime at each point. Again, we can say that Minkowski spacetime (together with SR) is a good linear approximation to the spacetime manifold, in the neighborhood of any point P . Hence, we see explicitly how the Principle of Equivalence embodies the geometrical interpretation of GR. From this standpoint, we could derive GR by simply providing a physical interpretation of each element of Riemannian geometry, such as the notions of connection, torsion, curvature, isometries, etc.

In the mathematical literature, the free-fall equation of motion goes under the name of ‘geodesic equation’. This refers to the related problem of finding ‘smallest length’ paths between two points in a curved manifold. If our manifold is equipped with a metric $dl^2 = \sum_{ij} g_{ij} dy^i dy^j$, the length along a given path can be obtained by (3.102). Extrema of this functional define the

⁷Incidentally, a famous theorem by the even more famous John Nash shows that any given Riemannian metric g_{ij} can be induced by embedding the manifold X^d in $\mathbb{R}^D$, for a sufficiently large dimension D .

⁸The attentive reader will notice the similarity between the vectors E_i and the analogous objects in the *frefo/fido* correspondence. Indeed, they play essentially the same role.

locally shortest paths, i.e. the geodesics. But we see that the mathematical problem is identical to ours, except for the use of positive definite metrics, rather than Lorentzian ones. Accordingly, the local geodesic equation has the form

$$\sum_i T^i \nabla_i T^j = 0, \quad (3.105)$$

where $\nabla \sim \partial + \Gamma$ in a completely analogous fashion and $T^i = dy^i/d\ell$ is the tangent vector to the geodesic curve, parametrized by the proper length ℓ . In particular, the explicit formula for the Christoffel symbols in terms of the first derivatives of the metric is exactly the same. So, we can adopt the geometry language and say that the motion in a gravitational field is equivalent to the motion along geodesics of a curved spacetime manifold. In our case, the geodesics are not ‘minimal length’ paths, but rather ‘maximum proper-time’ ones.

There are surprisingly few differences at the local level between the Riemannian formalism of Euclidean and Lorentzian signature manifolds. Since the signature of the metric can be changed by a formal analytic continuation in the eigenvalues of the metric, and this is a continuous procedure, the tensor differential calculus remains largely intact, up to some signs. At the global level, there are very important differences, associated to the existence of a causal structure in the Lorentzian signature case. This is related to the existence of restrictions on physical paths of particles and light rays, namely the locally measured speed of light must be maximal. These global differences become largest when dealing with cosmology and black holes.

The construction of Riemannian geometry from embeddings can be further generalized to define an induced metric for any p -dimensional submanifold Σ^p of X^d , with $p < d$, which of course will also be a submanifold of $\mathbb{R}^D$. Iterating the procedure gives the induced metric on Σ

$$g_{uv}^{(\Sigma)} = \sum_I \frac{\partial X^I}{\partial \sigma^u} \frac{\partial X^I}{\partial \sigma^v} = \sum_{ij} g_{ij} \frac{\partial y^i}{\partial \sigma^u} \frac{\partial y^j}{\partial \sigma^v}, \quad (3.106)$$

where σ^u label coordinates on Σ^p which can be specified by embedding functions $y^j(\sigma)$ into X^d or $X^I(y(\sigma))$ into $\mathbb{R}^D$.⁹

The trick of ‘inducing’ the metric from an ambient space can be repeated for any tensor. Given a covariant Diff tensor in X^d , with components $T_{ij\dots}$, we can define a Diff tensor on any submanifold Σ^p by

$$\left(T^{(\Sigma)}\right)_{uv\dots} = T_{uv\dots} \equiv \sum_{ij\dots} T_{ij\dots} \frac{\partial y^i}{\partial \sigma^u} \frac{\partial y^j}{\partial \sigma^v} \dots, \quad (3.107)$$

an operation which is also known as ‘pulling-back’ the tensor from X^d onto Σ^p . Once the tensor is pulled back on Σ^p , one can operate as usual for any Σ -intrinsic calculations, including the raising and lowering of indices with g_{uv} , taking Diff(Σ)-covariant derivatives, or defining invariant integrals and exterior calculus. The volume form for integration of any Diff-scalar over Σ^p is $d^p\sigma \sqrt{|\det(g_{uv})|}$ and the Levi-Civita tensor for integration of forms is $\epsilon_{u_1\dots u_p} = \sqrt{|\det(g_{uv})|} [u_1 \dots u_p]$.

⁹We will often abuse notation and identify tensors on Σ^p by the indices alone. In particular, we write $g_{uv}^{(\Sigma)} = g_{uv}$ when no confusion arises.

Parallel Transport, Holonomy and Curvature

In general, if a vector V^i is defined at each point of a curve $x^j(\sigma)$ with tangent vector $T^i = dx^i/d\sigma$, we can define the covariant derivative along the curve as

$$\frac{\nabla V^i}{d\sigma} \equiv \nabla_T V^i \equiv \sum_j T^j \nabla_j V^i . \quad (3.108)$$

If this covariant derivative vanishes, one says that the vector V is being defined by *parallel transport* along the curve $x^i(\sigma)$. The justification for this name comes from the fact that any two vectors V and W , carried by parallel transport along the curve, $\nabla_T V = \nabla_T W = 0$, satisfy

$$\frac{d}{d\sigma} \left(\sum_{ij} g_{ij} V^i W^j \right) = \sum_{ijk} T^k \nabla_k (g_{ij} V^i W^j) = 0 ,$$

as a consequence of the metric compatibility property, $\nabla_k g_{ij} = 0$. Hence, the local metric definition of angles and norms is preserved under parallel transport.¹⁰ We can write more explicitly the parallel-transport equation as

$$\frac{dV^i}{d\sigma} + \sum_j (\Gamma_T)^i_j V^j = 0 ,$$

where we define the matrix $(\Gamma_T)^i_j \equiv \sum_k T^k \Gamma_{kj}^i$. This equation can be solved to give the vector at some point Q , parallel-transported from some other point P , along the curve γ_{PQ} , in terms of a formal path-ordered product

$$V(Q) = \mathcal{P} \exp \left(- \int_{\gamma_{PQ}} \Gamma_T \right) V(P) . \quad (3.109)$$

For an infinitesimal displacement we have $V(Q) = (1 - \Delta\sigma \Gamma_T) V(P)$, so that the path-ordered product can be defined as the formal limiting product over an infinitesimal partition of the path

$$\mathcal{P} \exp \left(- \int_{\gamma_{PQ}} \Gamma_T \right) = \lim_{\Delta\sigma_s \rightarrow 0} \prod_s e^{-\Delta\sigma_s \Gamma_T(s)} .$$

The concept of parallel transport can be used to define a notion of curvature. Consider a closed curve γ_P , based at P , given by the function $x^i(\sigma)$ that starts and returns to a point P , i.e. $x^i(0) = x^i(2\pi)$, with σ an angular variable. The corresponding closed-path parallel transporter

$$U[\gamma_P] = \mathcal{P} \exp \left(- \oint_{\gamma_P} \Gamma_T \right)$$

depends both on the curve and the point P , and is called *holonomy*. Since parallel transport preserves norms and angles, $U(\gamma_P)$ is some matrix in the $SO(d)$ group (for an orientable manifold), where d is the dimension of the Riemannian manifold.¹¹ In Minkowski space, or a ‘flat’

¹⁰Notice that a geodesic is defined as a proper-length parametrized curve, $x^i(\ell)$, whose tangent vector, $T^i = dx^i/d\ell$, is ‘carried forward’ by parallel transport along itself, $\nabla_T T = 0$.

¹¹In four-dimensional Lorentzian signature the holonomy is a matrix in the proper orthochronous Lorentz group $SO_+(1,3)$, for an orientable and time-orientable spacetime.

space in general, the holonomy of any curve is the identity matrix. In this case, the parallel transport of vectors between two arbitrary points is independent of the path joining them. A local measure of the deviation from flatness, i.e. a mathematical definition of ‘curvature’, is obtained by considering the limit of very small closed curves.

Let us make one little ‘square’ composed of four curves intersecting in pairs, using the parameters of these curves as coordinates (x, y) of a two-dimensional submanifold with size of order ϵ . We consider tangent vector fields X and Y whose only nonvanishing components are $X^x = Y^y = 1$ respectively. Denoting $X^i = \partial z^i / \partial x$ and $Y^i = \partial z^i / \partial y$ the components in an arbitrary coordinate system, we have the following identity

$$[X, Y]^i \equiv \sum_j (X^j \partial_j Y^i - Y^j \partial_j X^i) = 0, \quad (3.110)$$

where the so-defined vector field $[X, Y]$ is known as the *Lie bracket* of X and Y . Hence, when two vector fields are used to coordinate a two-dimensional submanifold, their Lie bracket vanishes.¹²




Figure 3.5: A quadrilateral round-trip at P with direction vectors X, Y and holonomy $U(\gamma_P) = U_{-\epsilon Y} U'_{-\epsilon X} U'_{\epsilon Y} U_{\epsilon X}$.

The holonomy around this quadrilateral round trip is given by

$$U(\gamma_P) = U_{-\epsilon Y} U'_{-\epsilon X} U'_{\epsilon Y} U_{\epsilon X}$$

where we have, with quadratic precision in ϵ ,

$$U_{\epsilon X} = e^{-\epsilon \Gamma_X} = 1 - \epsilon \Gamma_X + \frac{1}{2} \epsilon^2 (\Gamma_X)^2 + O(\epsilon^3)$$

with a similar definition for $U_{\epsilon Y}$. The primed transporters are computed from the displaced versions of the connection

$$\Gamma'_Y = \Gamma_Y + \epsilon \partial_X \Gamma_Y + O(\epsilon^2), \quad \Gamma'_X = \Gamma_X + \epsilon \partial_Y \Gamma_X + O(\epsilon^2),$$

with the notation $\partial_X = \sum_i X^i \partial_i$ and analogously for Y . Upon explicit evaluation of the holonomy we find the first non-trivial term occurring at order ϵ^2 and given by

$$U(\gamma_P) = 1 - \epsilon^2 (\partial_X \Gamma_Y - \partial_Y \Gamma_X + \Gamma_X \Gamma_Y - \Gamma_Y \Gamma_X) + O(\epsilon^3).$$

¹²The converse statement is also true and it is a particular case of a theorem by Frobenius.

Evaluating now the derivatives in index notation and using $[X, Y] = 0$ one finds

$$U(\gamma_P)_j^i = \delta_j^i - \epsilon^2 \sum_{k,l} X^k Y^l R_{jkl}^i + O(\epsilon^3) , \quad (3.111)$$

where the coefficients

$$R_{jkl}^i = \partial_k \Gamma_{lj}^i + \sum_n \Gamma_{kn}^i \Gamma_{lj}^n - (k \leftrightarrow l) \quad (3.112)$$

define the Riemann tensor. Despite its explicit definition in terms of non-tensor objects, it is a Diff tensor with 20 independent components in four dimensions.¹³ Equivalently, it can be represented in terms of the spin connection via the mixed Diff-Lorentz tensor $R_{kl}^{ab} = \sum_{ij} e_i^a e^{jb} R_{jkl}^i$,

$$R_{kl}^{ab} = \partial_k \omega_l^{ab} - \partial_l \omega_k^{ab} + [\omega_k, \omega_l]^{ab} , \quad (3.113)$$

thus taking the form of a standard non-abelian field strength of Yang–Mills type, associated to the local $SO(1,3)$ gauge group.

The Riemann tensor has the purely geometrical interpretation of characterizing completely the local curvature: a Riemannian space is flat, i.e. metrically $\mathbb{R}^d$, if and only if the Riemann tensor vanishes identically. On the other hand, the Riemann tensor describes the deviation from flatness, isolating the combinations of second derivatives of the metric that cannot be eliminated by a clever choice of coordinates.

Hypersurfaces

For submanifolds Σ there is an important distinction between *intrinsic* and *extrinsic* curvature. The intrinsic curvature is the Riemannian curvature of the induced metric $g_{uv}^{(\Sigma)}$, whereas the extrinsic curvature defines how the submanifold is ‘bent’ inside the ambient manifold. When Σ^p is a codimension-one submanifold, $p = d - 1$, also called a *hypersurface*, the ‘bending’ is determined by the normal unit vector to Σ , which we denote by n^i . The two-index object $\Pi_{ij} = g_{ij} - n_i n_j$ is a projector that gives the induced metric when restricted to the tangent space of Σ , i.e. in intrinsic coordinates for Σ one has $g_{uv}^{(\Sigma)} = \Pi_{uv}^{(\Sigma)}$.

The bending of Σ inside X^d is then related to the failure of n^i to follow parallel transport when moved around Σ . Hence we define the *extrinsic curvature* as the hypersurface-tensor ‘induced’ on Σ by $\nabla_i n_j$:

$$K_{uv} = \sum_{ij} \nabla_i n_j \frac{\partial y^i}{\partial \sigma^u} \frac{\partial y^j}{\partial \sigma^v} . \quad (3.114)$$

To compute the covariant derivative of the normal vector, one can define n^i slightly away from Σ by extending it as the unit tangent vector of geodesics orthogonal to Σ . Then, one can see by explicit computation that $K_{uv} = K_{vu}$ is symmetric, using the symmetry of the Christoffel symbols.

A useful device when dealing with hypersurfaces is the ‘Gaussian normal coordinate system’. Given the hypersurface Σ , it is always possible to choose coordinates locally (near a point in Σ) so that the full metric takes the form

$$ds^2 = dn^2 + \sum_{u,v} g_{uv}(n, \sigma) d\sigma^u d\sigma^v , \quad (3.115)$$

¹³An equivalent, if more elegant expression with manifest tensorial form is $([\nabla_X, \nabla_Y] - \nabla_{[X, Y]}) V^i = \sum_{klj} X^k Y^l R_{jkl}^i V^j$, valid even for the case that $[X, Y] \neq 0$.

where n denotes a proper-distance coordinate normal to Σ , so that we can write the normal unit vector as $\sum_i n^i \partial_i = \partial/\partial n$. The orientation of n^i is conventionally chosen such that the system of coordinates (n, σ^u) is positively oriented in X^d . The lines of constant σ^u are geodesics orthogonal to Σ . In these coordinates $g_{nn} = 1$ and $g_{nu} = 0$, giving d conditions on the metric functions that can be fixed by adjusting the d functions defining a diffeomorphism.

Gaussian normal coordinates make the symmetry of K_{uv} completely manifest:

$$K_{uv} = \frac{1}{2} \partial_n g_{uv} = \frac{1}{2} \sum_{ij} \mathcal{L}_n g_{ij} \frac{\partial y^i}{\partial \sigma^u} \frac{\partial y^j}{\partial \sigma^v}, \quad (3.116)$$

where the second expression is the Lie derivative as defined in previous chapters.

Hypersurfaces can be boundaries of proper dimension- d subregions of the manifold. Then, using again the Gaussian normal coordinates, it is easy to prove Gauss' integration theorem on a general manifold, for the integral of a total derivative:

$$\int_{V^d} \sum_i \nabla_i J^i = \int_{\partial V^d} J_n, \quad (3.117)$$

where $J_n = \sum_i J_i n^i$ is the normal component of the vector field J^i , where now the normal vector is pointing 'outward' from the inside of V^d .

For Lorentzian signature, hypersurfaces can be spacelike, timelike or null according to whether the normal vector n^μ is respectively timelike, spacelike or null. This leads to some sign differences and specific constructions (not discussed here) for the null case. For timelike normal, $\tau^\mu \tau_\mu = -1$, the projector that defines the intrinsic metric is $\Pi_{\mu\nu} = g_{\mu\nu} + \tau_\mu \tau_\nu$ and the Gaussian normal coordinates are sometimes termed 'synchronous gauge':

$$ds^2 = -d\tau^2 + \sum_{ij} g_{ij}(\tau, \sigma) d\sigma^i d\sigma^j, \quad (3.118)$$

where τ now stands for the proper time along the timelike geodesics at fixed σ^i and orthogonal to Σ . One subtlety regarding sign conventions is the Lorentzian generalization of the divergence theorem, which retains the form (3.117) applied to regions such that ∂V^d is the union of piecewise timelike or spacelike hypersurfaces (no null pieces), with the rule that the normal vector is outward-pointing when spacelike and inward-pointing when timelike.

Non-Metrical Connections and Derivatives

The concept of connection (covariant derivative) is actually more primitive than the notion of metric. One can define ∇_μ axiomatically and from there one can reach the concepts of parallel transport, holonomy and curvature, since all these are concepts of the general theory of fiber bundles. This conceptual track emphasizes the similarity between GR and other gauge theories. The most general covariant derivatives are still of the form $\nabla \sim \partial + \Gamma$, but now the connection coefficients $\Gamma_{\mu\nu}^\alpha$ need not be symmetric in the lower indices, with the antisymmetric part $T_{\mu\nu}^\alpha = \Gamma_{\mu\nu}^\alpha - \Gamma_{\nu\mu}^\alpha$ defining the *torsion*. One can check explicitly that this is always a tensor.

Even in the presence of a metric, it is possible to have non-standard connections, in the sense that they might not be metric compatible, i.e. $\nabla_\alpha g_{\mu\nu} \neq 0$, and/or they might have torsion. However, demanding the connection to be symmetric (torsion free) and metric compatible, fixes

it completely to be the Levi–Civita connection, defined by the Christoffel symbols. In practice, one can always write an arbitrary connection as the sum of the Levi–Civita connection and a remainder. This remainder may depend on torsion terms and/or contain the degrees of freedom that violate the metric compatibility, and it can be regarded as a non-minimal coupling in covariant derivatives. The important fact about this remainder is that it is always a tensor. In this way, we may still use the standard Riemannian language for physics applications, with the non-standard connection degrees of freedom interpreted as a type of exotic tensor ‘matter’ that does not follow the minimal coupling prescription.

This freedom in the use of non-standard connections shows up in modern generalizations of GR, in the context of supergravity theories. Historically, this was also the arena for various forgotten alternative theories, tried by the ‘founding fathers’. In particular, Einstein contemplated connections with torsion in some of his attempts at constructing a ‘unified theory’ of gravity and electromagnetism. Even earlier, Weyl considered connections satisfying $\nabla_i g_{jk} = A_i g_{jk}$ for some vector field A_i . Such connections only preserve angles under parallel transport (rather than lengths).

Chapter 4

Curvature

The Principle of Equivalence says that the gravitational field trivializes locally when entering free fall. Therefore, the interesting gravitational dynamics must be encoded in the first non-trivial correction, the tidal forces. In this chapter we find that the relativistic treatment of tidal forces has a neat geometrical character in the formalism of Riemannian geometry: it is none other than curvature.

In our previous introduction to the notion of Riemannian curvature we emphasized the modern track based on parallel transport of vectors through round trips. However, the first encounter with the notion of curvature was historically associated to the famous Euclid’s ‘fifth postulate’, stating that initially parallel straight lines would stay so forever. The failure of this fifth postulate is an equivalent characterization of curvature. Interestingly, it is within this avatar that curvature arises most naturally in the physics of gravitation. Namely, geodesics that fail to stay parallel *accelerate* relative to one another. This is the tidal acceleration.

4.1 Relativistic Tidal Acceleration

The basic property of a fictitious gravitational field is the existence of a global free-fall frame that ‘transforms away’ gravity globally. In the free-fall frame, initially parallel trajectories stay so in time, with vanishing relative acceleration. On the other hand, in a real gravitational field, the Principle of Equivalence only removes the gravitational field at a point, leaving a residual relative acceleration between nearby particle trajectories. The corresponding residual forces are called ‘tidal forces’.

It should be clear that the tidal accelerations we are discussing are a property of the background gravitational field, and not a result of the gravitational attraction between the probe particles. What we have in mind is the radial infall trajectories around Earth for instance: two test particles falling with a relative angle with respect to the center of the planet will accelerate towards each other, even when we neglect their mutual gravitational attraction.

We can discuss tidal forces in the Newtonian theory by considering a family of free-falling particles along a curve $\vec{x}(s)$ at time $t = 0$. We also assume that the initial velocities of these particles vary continuously along the family, i.e. the function

$$\vec{v}(s, 0) = \frac{d\vec{x}}{dt}(s, 0)$$

is a continuous vector function of s . Now, we let them develop in time, so that the set of trajectories span a two-dimensional surface coordinated by (s, t) . The relative separation at time t of two particles with parametric distance Δs is

$$\Delta \vec{x} = \Delta s \frac{d\vec{x}}{ds}(s, t) ,$$

to first order in Δs . Denoting the transverse separation vector $\vec{\ell} = d\vec{x}/ds$, the rate of separation of the falling trajectories is $\Delta s d\vec{\ell}/dt$ and the relative acceleration $\Delta s d^2\vec{\ell}/dt^2$. Using now the equation of motion

$$\frac{d^2\vec{x}}{dt^2}(s, t) = -\vec{\partial} \phi_N(\vec{x}(s, t))$$

we obtain the final expression for the tidal accelerations,

$$a_{\text{tidal}}^i = \frac{d^2 \ell^i}{dt^2} = -(\vec{\ell} \cdot \vec{\partial}) \partial^i \phi_N = \sum_j \mathcal{R}_j^i \ell^j , \quad (4.1)$$

which indeed depend on the second derivatives of the gravitational potential and reproduces our previously introduced tidal tensor $\mathcal{R}_j^i = -\partial^i \partial_j \phi_N$.




Figure 4.1: Family of free-fall trajectories parametrized by global time and fiducial parameter s . The relative separation of the falling points is given by $\Delta s d\vec{x}/ds$.

In the relativistic case, we consider an entirely analogous situation, i.e. a family of free-falling geodesics $x^\mu(\tau, s)$, parametrized by the geodesic label, s , and proper time τ . The separation and velocity four-vectors are defined by

$$\ell^\mu = \frac{dx^\mu}{ds} , \quad u^\nu = \frac{dx^\nu}{d\tau} . \quad (4.2)$$

We can now define covariant derivatives along parametrized curves as

$$\frac{\nabla}{ds} \equiv \nabla_\ell \equiv \ell^\mu \nabla_\mu, \quad \frac{\nabla}{d\tau} \equiv \nabla_u \equiv u^\mu \nabla_\mu. \quad (4.3)$$

With this definition, the free-fall equation of motion may be rewritten as

$$\frac{du^\mu}{d\tau} + \Gamma_{\alpha\beta}^\mu u^\alpha u^\beta = \nabla_u u^\mu = u^\alpha \nabla_\alpha u^\mu = 0. \quad (4.4)$$

Using these formulas, one can explicitly write

$$\nabla_u \ell^\alpha = u^\mu \nabla_\mu \ell^\alpha = \frac{d\ell^\alpha}{d\tau} + \Gamma_{\mu\nu}^\alpha u^\mu \ell^\nu = \frac{d^2 x^\alpha}{d\tau ds} + \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{d\tau} \frac{dx^\nu}{ds}.$$

Since this expression is symmetric in s, τ , it is actually equal to

$$\frac{d^2 x^\alpha}{ds d\tau} + \Gamma_{\mu\nu}^\alpha \frac{dx^\mu}{ds} \frac{dx^\nu}{d\tau} = \ell^\mu \nabla_\mu u^\alpha.$$

Hence, we have proven

$$\nabla_u \ell^\alpha = u^\mu \nabla_\mu \ell^\alpha = \ell^\mu \nabla_\mu u^\alpha = \nabla_\ell u^\alpha \quad (4.5)$$

With these preliminaries, we are ready to define the relative acceleration of neighboring geodesics as

$$a_{\text{tidal}}^\mu = (\nabla_u)^2 \ell^\mu = u^\alpha \nabla_\alpha (u^\beta \nabla_\beta \ell^\mu) = u^\alpha \nabla_\alpha (\ell^\beta \nabla_\beta u^\mu), \quad (4.6)$$

where we have used (4.5) in the last equality. Manipulating this expression we find

$$a_{\text{tidal}}^\alpha = u^\mu (\nabla_\mu \ell^\nu) \nabla_\nu u^\alpha + u^\mu \ell^\nu \nabla_\nu \nabla_\mu u^\alpha + u^\mu \ell^\nu [\nabla_\mu, \nabla_\nu] u^\alpha. \quad (4.7)$$

Using again (4.5) in the first term and the geodesic condition $u^\alpha \nabla_\alpha u^\beta = 0$ we finally obtain

$$a_{\text{tidal}}^\alpha = u^\mu \ell^\nu [\nabla_\mu, \nabla_\nu] u^\alpha. \quad (4.8)$$

Thus, we have proven that the physical distinction between real and fictitious gravitational fields can be locally parametrized by the commutator of covariant derivatives acting on vector fields. Such an object defines the relativistic version of the tidal tensor, the so-called Riemann tensor.

4.1.1 The Riemann Tensor

The commutator of covariant derivatives acting on an arbitrary vector field, $[\nabla_\mu, \nabla_\nu] V^\alpha$, defines a tensor object that vanishes in a fictitious gravitational field (i.e. a reparametrized Minkowski space) because its flat-space version satisfies $[\partial_a, \partial_b] V^c = 0$ identically. Upon direct evaluation, we find

$$[\nabla_\mu, \nabla_\nu] V^\alpha = R^\alpha_{\rho\mu\nu} V^\rho, \quad (4.9)$$

where

$$R^\alpha_{\rho\mu\nu} = \partial_\mu \Gamma_{\nu\rho}^\alpha + \Gamma_{\mu\sigma}^\alpha \Gamma_{\nu\rho}^\sigma - (\mu \leftrightarrow \nu) \quad (4.10)$$

is the Riemann tensor. Although this expression in terms of the connection coefficients does not look obviously tensorial, the formula (4.9) ensures that the collection of coefficients $R^\alpha_{\rho\mu\nu}$ indeed transform as a tensor. As expected, it contains second derivatives of the metric that need

not vanish anywhere in a real gravitational field, even in a *frefo* frame. The explicit calculation above can be repeated for the case of a covariant vector field,

$$[\nabla_\mu, \nabla_\nu] V_\alpha = -R^\rho{}_{\alpha\mu\nu} V_\rho , \quad (4.11)$$

and for a general tensor

$$[\nabla_\mu, \nabla_\nu] T^{\dots\alpha\dots}_{\dots\beta\dots} = R^\alpha{}_{\rho\mu\nu} T^{\dots\rho\dots}_{\dots\beta\dots} + \dots - R^\rho{}_{\beta\mu\nu} T^{\dots\alpha\dots}_{\dots\rho\dots} - \dots . \quad (4.12)$$

The Riemann tensor controls tidal gravitational effects in the relativistic theory. Substituting back in (4.8) we find

$$a^\alpha_{\text{tidal}} = u^\mu \ell^\nu R^\alpha{}_{\rho\mu\nu} u^\rho , \quad (4.13)$$

whose non-relativistic approximation at low velocities and static weak fields $g_{00} \approx -1 - 2\phi_N$ recovers the Newtonian tidal tensor for static fields

$$R^i{}_{\mu\nu j} u^\mu u^\nu \approx R^i{}_{00j} = -R^i{}_{0j0} \approx -\partial_j \partial^i \phi_N . \quad (4.14)$$

Identities

The Riemann tensor has in principle 256 components, which are reduced to 20 independent ones due to a large set of symmetries. First, from the definition we immediately have antisymmetry in the second pair of indices:

$$R^\alpha{}_{\rho\mu\nu} = -R^\alpha{}_{\rho\nu\mu} . \quad (4.15)$$

We also have the so-called cyclic identity of the last three indices:

$$R^\alpha{}_{\rho\mu\nu} + R^\alpha{}_{\nu\rho\mu} + R^\alpha{}_{\mu\nu\rho} = 0 = R^\alpha{}_{[\rho\mu\nu]} . \quad (4.16)$$

In addition, working with the completely covariant form $R_{\sigma\rho\mu\nu} = g_{\sigma\alpha} R^\alpha{}_{\rho\mu\nu}$, there is antisymmetry in the first pair of indices and symmetry in the swap of the first and second pairs of indices:

$$R_{\alpha\rho\mu\nu} = -R_{\rho\alpha\mu\nu} , \quad R_{\alpha\rho\mu\nu} = R_{\mu\nu\alpha\rho} . \quad (4.17)$$

The simplest way of proving all these identities is to pass to a *frefo* frame at a given point P . In this frame the connection coefficients vanish at P and the Riemann tensor can be written:

$$R_{abcd} \Big|_{P-\text{frefo}} = \frac{1}{2} (\partial_b \partial_c g_{ad} + \partial_a \partial_d g_{bc} - \partial_b \partial_d g_{ac} - \partial_a \partial_c g_{bd}) , \quad (4.18)$$

so that the algebraic identities can be verified directly. Once established in the *frefo* frame, being tensorial relations, they are valid in an arbitrary frame.

In addition to these purely algebraic properties, the Riemann tensor also satisfies a set of differential identities called *Bianchi identities*:

$$\nabla_{[\sigma} R_{\alpha\rho]\mu\nu} = 0 . \quad (4.19)$$

These too can be proven using the trick of the *frefo* frame, since the covariant derivative of the Riemann tensor simplifies as

$$\nabla_e R^d{}_{cab} \Big|_{P-\text{frefo}} = \partial_e \partial_a \Gamma^d_{bc} - \partial_e \partial_b \Gamma^d_{ac} , \quad (4.20)$$

and further writing this expression in terms of second derivatives of the metric, the Bianchi identities are easily proved directly. In more mathematical terms, the Bianchi identities are nothing but the Jacobi identity for the covariant derivative operator:

$$0 = [\nabla_\mu, [\nabla_\nu, \nabla_\rho]] + \text{cyclic permutations}.$$

It is worth mentioning at this point that the notion of curvature, and the associated Riemann tensor, can be introduced provided one is handed a connection (a covariant derivative operator) independently of the existence of a metric. However, some of the properties listed above, such as the quoted forms of the cyclic (4.16) and Bianchi (4.19) identities and the simplifying *frefo* expression (4.20), require that the torsion vanishes (symmetric connections). The rest of symmetry properties which depend upon lowering indices and using (4.18) further require that the connection be metric compatible, $\nabla_\alpha g_{\mu\nu} = 0$. Since we only use the Levi-Civita connection in these notes, we make no such distinctions, but the reader should be aware of these subtleties when consulting mathematical references.

Riemann Is all You Need

At this point we have seen that the Riemann tensor gives a mathematical characterization of curvature and at the same time controls tidal gravitational effects. It is a Diff tensor depending on second derivatives of the metric. Since the metric defines the relativistic analog of a gravitational potential, the Riemann tensor is a natural candidate to enter the Lagrangian of the gravitational field. One natural question at this point is whether there might exist other independent tensors constructed from second derivatives of the metric which deserve consideration. We will give a negative answer, using an argument that is motivated by a somewhat different question.

The condition (3.63) gives a mathematically precise statement of the Principle of Equivalence. It is interesting to ask whether the required *frefo* can always be introduced at an arbitrary point, for a *generic* $g_{\alpha\beta}$, or some restriction must be put on the metric of spacetime. In other words, is it always possible to enter free fall in any gravitational field? To study this, consider constructing the map between the *frefo* frame X_P^a and the *fido* frame x^α in a perturbative expansion. Without loss of generality we may define the additive normalization of the coordinates so that $x^\alpha(P) = X_P^a(P) = 0$. Then we can expand the function $x^\alpha(X_P^a)$ as

$$x^\alpha(X_P^a) = A_a^\alpha X_P^a + \frac{1}{2} B_{ab}^\alpha X_P^a X_P^b + \frac{1}{6} C_{abc}^\alpha X_P^a X_P^b X_P^c + O(X_P^4), \quad (4.21)$$

where the A, B, C are collections of constant coefficients arising from the Taylor expansion. Using now (3.62) we find for the metric in the *frefo* frame

$$g_{ab}(P) = A_a^\alpha A_b^\beta g_{\alpha\beta}(P),$$

so that $g_{ab}(P) = \eta_{ab}$ by simply choosing the A_a^α to diagonalize and rescale appropriately the original metric matrix at point P . There is in fact an ambiguity by multiplication of A_a^α by a Lorentz transformation, since η_{ab} is invariant under those. Computing now the Christoffel symbols in the X_P -frame, using for example (3.38), we find

$$\Gamma_{bc}^a(P) = (A^{-1})_\alpha^a A_b^\beta A_c^\gamma \Gamma_{\beta\gamma}^\alpha(P) + (A^{-1})_\alpha^a B_{bc}^\alpha,$$

so that $\Gamma_{bc}^a(P) = 0$ can be readily arranged by setting

$$B_{bc}^\alpha = -A_b^\beta A_c^\gamma \Gamma_{\beta\gamma}^\alpha(P) .$$

We conclude that the Principle of Equivalence, as given by (3.63), can always be reached in an appropriate *frefo*. On the other hand, the second derivatives of the metric at P , or equivalently the first derivatives of the connection, will be affected by the 80 independent coefficients C_{abc}^α . All in all there are 100 independent components in the second derivatives of the metric $\partial_\alpha \partial_\beta g_{\mu\nu}(P)$, so that we cannot remove all of them by a clever choice of *frefo*. The offset of 20 components gives the degrees of freedom of the *curvature* in a four-dimensional spacetime. The argument just given here is known in the mathematical literature as the ‘local flatness theorem’, since it specifies that any Riemannian space is ‘flat’ to linear order in a local expansion, with ‘intrinsic’ curvature effects arising only in second order in this local expansion.

These 20 non-removable degrees of freedom in second derivatives of the metric equal in number the 20 independent components of the Riemann tensor, and this is no coincidence. Expanding the metric to second order in the expansion (4.21) we have

$$g_{ab}(X_P) \Big|_{P-\text{frefo}} = \eta_{ab} - \frac{1}{3} E_{acbd} X_P^c X_P^d + O(X_P^3) , \quad (4.22)$$

where the set of coefficients E_{acbd} , proportional to the second derivatives of the metric, must be symmetric in the (ab) and (cd) pairs of indices. Computing these second derivatives we find

$$\partial_c \partial_d g_{ab} \Big|_P = -\frac{1}{3} (E_{acbd} + E_{adbc})$$

which we can substitute back into (4.18) to prove explicitly that $E_{abcd} = R_{abcd}$. Therefore, we see that the Riemann tensor characterizes completely the quadratic deviation from the Minkowski spacetime. More invariantly, we say that $T_P X^{1+3}$, the tangent space of X^{1+3} at P is not only a copy of $\mathbb{R}^{1+3}$, Minkowski spacetime, but it also provides a linear approximation to the geometry of X^{1+3} , with the deviation from flatness arising at the level of second derivatives of the metric, and invariantly characterized by the Riemann tensor.

From the physical point of view, the main conclusion is that Riemann is the essentially unique tensor that can be constructed with up to two derivatives of the metric tensor.

4.1.2 Ricci, Einstein and Weyl Tensors

The symmetry properties imply that the Riemann tensor produces only one independent second rank tensor under contraction, the so called Ricci tensor

$$R_{\mu\nu} = R^\alpha{}_{\mu\alpha\nu} , \quad (4.23)$$

which is symmetric $R_{\mu\nu} = R_{\nu\mu}$. There is also a unique scalar that can be formed from contraction of Ricci, known as the Ricci scalar

$$R = g^{\mu\nu} R_{\mu\nu} . \quad (4.24)$$

Contracting the Bianchi identities we obtain their projection over the Ricci tensors,

$$\nabla^\mu R_{\mu\nu} = \frac{1}{2} \partial_\nu R , \quad (4.25)$$

so that it becomes natural to define the so-called Einstein tensor

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \quad (4.26)$$

for which the Bianchi identities amount to ‘covariant conservation’:

$$\nabla^\mu G_{\mu\nu} = 0 . \quad (4.27)$$

Another interesting object is the Weyl tensor, defined by

$$C_{\mu\nu}{}^{\rho\sigma} \equiv R_{\mu\nu}{}^{\rho\sigma} - 2R_{[\mu}{}^{[\rho}\delta_{\nu]}{}^{\sigma]} + \frac{1}{3} R \delta_{[\mu}{}^{\rho}\delta_{\nu]}{}^{\sigma} . \quad (4.28)$$

It can be interpreted as the part of the Riemann tensor that is not controlled by the Ricci component, as all its contractions vanish identically. Its main application is as a criterion for conformal equivalence of metrics, namely two metrics related by a local rescaling $g_{\mu\nu}(x) \rightarrow \Omega^2(x)g_{\mu\nu}(x)$ have the same Weyl tensor $C^\mu{}_{\nu\rho\sigma}$ (with one index up; other index positions pick up powers of Ω).

Chapter 5

Einstein's Equations

With the formal machinery introduced so far, we are able to solve for the dynamics of arbitrary matter systems in a prescribed gravitational field, as specified by some metric tensor $g_{\mu\nu}$. In this so-called *probe approximation*, the spacetime metric is considered as a fixed external background. The remaining part of the problem is the specification of the dynamics of the gravitational field itself or, in other words, the Lagrangian of the metric tensor, which ultimately determines which metrics actually occur depending on the energy content of spacetime.

5.1 The Matter-Gravitation Dynamics

Since $g_{\mu\nu}$ plays the role of a gravitational potential, we seek the relativistic analog of the Newtonian action

$$S_{\text{Newton}} = \frac{1}{8\pi G} \int dt d^3x \phi_N \vec{\partial}^2 \phi_N$$

in a scalar Lagrangian linear in second derivatives of the metric, i.e.

$$S = S_G + S_m = \int d^4x \sqrt{-g} (\mathcal{L}_G(g_{\mu\nu}) + \mathcal{L}_m(\Psi, \nabla_\mu)) , \quad (5.1)$$

with $\mathcal{L}_m$ the Lagrangian of generic matter degrees of freedom Ψ , minimally coupled to $g_{\mu\nu}$ as in the previous sections, and $\mathcal{L}_G \sim \partial\partial g$ is the purely gravitational Lagrangian, a Diff scalar. The obvious choice $\nabla^2 g$ vanishes because of the metric compatibility of the affine connection. However, we have argued that there is essentially a unique tensor containing up to two derivatives of the metric, the Riemann tensor. Therefore, we are forced to choose as the Lagrangian for the gravitational field the unique scalar constructed from the Riemann tensor, the Ricci scalar. This leads to the Hilbert action ¹

$$S_H = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} R , \quad (5.2)$$

where κ^2 is proportional to G , the precise constant of proportionality to be determined below.

In fact, there is a superficially more important term that can be added to the purely gravitational Lagrangian: an additive renormalization of R by a constant. It is more important

¹In a dramatic episode in November 1915, Hilbert derived the field equations from this action almost simultaneously with Einstein; the question of priority is still debated by historians.

because it does not contain derivatives and thus dominates the long distance physics. This is the famous *cosmological constant*, Einstein's 'biggest blunder', by his own reported admission and, in hindsight, one of his major discoveries!

The complete Lagrangian of gravity and matter then reads,

$$S[g, \Psi] = S_H + S_m = \int d^4x \sqrt{-g} \left(\frac{1}{2\kappa^2} (R - 2\Lambda) + \mathcal{L}_m(\Psi, \nabla) \right) + \dots, \quad (5.3)$$

with the dots standing for corrections in higher derivatives, in the spirit of Effective Field Theory. It can be shown that, in the purely gravitational sector, the terms that can appear are any Diff-scalars made of polynomial combinations of covariant derivatives and powers of the Riemann tensor so that, to some extent, the Riemann tensor is the fundamental building block of all terms in the higher-derivative Lagrangian.

5.1.1 Einstein's Equations

We are now ready to derive Einstein's equations for the gravitational field, by simply stabilizing the variation of the Lagrangian with respect to variations of the metric $g_{\mu\nu}$. We first consider the variation of the gravitational Lagrangian. It contains three terms,

$$\delta(\sqrt{-g} (R - 2\Lambda)) = \delta(\sqrt{-g})(R - 2\Lambda) + \sqrt{-g} \delta g^{\mu\nu} R_{\mu\nu} + \sqrt{-g} g^{\mu\nu} \delta R_{\mu\nu}.$$

Using the relations $\delta\sqrt{-g} = \frac{1}{2} \sqrt{-g} g^{\mu\nu} \delta g_{\mu\nu}$ and $\delta g^{\mu\nu} = -g^{\mu\alpha} g^{\nu\beta} \delta g_{\alpha\beta}$ we can evaluate the first two terms. As for the last one we have

$$\delta R_{\mu\nu} = \partial_\alpha \delta \Gamma_{\mu\nu}^\alpha - \partial_\mu \delta \Gamma_{\alpha\nu}^\alpha + O(\Gamma \delta \Gamma).$$

The terms proportional to $\Gamma \delta \Gamma$ vanish on a free fall frame at a given point. Now, although the connection Γ is not a tensor, the difference between two connections is a tensor, since the offending inhomogeneous terms vanish. So we can find the complete expression by covariantizing the previous one,

$$\delta R_{\mu\nu} = \nabla_\alpha \delta \Gamma_{\mu\nu}^\alpha - \nabla_\mu \delta \Gamma_{\alpha\nu}^\alpha. \quad (5.4)$$

The same type of reasoning shows that

$$\delta \Gamma_{\mu\nu}^\alpha = \frac{1}{2} g^{\alpha\sigma} (\nabla_\mu \delta g_{\sigma\nu} + \nabla_\nu \delta g_{\sigma\mu} - \nabla_\sigma \delta g_{\mu\nu}). \quad (5.5)$$

So, the final result is that $g^{\mu\nu} \delta R_{\mu\nu} = \nabla_\mu f^\mu$, with f^μ a four-vector field linear in covariant derivatives of $\delta g_{\mu\nu}$. Since this variation contributes a total derivative, it is tempting to neglect it on the grounds that the metric is fixed on the boundary of spacetime, in accord with common practice in the Lagrangian formalism. There is a subtlety though, in that f^μ depends on $\delta g_{\mu\nu}$ through its covariant derivative, and the condition that these vanish is not quite the same as the vanishing of $\delta g_{\alpha\beta}$. In fact, as remarked by York, Gibbons and Hawking (YGH), one can add a boundary term to Hilbert's action in such a way that those terms are cancelled out, and one is left with a standard variational principle.² The resulting equations are identical to those obtained by a naive procedure in which one simply neglects the f^μ term, so that we will not worry about this issue at this point.

²The York–Gibbons–Hawking term is given by $S_{YGH} = \frac{1}{\kappa^2} \oint K$, where the integral is defined over the boundary, and K is the trace of the extrinsic curvature associated to the induced metric.

Collecting all terms, the variation of the gravitational action is

$$\delta S_H = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left(-R^{\mu\nu} + \frac{1}{2} g^{\mu\nu} (R - 2\Lambda) \right) \delta g_{\mu\nu} . \quad (5.6)$$

Next we turn to the matter sector. Here, the first order variation defines the *energy-momentum tensor*, according to (3.89) and (3.90), through

$$\delta S_m = \int d^4x \sqrt{-g} \frac{1}{2} T^{\mu\nu} \delta g_{\mu\nu} . \quad (5.7)$$

In terms of the Lagrangian,

$$T^{\mu\nu} = 2 \frac{\partial \mathcal{L}_m}{\partial g_{\mu\nu}} + g^{\mu\nu} \mathcal{L}_m . \quad (5.8)$$

Finally, we can put all the terms together and write down Einstein's equations as

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} (R - 2\Lambda) = \kappa^2 T_{\mu\nu} , \quad (5.9)$$

a set of ten non-linear second-order partial differential equations. The non-linearity, which persists even for vacuum solutions, makes it highly non-trivial to find exact solutions. Flat Minkowski space $g_{\alpha\beta} = \eta_{\alpha\beta}$ is of course a solution with $T_{\mu\nu} = \Lambda = 0$. The vacuum equations are obtained by setting to zero the matter sector and tracing the equation to obtain $R - 2(R - 2\Lambda) = -R + 4\Lambda = 0$. Hence,

$$R_{\mu\nu} = \Lambda g_{\mu\nu} \quad (5.10)$$

gives the vacuum equations. Manifolds solving the vacuum Einstein equations are called *Einstein manifolds*. For $\Lambda = 0$ we have the so-called *Ricci-flat manifolds*.

The cosmological constant term is equivalent to a vacuum energy on the matter energy-momentum tensor. For example, if the matter sector contains a scalar field with energy-momentum tensor

$$T_{\mu\nu}^{(\phi)} = \partial_\mu \phi \partial_\nu \phi - g_{\mu\nu} \left(\frac{1}{2} (\partial\phi)^2 + V(\phi) \right) ,$$

we see that the shift $V(\phi) \rightarrow V(\phi) + \Lambda/\kappa^2$ in the origin of potential energies introduces the cosmological constant if it was zero on the left hand side of Einstein's equations. Alternatively, for a perfect fluid *ansatz*,

$$T_{\mu\nu}^{(\text{fluid})} = (p + \rho) U_\mu U_\nu + p g_{\mu\nu} ,$$

the cosmological constant is equivalent to a matter component with the equation of state $p = -\rho = -\Lambda/\kappa^2$, so that we may fold Λ into the non-gravitational Lagrangian as a matter of convention and write the equations as ³

$$G_{\mu\nu} = \kappa^2 T_{\mu\nu} . \quad (5.11)$$

It remains to determine the absolute normalization of the action, i.e. the relation between κ and ordinary Newton's constant G . This can be fixed by taking the Newtonian limit. We begin by rewriting (5.11) in the equivalent form

$$R_{\mu\nu} = \kappa^2 (T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T) , \quad (5.12)$$

³This innocent-looking statement causes the so-called 'cosmological constant problem'; the astonishing fact that the measured value of Λ , given by $G\Lambda \sim 10^{-120}$, is so incredibly small, despite the multitude of independent contributions to Λ from the matter sector, each of them potentially much larger than the experimental bound.

where $T = g^{\alpha\beta}T_{\alpha\beta}$ is the trace of the energy-momentum tensor. In the non-relativistic limit for static and small fields, we can replace the metric with the Newtonian form in computing the Ricci tensor,

$$ds^2 \approx -(1 + 2\phi_N) dt^2 + \dots, \quad (5.13)$$

where we assume that the g_{0i} components are negligible once we choose a frame without rotation, so that $G_{0i} \approx R_{0i} \approx 0$. Then, for any *static* metric with $g_{0i} \approx 0$ one has $R_{00} \approx -\frac{1}{2}\vec{\partial}^2 g_{00} \approx \vec{\partial}^2 \phi_N$, quite independently of the value of g_{ij} . Then, the time-time component of Einstein's equations in the 'trace-reversed' form (5.12) gives ⁴

$$\vec{\partial}^2 \phi_N \approx R_{00} = \kappa^2 (T_{00} + \frac{1}{2} T), \quad (5.14)$$

where we have used $1 + 2\phi_N \approx 1$ on the right hand side. For a matter source composed of a non-relativistic system of slow particles, we have $T_{00} \approx \rho_m$ and $T = -T_{00} + \sum_i T_{ii} \approx -T_{00}$, so that the right-hand side of (5.14) is $\frac{\kappa^2}{2} \rho_m$, which agrees with the Poisson equation for $\kappa^2 = 8\pi G$. The final form of Einstein's equations being

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi G T_{\mu\nu}. \quad (5.15)$$

Einstein's equations can be viewed as a set of partial differential equations determining the metric $g_{\mu\nu}$ on a spatial surface at time t , given initial data on a spatial surface at time $t = t_0$. The initial data are 'positions' $g_{\mu\nu}(t_0)$, and 'velocities' $\partial_t g_{\mu\nu}(t_0)$. Apparently, the system of ten equations just determines the ten components of the metric, but this is actually deceptive, since there are four degrees of freedom not determined by the equations, related to Diff covariance. Indeed, only the six spatial equations

$$G_{ij} = 8\pi G T_{ij}$$

contain second derivatives in time and are true 'dynamical equations'. The remaining four,

$$G^{0\mu} = 8\pi G T^{0\mu} \quad (5.16)$$

only contain up to first derivatives in time of $g_{\mu\nu}$. This follows from the Bianchi identities, $\nabla_\mu G^{\mu\nu} = 0$, which can be written as

$$\nabla_0 G^{0\mu} = - \sum_i \nabla_i G^{\mu i}.$$

The right hand side contains derivatives up to at most second order in time, thus $G^{0\mu}$ can only contain, at most, *first* time derivatives of the metric. The conclusion is that the four equations (5.16) must be imposed as constraints on initial data. This remark is very useful in practice, when faced with the task of solving Einstein's equations. In many situations with high symmetry, it is useful to evaluate the $G_{00} = 8\pi G T_{00}$ equation first, since it often expresses "constants of the motion" in the combined gravity/matter system. More generally, the method of solving Einstein's equations in 'Hamiltonian' form, emphasizing the initial-value formulation, is the starting point for numerical simulations of Einstein equations, a subject that has grown enormously in precision and scope in the last decade.

⁴Notice that we have been careful in not specifying that $g_{ij} \approx \delta_{ij}$. This is actually not true, since g_{ij} contains pieces proportional to ϕ_N that are not visible to Newtonian test particles moving at slow velocities. This will be further clarified in a later chapter of these notes.

5.1.2 Energy-Momentum ‘Local’ Conservation

The Bianchi identity on the Einstein tensor $\nabla_\mu G^{\mu\nu} = 0$ implies, via the Einstein equations, a covariant conservation law for the energy-momentum tensor of all non-gravitational degrees of freedom,

$$\nabla_\mu T^{\mu\nu} = 0 . \quad (5.17)$$

We can derive this relation, as well as the Bianchi identity on $G_{\mu\nu}$, from the invariance of the action under reparametrizations. Since S_m is manifestly invariant under smooth reparametrizations: $x \rightarrow x'(x)$, we have for infinitesimal ones $x' = x - \xi(x) + O(\xi^2)$,

$$\delta_\xi S_m = \int d^4x \sqrt{-g} \frac{1}{2} T^{\mu\nu} \delta_\xi g_{\mu\nu} = 0 , \quad (5.18)$$

straight from the definition of $T_{\mu\nu}$ in (5.7), applied to a particular variation. Here $\delta_\xi g_{\mu\nu}$ is the variation of the metric *functions* under the Diff map, that is to say

$$g'_{\mu\nu}(x) - g_{\mu\nu}(x) \equiv \delta_\xi g_{\mu\nu}(x) + O(\xi^2) . \quad (5.19)$$

Using now the Taylor expansion

$$g'_{\mu\nu}(x) = g'_{\mu\nu}(x' + \xi) = g'_{\mu\nu}(x') + \xi^\alpha \partial_\alpha g'_{\mu\nu}(x') + O(\xi^2)$$

and the tensor law

$$g'_{\mu\nu}(x') = \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu} g_{\alpha\beta}(x)$$

to linear order in ξ , we find the Lie derivative like in our previous discussion for symmetries in Minkowski spacetime,

$$\delta_\xi g_{\mu\nu} \equiv \mathcal{L}_\xi g_{\mu\nu} = \xi^\alpha \partial_\alpha g_{\mu\nu} + g_{\mu\alpha} \partial_\nu \xi^\alpha + g_{\nu\alpha} \partial_\mu \xi^\alpha = \nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu , \quad (5.20)$$

where the last expression using covariant derivatives of the vector field ξ^μ is special to the Lie derivative of the metric.

Introducing (5.20) back into the variation of the action (5.18) we obtain, upon integration by parts,

$$0 = \int d^4x \sqrt{-g} \nabla_\mu T^{\mu\nu} \xi_\nu \quad (5.21)$$

for all ξ_ν , which leads to the covariant conservation relation (5.17).

The covariant conservation equation $\nabla_\mu T^{\mu\nu} = 0$ is quite surprising at first sight, for $T_{\mu\nu}$ is the energy-momentum tensor of *matter* alone. When we discussed the electromagnetic field coupled to charged particles, the energy-momentum tensor of the particles was not conserved by itself, as some energy is transferred from the particles to the field degrees of freedom and vice versa (2.134). Only after we added the intrinsic energy-momentum density of the Maxwell field we obtained a conserved energy-momentum tensor. For that reason, the conservation of the matter energy-momentum, without apparent need for a ‘purely gravitational’ piece of the energy-momentum tensor, is somewhat puzzling.

On the other hand (5.17) is forced upon us by the Principle of Equivalence, since it is the Diff-covariant version of the SR equation, valid in a *frefo*

$$\partial_a T^{ab} \Big|_{P-\text{frefo}} = 0 . \quad (5.22)$$

Clearly, the gravitational field, and whatever its energy density might be, disappears locally in free fall. Therefore, no strictly local notion of ‘gravitational energy density’ is possible in GR, despite the fact that the Newtonian approximation does contain the notion of ‘gravitational self-energy’, which we may associate to ‘the gravitational field’. For instance, the Newtonian gravitational self-energy of a mass distribution, defined as the work needed to assemble it from infinity, can be rewritten, using the field equation and integration by parts, as the integral⁵

$$U_N = -\frac{G}{2} \int d^3x d^3x' \frac{\rho_m(\vec{x}) \rho_m(\vec{x}')}{|\vec{x} - \vec{x}'|} = \frac{1}{2} \int d^3x \rho_m(\vec{x}) \phi_N(\vec{x}) = -\frac{1}{8\pi G} \int d^3x \vec{\partial} \phi_N \vec{\partial} \phi_N , \quad (5.23)$$

which, in the third form, may be interpreted as an energy stored ‘in the field’. Notwithstanding this, it is clear that the physical notion of gravitational self-energy has a global character by the way it is defined, using the infinitely dilute limit as a reference. Accordingly, we will see in future sections that global notions of energy can be defined in GR, and local notions of energy can be defined in approximate descriptions in which some spacetime structure is maintained as ‘non-dynamical’.

Problem: Free Fall from Einstein’s Equations

Consider a dust fluid with energy-momentum tensor $T_{\mu\nu} = \rho U_\mu U_\nu$. Show that its covariant conservation, which can be viewed as a consequence of Einstein’s equations, is powerful enough to determine the free fall of the dust, namely that the fluid flow is geodesic.

5.1.3 Isometries and Conservation Laws

There remains the interesting question of what conditions must be imposed on a spacetime so that matter energy-momentum conservation can be expressed as an ordinary ‘continuity equation’ for a vector current, which is the mathematical vehicle for the notion of ‘local conservation’. In Minkowski spacetime without gravity, one defines the local *density* of four-momentum as $J_u^a = -u_b T^{ab}$, associated to a field of local observers with four-velocity u^a . If all four-velocities are parallel, so that the test observers are at relative rest with respect to one another, we have $\partial_a u_b = 0$ and the so-defined current is conserved, $\partial_a J_u^a = 0$. The analog in curved spacetime is the covariant conservation $\nabla_\mu J_\xi^\mu = 0$ for

$$J_\xi^\mu = -T^{\mu\nu} \xi_\nu , \quad (5.24)$$

defined in terms of some vector field ξ^μ which would be somehow related to a set of physical observers. Now, given $\nabla_\mu T^{\mu\nu} = 0$, the current J_ξ^μ is conserved if and only if the vector field satisfies

$$\nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu = 0 . \quad (5.25)$$

Such a vector field is known as a Killing field. According to equation (5.20), the Killing condition (5.25) coincides with the vanishing of the Lie derivative of the metric along the directions selected

⁵Notice the negative sign, due to the attractive nature of gravity, in contrast with the analogous expression in electrostatics.

at each point by the vector ξ^μ . In general $\xi^\mu \xi_\mu \neq -1$, so that we cannot claim that the Killing vector is some physical four-velocity. Rather, a standard observer moving along the integral curves of a timelike Killing vector, has four-velocity $u^\mu = \xi^\mu / \sqrt{-\xi^2}$. At any rate, the covariant conservation of J_ξ^μ defined by (5.24) does hold for any Killing vector field, irrespective of whether ξ^μ is timelike, null or spacelike. A metric supporting a Killing field is said to have an *isometry* in the ξ -directions.

The geometrical meaning of an isometry is that of ‘rigidity’ of the metric structure under motions within the manifold. Let us consider the integral curves generated by the vector field ξ^μ , that is to say, the solutions of the equation

$$\frac{dx^\mu}{d\lambda} = \xi^\mu(x(\lambda)) , \quad (5.26)$$

determining curves $x^\mu(\lambda)$ as a function of the parameter λ . We may use these integral curves as part of a particular coordinate chart ‘adapted’ to the vector field ξ^μ . In doing so, we can write $\xi^\mu \partial_\mu = \partial_\lambda$ and use λ as one of the coordinates. In this frame we have $\xi^\mu = \delta_\lambda^\mu$ and the Killing equation collapses to $\partial_\lambda g_{\mu\nu} = 0$. Therefore, existence of a Killing field is tantamount to the existence of a set of curves, the integral curves, along which the metric (aka the geometry) does not change. The intuition is that of a flat plane, or Minkowski spacetime for that matter, which are invariant under translations, rotations or boosts, each independent isometric motion being associated to a Killing vector. For a curved manifold, the standard example is a round sphere, which is isometric with respect to rotation around any polar axis.

In order to elucidate further the notion of isometry, let us take a more physical attitude, using particles as a probe of the spacetime structure. Working with the linearized version of the action for simplicity,

$$S_P = \frac{1}{2} m \int d\tau (g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu - 1) , \quad (5.27)$$

we identify symmetries of the geometry as symmetries of the particle action for *arbitrary histories* of the particle. Infinitesimally, the motion along the integral curves solving (5.26) is $x^\mu \rightarrow x^\mu + \lambda \xi^\mu(x) + O(\lambda^2)$. We now give a physical definition of an *isometry* as those vector-field flows that leave the particle action invariant, i.e. these transformations leave the metric ‘rigid’ as far as particle propagation is concerned. Since the action itself is invariant under reparametrizations, this criterion is independent of any coordinate system that we might use.

The variation of the particle action under the transformation $\delta_\lambda x^\mu = \lambda \xi^\mu$ is

$$\delta_\lambda S_P = \frac{1}{2} m \lambda \int d\tau (\mathcal{L}_\xi g_{\mu\nu}) \dot{x}^\mu \dot{x}^\nu , \quad (5.28)$$

where the integrand is controlled by the *Lie derivative* of the metric

$$\mathcal{L}_\xi g_{\mu\nu} \equiv \xi^\alpha \partial_\alpha g_{\mu\nu} + \partial_\mu \xi^\alpha g_{\alpha\nu} + \partial_\nu \xi^\alpha g_{\alpha\mu} = \nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu . \quad (5.29)$$

Hence, the condition on the metric for the action to be invariant is that ξ^α is a *Killing* vector field, satisfying (5.25).

In this case, we can define a conserved quantity along physical trajectories using the standard formula for the Noether charge,

$$Q_\xi = -\frac{\partial \mathcal{L}_P}{\partial \dot{x}^\mu} \xi^\mu = -m g_{\mu\nu} \dot{x}^\mu \xi^\nu = -p_\mu \xi^\mu . \quad (5.30)$$

The constancy of $Q_\xi = -p_\mu \xi^\mu$ along timelike geodesics generalizes to the case of null geodesics, where $Q_\xi = -k \cdot \xi$ and k^μ is the vector tangent to the light ray trajectory. In fact, it generalizes to any geodesic with tangent T^μ , even spacelike ones, for $Q_\xi = T_\mu \xi^\mu$ is covariantly constant along the geodesic,

$$\nabla_T(\xi \cdot T) = T^\alpha \nabla_\alpha (\xi_\mu T^\mu) = T^\alpha T^\mu \nabla_\alpha \xi_\mu + \xi_\mu T^\alpha \nabla_\alpha T^\mu = 0, \quad (5.31)$$

the first term vanishing because of the Killing condition and the second because of the geodesic condition.

Hence, for spacetimes with special symmetries, local conservation of matter energy-momentum still holds. There are locally conserved energy-momentum currents constructed out of the matter energy-momentum tensor and suitable projections of four-momenta of particles are also conserved along their free-fall paths. It should be understood that the requirement of a symmetry on a manifold implies a corresponding stringent requirement on the distribution and motion of matter in this spacetime, because matter dynamics and spacetime geometry are linked locally through Einstein's equations.

5.2 Finding Solutions

Exact solutions of Einstein's equations with a definite physical interpretation are hard to come by. In a sense, any metric can be regarded as a solution, provided we 'invent' an exotic energy-momentum tensor that fulfills the right hand side of the equations. In fact, the Bianchi identity ensures that such energy-momentum tensor is at least conserved. However, it will have unphysical features, such as locally measurable negative energy density. If one postulates that $T_{\mu\nu}$ is computed from some standard matter model of particles, fluids or field theories, then no general methods exist to find exact solutions of the gravitational equations, due to their strong non-linearity. Numerical analysis is always an option, and one that has gained a lot of interest in recent times, due to the wealth of new data about black-hole collisions as 'heard' by gravitational wave detectors.

On the analytic side, one can simplify matters by looking for solutions with some prescribed symmetry that is interesting for physical reasons. For example, in the context of cosmological models, we seek metrics with the properties of *homogeneity* and *isotropy* which appear to be satisfied by the large-scale structure of the observable universe. For problems involving motion in the solar system, we are interested in metrics with axial and/or spherical symmetry.

5.2.1 More on Isometries

We have discussed in the previous section the basic diagnostic for a geometrical symmetry, namely the existence of a Killing vector. The main consequence for the simplification of Einstein equations is that one can introduce special coordinate systems in which the metric depends on fewer variables. Adapting the coordinates to a Killing vector, so that $\xi^\mu \partial_\mu = \partial_\lambda$, results in the metric being λ -independent. For example, a metric is called *stationary* if it has a timelike Killing vector. Choosing the time coordinate adapted to this Killing vector, we have $\xi = (1, 0, 0, 0)$ in this frame, and $\partial_t g_{\mu\nu} = 0$, hence

$$ds_{\text{stationary}}^2 = g_{00}(\vec{x}) dt^2 + 2 \sum_i g_{0i}(\vec{x}) dx^i dt + \sum_{ij} g_{ij}(\vec{x}) dx^i dx^j. \quad (5.32)$$

Furthermore, a stationary metric is called *static* if it is invariant under time reversal $t \rightarrow -t$. This leads to

$$ds_{\text{static}}^2 = g_{00}(\vec{x}) dt^2 + \sum_{ij} g_{ij}(\vec{x}) dx^i dx^j . \quad (5.33)$$

Analogous simplifications are available when the metric has a spacelike or null Killing vector.

If a metric has more than one Killing vector, one can consider the algebra of transformations that they form under commutation. To define this notion, we consider the effect of the translation along integral curves of a vector field ξ^μ when acting on arbitrary scalar functions:

$$\delta_\xi f(x) = \mathcal{L}_\xi f(x) = \xi^\mu \partial_\mu f(x) . \quad (5.34)$$

Then, one can write

$$[\delta_\xi, \delta_\eta] f(x) = \delta_{[\xi, \eta]} f(x) , \quad (5.35)$$

for any function, provided we define the so-called *Lie bracket* as ⁶

$$[\xi, \eta]^\mu = \xi^\alpha \partial_\alpha \eta^\mu - \eta^\alpha \partial_\alpha \xi^\mu . \quad (5.36)$$

This commutator algebra is very transparent if we adopt the view that vector fields expressed as $\xi = \xi^\mu \partial_\mu$ are literally interpreted as differential operators acting on scalar functions. Hence, if a coordinate system can be defined as adapted to two Killing vectors simultaneously, it is clear that their commutator vanishes because, in adapted coordinates $\xi = \partial_\lambda$, $\eta = \partial_\zeta$, one has $[\delta_\xi, \delta_\eta] f = [\partial_\lambda, \partial_\zeta] f = 0$ for any smooth scalar $f(x)$. The converse is also true, i.e. the necessary and sufficient condition for the integral curves of two vector fields to coordinate a two-dimensional submanifold is that they commute. In more detail, a theorem by Frobenius states that the integral curves of a set of vector fields $\xi_{(a)}$ are contained in a submanifold if and only if their Lie brackets close on themselves, $[\xi_{(a)}, \xi_{(b)}] = \sum_c f_{ab}^c(x) \xi_{(c)}$, with coefficients $f_{ab}^c(x)$ that may be functions (constant coefficients correspond to the Lie algebra of a group of isometries). Moreover, these integral curves are good coordinates when they form an abelian subalgebra.

Problem: Poincaré Isometries

Find all the Killing vector fields of Minkowski spacetime and recover the Poincaré group as the associated Lie algebra under commutation through the Lie brackets of the Killing vector fields.

Non-abelian isometry groups do not give such simple rules as abelian ones for the metric components. In general, for non-abelian isometries, one must list all possible invariants of the coordinates and the differentials dx^a and restrict the metric accordingly. As an example, look at a static spacetime (5.33) and impose rotation $SO(3)$ symmetry of the spatial sections. If we let the rotation group $SO(3)$ act on $\vec{x} \in \mathbb{R}^3$ in the standard fashion, the basic invariants are $\vec{x}^2$, $d\vec{x}^2$ and $\vec{x} \cdot d\vec{x}$. So the general *ansatz* for a static, spherically symmetric metric is

$$ds^2 = -F(\rho) dt^2 + D(\rho) (\vec{x} \cdot d\vec{x})^2 + C(\rho) d\vec{x}^2 , \quad (5.37)$$

⁶It is non-trivial but easy to check that this operation always defines a Diff vector.

where $\rho^2 = \vec{x}^2$ and $F(\rho), D(\rho), C(\rho)$ are general functions. Picking spherical coordinates on $\mathbb{R}^3$ we have $d\vec{x}^2 = d\rho^2 + \rho^2 d\Omega^2$ and $(\vec{x} \cdot d\vec{x})^2 = \rho^2 d\rho^2$, so that combining all terms we end up with a form

$$ds^2 = -F(\rho)dt^2 + G(\rho)d\rho^2 + H(\rho)d\Omega^2, \quad (5.38)$$

where $d\Omega^2 = d\ell_{S^2}^2 = d\theta^2 + \sin^2 \theta d\phi^2$. We may now change coordinates to $r^2 = H(\rho)$ so that the area of fixed radius S^2 spheres is $4\pi r^2$. So the final form of the static and $SO(3)$ -symmetric *ansatz* is

$$ds^2 = -A(r)dt^2 + B(r)dr^2 + r^2 d\Omega^2. \quad (5.39)$$

A full classification of standard metrics for different isometry groups is beyond the scope of these notes. We shall content ourselves with introducing the required forms as they are needed by the physics discussion at hand. In general, if we have sufficient symmetry to leave just one nontrivial coordinate, such as the situation in (5.39), the Einstein partial differential equations may simplify into a set of coupled ordinary differential equations, which are much easier to solve.

5.2.2 Spherically Symmetric Vacuum Solutions

With the *ansatz* (5.39) in place, $ds^2 = -A(r)dt^2 + B(r)dr^2 + r^2 d\Omega^2$, we may look for spherically symmetric and static solutions of the vacuum equations,

$$R_{\mu\nu} = \Lambda g_{\mu\nu}. \quad (5.40)$$

Upon direct calculation, the components of the Ricci tensor for this metric read

$$\begin{aligned} R_{tt} &= \frac{A''}{2B} - \frac{A'}{4B} \left(\frac{A'}{A} + \frac{B'}{B} \right) + \frac{A'}{rB} \\ R_{rr} &= -\frac{A''}{2A} + \frac{A'}{4A} \left(\frac{A'}{A} + \frac{B'}{B} \right) + \frac{B'}{rB} \\ R_{\theta\theta} &= 1 - \frac{r}{2B} \left(\frac{A'}{A} - \frac{B'}{B} \right) - \frac{1}{B} \\ R_{\phi\phi} &= \sin^2 \theta R_{\theta\theta}, \end{aligned} \quad (5.41)$$

where the primes denote derivative with respect to r .⁷

The *ansatz* (5.39) obtains the Minkowski metric in polar coordinates

$$ds^2 = -dt^2 + dr^2 + r^2 d\Omega^2$$

for $A = B = 1$. This is the trivial vacuum solution with vanishing cosmological constant $\Lambda = 0$. It is not too difficult to generalize the Minkowski solution to vacuum spacetimes with positive or negative cosmological constant. Since Λ defines a length scale, let us denote $|\Lambda| = 3/R^2$, where R is the radius at which the effects of the cosmological constant are felt.

⁷The fastest way of obtaining this result is to compute the Christoffel symbols by variation of the particle action in the metric *ansatz*:

$$S_P = \frac{1}{2} m \int d\tau (g_{\mu\nu} \dot{x}^\mu \dot{x}^\nu - 1) = \frac{1}{2} m \int d\tau \left(-A(r) \dot{t}^2 + B(r) \dot{r}^2 + r^2 (\dot{\theta}^2 + \sin^2 \theta \dot{\phi}^2) - 1 \right),$$

and match to the geodesic equation. This produces naturally all the non-vanishing Christoffel components. Then, one must compute explicitly the Ricci tensor from these connection coefficients.

Inserting the *ansatz* (5.39) in the vacuum Einstein equations, $R_{\mu\nu} = \Lambda g_{\mu\nu}$, we find

$$\frac{R_{rr}}{B} + \frac{R_{tt}}{A} = \frac{1}{rB} \left(\frac{A'}{A} + \frac{B'}{B} \right) = 0 ,$$

which implies $A(r)B(r) = \text{constant}$. Since $A(0) = B(0) = 1$ to get back Minkowski spacetime at short distances, we find $A(r) = 1/B(r)$. Inserting this result now into $R_{\theta\theta} = \Lambda g_{\theta\theta} = r^2 \Lambda$ we obtain

$$(rA)' = 1 - \Lambda r^2 ,$$

which, together with the boundary condition at $r = 0$, is solved by $A(r) = 1 - \Lambda r^2/3$. These metrics

$$ds_\Lambda^2 = -dt^2 \left(1 - \frac{1}{3} \Lambda r^2 \right) + \frac{dr^2}{1 - \frac{1}{3} \Lambda r^2} + r^2 d\Omega^2 \quad (5.42)$$

are known as de Sitter spacetime (dS) for $\Lambda > 0$ and Anti de Sitter spacetime (AdS) for $\Lambda < 0$.

AdS is a smooth non-compact spacetime of constant negative curvature. Defining a new radial variable ρ as $r = R \sinh(\rho/R)$, we learn that the spatial sections at constant t have 3-metric

$$ds_{\mathbf{H}^3}^2 = d\rho^2 + R^2 \sinh^2(\rho/R) d\Omega^2 ,$$

which is the three-dimensional generalization of the famous Lobachevsky (hyperbolic) plane, a space in which volumes enclosed by large spheres scale like the surface area.

The dS metric is singular at $r = R = \sqrt{3/\Lambda}$, but this is only a singularity of the particular frame used to write down the metric. Clearly, the Ricci curvature stays constant throughout all dS spacetime, and one can explicitly show that the dS metric can be analytically extended beyond $r = R$. To see this, define

$$z_0 = \sqrt{R^2 - r^2} \sinh(t/R) , \quad z_1 = \sqrt{R^2 - r^2} \cosh(t/R) ,$$

and let $z_2^2 + z_3^2 + z_4^2 = r^2$ parametrize an S^2 of radius r embedded in $\mathbb{R}^3$. Then, it is easy to see that dS spacetime is the hyperboloid

$$-z_0^2 + z_1^2 + z_2^2 + z_3^2 + z_4^2 = R^2$$

embedded in five-dimensional $\mathbb{R}^5$ with Minkowski metric

$$ds_{\mathbb{R}^5}^2 = -dz_0^2 + dz_1^2 + dz_2^2 + dz_3^2 + dz_4^2 .$$

Hence, we see that dS is a homogeneous spacetime without singularities. In this description, it is evident that the full isometry group of dS is the Lorentz group in five dimensions, namely $O(1, 4)$.

There is another intrinsic representation of the dS metric that covers the entire manifold. Setting

$$z_0 = R \sinh(t'/R) , \quad z_k = R \cosh(t'/R) N_k , k = 1, 2, 3, 4 ,$$

where N_k is a unit vector in $\mathbb{R}^4$, parametrizing an S^3 of unit radius, we have

$$ds_{\text{dS}}^2 = -dt'^2 + R^2 \cosh^2(t'/R) d\ell_{S^3}^2 . \quad (5.43)$$

In this form, the dS metric looks very different from (5.42). It consists of a 3-sphere that contracts from the infinite past to a radius R and then re-expands again into the future with an asymptotically exponential rate. Hence, dS space is a *cosmological* spacetime, something clearly surprising if we are given only the (5.42) static form.⁸

Problem: May “The Force” Be with You

Study the radial geodesics of massive test particles in the vacua with nonvanishing cosmological constant. Show that Λ behaves as a repulsive cosmic potential for dS and as a “confining” cosmic potential for AdS.

5.2.3 The Schwarzschild Solution

The Schwarzschild metric is the unique asymptotically flat vacuum solution with $SO(3)$ symmetry and static. It is therefore the appropriate metric for the exterior of a spherically symmetric and time independent matter distribution. In fact, it turns out that the static condition is redundant, as proved by Birkhoff, all vacuum solutions of $SO(3)$ isometry group are static (there is no purely monopole gravitational radiation). This result is entirely analogous to its electromagnetic cousin.

Plugging the *ansatz* (5.39) into the vacuum Einstein equation $R_{\mu\nu} = 0$ one finds

$$\frac{R_{rr}}{B} + \frac{R_{tt}}{A} = \frac{1}{rB} \left(\frac{A'}{A} + \frac{B'}{B} \right) = 0, \quad (5.44)$$

where primes stand for radial derivatives. From here, we obtain $A(r)B(r) = \text{constant}$. This much is the same as in the discussion of isotropic vacuum solutions but now, instead of demanding smoothness at the origin of polar coordinates, we require that the metric approaches Minkowski at infinity, so that we get, again, $A(r)B(r) = 1$. Now, it is enough to concentrate on the $R_{\theta\theta}$ equation, that reads

$$R_{\theta\theta} = 1 - A'(r)r - A(r) = 0,$$

with solution $rA(r) = r + \text{constant}$. Since we must approach the Newtonian form $g_{tt} \rightarrow -1 + 2GM/r + \dots$ at large radius, this is enough to fix the complete metric as

$$ds^2 = - \left(1 - \frac{2GM}{r} \right) dt^2 + \left(1 - \frac{2GM}{r} \right)^{-1} dr^2 + r^2 d\Omega^2, \quad (5.45)$$

where M accounts for the total energy contained in the spacetime.⁹

⁸Originally, de Sitter found the static form (5.42), describing only one patch of the full dS spacetime. Despite this, Hubble did incorporate a cosmological interpretation of the static dS metric in his fits of galactic recession velocities.

⁹It is also possible to generalize the asymptotically flat Schwarzschild solution to analogous solutions with dS or AdS asymptotics. The *naive* answer obtained in the limit $(GM)^2\Lambda \ll 1$ turns out to be exact, i.e. $A = B^{-1} = 1 - \frac{1}{3}\Lambda r^2 - \frac{2GM}{r}$.

Problem: Birkhoff Theorem

Show that the time-independence of the metric *ansatz* for the Schwarzschild solution is redundant, namely *vacuum* solutions with $SO(3)$ symmetry are necessarily static.

The metric components are singular at the so-called Schwarzschild radius $r_s = 2GM$. For standard astrophysical bodies, such as stars and planets, one has $r_s \approx 3(M/M_\odot)$ km, so that this singularity is irrelevant in many applications, since the Schwarzschild solution must be matched to some ‘interior’ solution at the surface of the body. Any object for which the vacuum Schwarzschild solution would apply down to $r = r_s$ is called a *black hole*.

The behavior of the $r = r_s$ surface is similar to the analogous surface in de Sitter space, the singularity being just an artifact of the particular coordinate system. A freely falling observer is only sensitive to the tidal forces, which are controlled by the Riemann tensor. By direct calculation, it is possible to check that all the non-vanishing components of the Riemann tensor in the vicinity of $r = r_s$ are of order $1/r_s^2$. Therefore, tidal accelerations have the expected value inferred by extrapolation from Newtonian scaling: for an object of length ℓ , $a(\ell)_{\text{tidal}} \sim GM\ell/r_s^3 \sim \ell/r_s^2$. This shows that an infalling observer will not notice any singular behavior upon crossing the surface $r = r_s$. This crossing occurs in finite proper time. The proper time to radially fall from some initial radius $r_0 > r_s$ down to r_s is

$$\Delta\tau = \int_{r_s}^{r_0} \frac{dr}{\sqrt{\frac{r_s}{r} - \frac{r_s}{r_0}}} < \infty.$$

Hence, the proper time for radial fall is finite, as well as the tidal forces at $r = r_s$, and there is no physical singularity.

5.3 The Classic Tests

With the Schwarzschild solution at hand, we can now discuss the three classic tests of General Relativity. We demonstrate the advance of perihelia, the bending of light and the gravitational red shift effect in the background of the Schwarzschild solution. We also include a fourth ‘classic’ test for radar echo delay in the Solar System. Today, the precision measurements based on these effects confirm the theory down to the 10^{-5} level.

5.3.1 Orbital Precession

Planets will be approximated as pointlike particles of mass m following free-fall geodesics in the Schwarzschild metric. In strict analogy with Kepler’s problem, this system can be completely integrated once the symmetries are taken into account. As a consequence of the spherical symmetry, particle orbits lie on a plane, which can be regarded as equatorial. Restricted to this plane, the spacetime enjoys two commuting Killing vectors: the temporal one $\xi_t = \partial/\partial t$ and the polar one $\xi_\phi = \partial/\partial\phi$. The associated Noether charges are the energy and angular momentum

$$E = -p_\alpha \xi_t^\alpha = m(1 - 2GM/r) \dot{t}, \quad L = p_\alpha \xi_\phi^\alpha = mr^2 \dot{\phi}. \quad (5.46)$$

Defining the energy and angular momentum per unit mass $\varepsilon \equiv E/m$, $\ell = L/m$, we have, from the dispersion relation $p_\alpha p^\alpha + m^2 = 0$, or $u^2 = -1$,

$$-1 = -\frac{\varepsilon^2}{1 - 2GM/r} + \frac{\dot{r}^2}{1 - 2GM/r} + \frac{\ell^2}{r^2} . \quad (5.47)$$

This equation is equivalent to the classical one-dimensional motion of a unit-mass particle of energy per unit mass equal to $\varepsilon_{\text{eff}} = (\varepsilon^2 - 1)/2$, subject to the effective potential

$$V_{\text{eff}} = -\frac{GM}{r} + \frac{\ell^2}{2r^2} - \frac{GM\ell^2}{r^3} . \quad (5.48)$$

So that $\dot{r}$ can be solved from

$$\frac{1}{2} \dot{r}^2 + V_{\text{eff}}(r) = \varepsilon_{\text{eff}} . \quad (5.49)$$

Compared to the Newtonian problem, the crucial new feature of this potential is the last term $-GM\ell^2/r^3$ which dominates over the Newtonian one at small enough radii, and is negative definite, representing a clear tendency for the orbits to be destabilized towards small radius. The centrifugal barrier is no longer infinite. In fact, it is not present at all if $\ell^2 < 12(GM)^2$. In this case, any particle on a trapped orbit, $\varepsilon^2 < 1$ will fall to the Schwarzschild radius.¹⁰




Figure 5.1: *Effective potential for massive particles with $\ell^2 > 12(GM)^2$. For $\ell^2 < 12(GM)^2$ the centrifugal barrier is gone.*

For $\ell^2 > 12(GM)^2$ the potential has a stable minimum at R_+ and an unstable maximum at R_- , with

$$R_{\pm} = \frac{\ell^2 \pm \sqrt{\ell^4 - 12(\ell GM)^2}}{2GM} . \quad (5.50)$$

Thus, stable circular orbits exist at R_+ , which is restricted to be greater than $6GM$. Unstable circular orbits exist at R_- which is restricted to $3GM < R_- < 6GM$. Any particle in a circular orbit has angular momentum per unit mass

$$\ell = \sqrt{\frac{GMR_{\pm}^2}{R_{\pm} - 3GM}} , \quad (5.51)$$

¹⁰Notice that the effective energy ε_{eff} is nothing but the Newtonian energy per unit mass in the nonrelativistic limit.

and energy per unit mass equal to (obtained setting $\dot{r} = 0$),

$$\varepsilon = \frac{R_{\pm} - 2GM}{\sqrt{R_{\pm}(R_{\pm} - 3GM)}} . \quad (5.52)$$

For $3GM < R_- < 4GM$, ε ranges between ∞ and unity, so that any particle in unstable circular orbits with $R_- < 4GM$ will escape to infinity if perturbed outwards.

Regarding the stable circular orbits, the lowest possible one (known as the ISCO) sits at $R_+ = 6GM$, and its binding energy per unit mass is

$$\varepsilon_b = 1 - \varepsilon = 1 - \sqrt{8/9} \approx 0.06 . \quad (5.53)$$

Hence, particles in circular orbits at very large radius, with $\varepsilon \approx 1$, will steadily lose energy and angular momentum by gravitational radiation emission. Gradually, they will spiral down to the last possible circular orbit at $R_+ = 6GM$, and then fall rapidly to the Schwarzschild radius. In the quasistatic process of gravitational radiation emission, up to six per cent of the initial energy is radiated, a quantity which is not negligible. So, we see that gravitational radiation, being cumulative, can have significant effects over time, even if the radiation power is so small. If the radiated energy is mediated by electromagnetic interactions, as happens through viscous (turbulent) friction in accretion disks, the release of energy with such efficiencies is much more powerful than thermonuclear reactions in stars.

The parametric equation for the orbits can be obtained by writing $\dot{r} = \sqrt{2(\varepsilon_{\text{eff}} - V_{\text{eff}})}$ and dividing by $\dot{\phi} = L/mr^2$ to obtain

$$\Delta\phi = \int_{r_i}^{r_f} \frac{d\phi}{dr} dr = \int_{r_i}^{r_f} \frac{Ldr}{\sqrt{2m^2r^4(\varepsilon_{\text{eff}} - V_{\text{eff}})}} . \quad (5.54)$$

Setting r_i and r_f at the turning points, which solve $V_{\text{eff}}(r_{\pm}) = \varepsilon_{\text{eff}}$, computes the perihelion advance for closed orbits. We can derive a simple approximation to the orbital precession in the case of nearly circular orbits (small eccentricity). For a nearly circular orbit, we can study the harmonic approximation to the potential near R_+ , and we find a characteristic frequency

$$\omega_r^2 = V_{\text{eff}}''(R_+) = \frac{GM(R_+ - 6GM)}{R_+^3(R_+ - 3GM)} . \quad (5.55)$$

On the other hand, the angular frequency per revolution in the polar angle ϕ is

$$\omega_{\phi}^2 = \frac{\ell^2}{R_+^4} = \frac{GM}{R_+^2(R_+ - 3GM)} . \quad (5.56)$$

The inequality of these two frequencies translates into a precession of the orbit. The precession frequency being simply the difference

$$\omega_p = \omega_{\phi} - \omega_r = \omega_{\phi} \left(1 - \sqrt{1 - 6GM/R_+} \right) . \quad (5.57)$$

To lowest non-vanishing order for large R_+ , we find

$$\omega_p \approx \frac{3(GM)^{3/2}}{R_+^{5/2}} , \quad (5.58)$$

or, in terms of the angular advance per period $T_{\text{orb}} \approx 2\pi R_+ \sqrt{R_+/GM}$:

$$\Delta\phi_p \approx \frac{6\pi GM}{R_+} . \quad (5.59)$$

A direct evaluation of (5.54) for arbitrary orbits yields the general result

$$\Delta\phi_p \approx \frac{6\pi GM}{(1-e^2)a} , \quad (5.60)$$

in terms of the eccentricity e and the semimajor axis of the ellipse a . This gives the famous 43 seconds of arc per century for Mercury. For the equally famous binary pulsar of Hulse and Taylor, the precession is about 4 degrees per year!

Problem: Planetary Precession in a Wrong Theory of Gravity

Compute the precession of Mercury in a scalar theory of gravity and compare with the result in GR.

5.3.2 Gravitational Lensing

We discuss light (photon) propagation taking the $m \rightarrow 0$ limit of the particle propagation. In this limit the proper time becomes a bad parameter and we need to re-parametrize the trajectory in terms of an affine parameter or in terms of coordinate time. For the purposes of describing photon orbits we can avoid such choices by focusing on the parametric equation (5.54)

$$\frac{d\phi}{dr} = \left[\frac{2m^2 r^4}{L^2} (\varepsilon_{\text{eff}} - V_{\text{eff}}) \right]^{-1/2}$$

which does admit a non-singular $m \rightarrow 0$ limit. In this limit the photon angular momentum is given by $L_\gamma = |\vec{p}_\gamma| b = E_\gamma b$, where b is the impact parameter for unbounded orbits. The resulting photon parametric equation is

$$\frac{d\phi}{dr} = \left[\frac{r^4}{b^2} - r(r - 2GM) \right]^{-1/2} . \quad (5.61)$$

The expression in brackets vanishes at the turning points r_0 of the orbit, which are then localized at the largest positive root of the equation

$$r_0^3 - b^2(r_0 - 2GM) = 0 . \quad (5.62)$$

Such a root only exists for $b \geq b_c = 3^{3/2}GM$, which corresponds to the smallest possible impact parameter such that the photon is not captured, giving the capture cross section

$$\sigma_{\text{capture}} = \pi b_c^2 = 27\pi(GM)^2 . \quad (5.63)$$

It is interesting that the critical value of the impact parameter corresponds to a turning point $r_0 = 3GM$, which happens to be degenerate, in the sense that the photon admits a circular (albeit unstable) orbit at this radius.




Figure 5.2: Gravitational bending of a light ray grazing at a distance r_0 from a central field.

The light that is not captured will scatter off the gravitational field. The deflection of the light ray is defined by

$$\delta\phi = \Delta\phi - \pi, \quad (5.64)$$

with $\Delta\phi = \phi(\sigma = +\infty) - \phi(\sigma = -\infty)$.

So, the total $\Delta\phi$ is twice the amount from the turning point r_0 out to infinity:

$$\Delta\phi = 2 \int_{r_0}^{\infty} \frac{dr}{\sqrt{r^4 b^{-2} - r(r - 2GM)}}. \quad (5.65)$$

To estimate this, we change variables to $u = r_0/r$ and eliminate b via (5.62) to obtain

$$\Delta\phi = 2 \int_0^1 \frac{du}{\sqrt{1 - u^2 - \frac{2GM}{r_0}(1 - u^3)}}. \quad (5.66)$$

Expanding to leading order in GM/r_0 we finally obtain

$$\Delta\phi = \pi + \frac{4GM}{r_0} + \dots = \pi + \frac{4GM}{b} + \dots, \quad (5.67)$$

since the difference between r_0 and b is already of order GM/r_0 . Hence, the deflection of light is, to leading order

$$\delta\phi \approx \frac{4GM}{b}. \quad (5.68)$$

For starlight grazing the Sun, one obtains the famous 1.75 seconds of arc that were measured by Eddington in 1919 and that started Einstein's life as a pop icon.

Problem: Newtonian Light Bending

Compute the light bending effect in the purely Newtonian metric $ds_N^2 = -(1 + 2\phi_N)dt^2 + d\vec{x}^2$, a computation done by Einstein in 1911. How does the answer compare with GR and with the purely Newtonian answer that Newton himself would have obtained in his corpuscular theory of light?

Today, gravitational light bending is a routine tool for Astronomy. Heavy objects, such as a galaxy, sitting in between us and some distant source (another galaxy) can act as a gravitational




Figure 5.3: Gravitational lensing by a point mass M_L (angles exaggerated). (a) Side view. The source S is seen at an angle β from the optical axis through the lens. The lens is at distance D_L from the observer O , the source plane is at distance D_S , and $D_{LS} \approx D_S - D_L$. A light ray passing the lens at impact parameter $b \approx \theta_+ D_L$ is bent by the angle $\delta\phi = 4GM_L/b$ of (5.68). It reaches O from the direction θ_+ , so that S appears at the image position S_+ . The heights on the source plane give the lens equation $\theta D_S \approx \beta D_S + \delta\phi D_{LS}$. A second ray (gray) passes on the other side of the lens with a smaller impact parameter and a larger deflection, and produces the image S_- at $\theta_- < 0$. (b) The sky seen by O for $\beta = \theta_E/2$. The images of a small circular source (dotted) lie on opposite sides of the lens: at θ_+ , outside the Einstein ring of radius θ_E , and at θ_- , inside it. Both are stretched tangentially, with areas, relative to the source, approximately equal to the magnifications 1.59 and 0.59. For $\beta = 0$ they merge into the ring. (c) The image positions $\theta_{\pm} = \frac{1}{2}(\beta \pm \sqrt{\beta^2 + 4\theta_E^2})$ as functions of β , with $\theta_+ \theta_- = -\theta_E^2$. The dots mark the case shown in (b). For $\beta \gg \theta_E$ the primary image approaches the true position, $\theta_+ \rightarrow \beta$, while the secondary one approaches the lens and fades.

lens. Consider the geometry of the figure, featuring a source at distance D_S and a lens of mass M_L at distance D_L , slightly displaced a small angle β from the line of sight of the source, which is itself seen at an angle θ with respect to the line of the lens. Then, the impact parameter for the bent light ray is $b \approx \theta D_L$ and from the figure we find, in the approximation that all angles are very small

$$\theta D_S \approx \beta D_S + \delta\phi D_{LS} , \quad (5.69)$$

where $D_{LS} \approx D_S - D_L$ is the lens-source distance and $\delta\phi$ is the GR deflection angle (5.68). Using $\delta\phi \approx 4GM_L/b \approx 4GM_L/\theta D_L$, we may solve for θ as

$$\theta_{\pm} = \frac{1}{2} \left(\beta \pm \sqrt{\beta^2 + 4\theta_E^2} \right) , \quad (5.70)$$

where

$$\theta_E = \sqrt{\frac{4GM_L D_{LS}}{D_L D_S}} \quad (5.71)$$

is called the *Einstein angle*. If the source and the lens are approximated as pointlike and are exactly aligned, one sees a ring subtended by the Einstein angle θ_E . If source and lens are slightly misaligned one sees two distorted images around the angles $\theta_{\pm}$ in (5.70). When the lens has a more complicated structure, the lensing produces multiple images in complicated deformation patterns. If one can measure θ_E and estimate the distances to the source and the lens, one can obtain a measurement of M_L . For typical cosmological situations of galaxies lensing background galaxies, one has $\theta_E = O(1)$ arcseconds, perfectly detectable with modern telescopes. For compact objects of $M_L \sim M_{\odot}$ lensing stars within galactic scales, one has Einstein angles a thousand times smaller, not directly detectable. However, the slight increase in brightness caused by the focusing *is* detectable, a situation that is described as *microlensing* and can be used to scan statistically for such compact dark objects acting as lenses. Since those objects have characteristic velocities in the galaxy in the range of the average orbital velocity, about 200 km/s, the crossing time at a typical distance $D_L \sim 10$ kpc is about

$$\Delta t_{\text{crossing}} \sim \frac{\theta_E D_L}{v_L} \sim 10^{-3} \text{ arcsec} \frac{10 \text{ kpc}}{200 \text{ km/s}} \sim 0.2 \text{ years} . \quad (5.72)$$

Therefore, month-long peaks in luminosity are signals of microlensing by such compact objects within the galaxy.

5.3.3 Gravitational Redshift

In a general spacetime with metric $g_{\alpha\beta}$ the locally measured frequency of light depends on the point of measurement, as well as the state of motion of the observer. An invariant definition is given by the expression (2.88)

$$\omega = -k_{\mu} u^{\mu} , \quad (5.73)$$

where k_{μ} is the wave vector of the light ray and u^{μ} is the four-velocity of the observer. Hence, in order to relate the measured frequency at two different events, one typically needs to solve for the null geodesic of the light path between them. Things are greatly simplified if the observers' trajectories are tangent to a Killing vector. In general, given a Killing vector ξ and a tangent vector T^a to some geodesic curve, the contraction $\xi_{\mu} T^{\mu}$ is constant along the geodesic (cf. (5.31)). Applying this lemma to a pair of static observers with four-velocity parallel to the

temporal Killing vector of Schwarzschild, $\xi = \partial/\partial t$, we find that $\xi_\alpha k^\alpha = \text{constant}$ along the light path and therefore is the same at emission and reception. The four-velocity of static observers is normalized by definition $u^\alpha u_\alpha = -1$, so

$$u^\alpha = \frac{\xi^\alpha}{\sqrt{-\xi_\mu \xi^\mu}} . \quad (5.74)$$

From $(\xi \cdot k)_{\text{emission}} = (\xi \cdot k)_{\text{reception}}$ we deduce

$$\frac{\omega_{\text{em}}}{\omega_{\text{rec}}} = \frac{(-\xi^2)_{\text{rec}}^{1/2}}{(-\xi^2)_{\text{em}}^{1/2}} = \sqrt{\frac{1 - 2GM/r_{\text{rec}}}{1 - 2GM/r_{\text{em}}}} . \quad (5.75)$$

Hence, light redshifts as it climbs through the gravitational field. For small trips of height $h = |r_{\text{rec}} - r_{\text{em}}|$ one finds $\Delta\omega/\omega \approx |\vec{g}|h/c^2$, an effect of order 10^{-15} that was tested by Pound and Rebka in 1959, with $h \approx 22.5$ m, the height of the tower of the Jefferson Physical Laboratory at Harvard University. Today, accurate consideration of this effect is essential for the correct performance of GPS systems.

Problem: Those Twins Again

Consider our proverbial twins again, but now the traveling twin embarks on a trip around a trapped solar orbit with initial velocity very close to the escape velocity from the Sun. After a very long time he falls back to Earth. Who is younger?

5.3.4 Echo Delay

The unofficial ‘fourth classic test’ of General Relativity is Shapiro’s echo delay effect, proposed by Shapiro in 1964 and first tested in the late 1960’s by bouncing radar pulses off Mercury and Venus (and later off interplanetary spacecraft). The idea is that lensed electromagnetic signals (or any other massless carrier of information) passing near a massive body take longer time than they would in flat spacetime.

For any static spacetime null geodesics satisfy

$$ds^2 = 0 = g_{00}(\vec{x})dt^2 + \sum_{ij} g_{ij}(\vec{x}) dx^i dx^j , \quad (5.76)$$

Assuming that g_{00} does not vanish in the region of consideration, the null geodesics also satisfy

$$\tilde{ds}^2 = 0 = -dt^2 + \sum_{ij} \tilde{g}_{ij}(\vec{x}) dx^i dx^j , \quad (5.77)$$

where $\tilde{g}_{ij} \equiv g_{ij}/|g_{00}|$ is the so-called *optical* or *Fermat metric*. It is a 3-metric of Euclidean signature whose geodesics give the *spatial* trajectories of light rays. In particular, the coordinate time lapse taken to go along some geodesic curve C is

$$t_C = \int_C \sqrt{\tilde{g}_{ij} dx^i dx^j} . \quad (5.78)$$




Figure 5.4: Arrangement for the test of the echo time delay.

Now, it is clear that a null geodesic that grazes the Schwarzschild radius (coming close to a black hole) will encounter very small values of g_{00} , so that Fermat metric is greatly stretched in these near-horizon regions and the signal will be greatly delayed, compared to the answer in flat space.

To be more quantitative, in the Schwarzschild metric we can pass to the so-called *isotropic* radial coordinate defined by

$$r = \rho \left(1 + \frac{GM}{2\rho} \right)^2, \quad (5.79)$$

in terms of which the Schwarzschild metric reads

$$ds^2 = - \left(\frac{1 - \frac{GM}{2\rho}}{1 + \frac{GM}{2\rho}} \right)^2 dt^2 + \left(1 + \frac{GM}{2\rho} \right)^4 (d\rho^2 + \rho^2 d\Omega^2). \quad (5.80)$$

This leads to a Fermat metric

$$\tilde{ds}^2 = -dt^2 + n(\vec{x})^2 d\vec{x}^2, \quad n(\vec{x}) = \frac{\left(1 + \frac{GM}{2|\vec{x}|} \right)^3}{\left(1 - \frac{GM}{2|\vec{x}|} \right)}, \quad (5.81)$$

where we have written $d\vec{x}^2 = d\rho^2 + \rho^2 d\Omega^2$. In the metric (5.81) the Fermat geodesics C are straight lines in $\vec{x}$ coordinates, to leading order in GM . Denoting by ℓ a length parameter along these straight lines, the elapsed time becomes

$$t_C = \int_C d\ell n(\vec{x}(\ell)), \quad (5.82)$$

showing that $n(\vec{x})$ behaves as an effective ‘refraction index’. Since $n(\vec{x})$ approaches unity far away from the gravitating body and it is otherwise larger than unity, any signal that approaches

a massive body on its way suffers a time delay with respect to what it would be in flat spacetime. If we have a signal between two points in near opposition at radii R_A and R_B from the gravitating body of mass M , and the distance of closest approach of the null geodesic with the central body is R_0 , we have a delay

$$\Delta t = \int_C d\ell \left(n(\vec{x}(\ell)) - 1 \right) \approx 2GM \log \left(\frac{4R_A R_B}{R_0^2} + 1 \right), \quad (5.83)$$

on doing the integral in the regime $GM \ll R_0 \ll R_{A,B}$. In an echo experiment where the signal is sent from A to B and back, the delay comes with an extra factor of two.¹¹

¹¹We are quoting in (5.83) the time delay in the t coordinate. A real clock at point A would tick to proper time rather than coordinate time. The two are related by the usual factor of $\sqrt{-g_{00}(A)} = \sqrt{1 - 2GM/R_A}$, which we may neglect when A is far away from the gravitating source.

Chapter 6

Faraway GR

Despite the fact that the long-distance physics of gravity is dominated by the cosmological constant term, Λ , whose physical interpretation will be explained when we discuss cosmology in more detail, its extremely small value in nature implies that the approximation $\Lambda \approx 0$ is very good at any length scale below Megaparsecs. On such scales, strong gravitational fields appear near localized sources, the field becoming very weak far away from them. Therefore, it is a good approximation for most non-cosmological purposes to model spacetime as asymptotically flat, solving Einstein's equations with $\Lambda = 0$ and assuming asymptotic conditions at 'infinity' approaching Minkowski spacetime. The weak-field approximation is actually good in most places, either because the gravitational fields are weak everywhere, or because one measures them far away from the possibly strong sources. In this chapter we consider the weak-field approximation to Einstein's equations in asymptotically flat spacetime. This will allow us to set the language used in discussing actual tests of small relativistic effects, the physics of gravitational radiation and some more formal topics like the definition of global notions of energy.

6.1 Weak Gravitational Fields

We begin our analysis of weak gravitational physics with the linearized approximation of Einstein's equations with $\Lambda = 0$, valid for the physical situation of weak gravitational fields in a Minkowski background or more generally for weak gravitational fields in any spacetime whose radius of curvature is much larger than the length scales of interest. In the course of this analysis, we will make contact with the Newtonian theory, define global notions of mass and angular momentum in asymptotically flat spaces, and obtain an elementary discussion of gravitational radiation.

6.1.1 Linearized GR

The basic situation under consideration is that of metrics of the form

$$g_{\alpha\beta} = \eta_{\alpha\beta} + h_{\alpha\beta} , \tag{6.1}$$

with $h_{\alpha\beta}$ 'small'. The appropriate notion of 'smallness' must be qualified because of the basic ambiguity by reparametrizations. Even flat Minkowski space can show a metric tensor that has 'large' components, simply because it was written in a frame with large accelerations. Hence,

we shall split the metric as above, as a Minkowskian part plus a perturbation, adopting a *preferred* Minkowskian frame in which the flat part of the metric is the standard Minkowski one η_{ab} . After this preferred frame x^a has been chosen, only Lorentz transformations on the x^a are allowed, leaving the background metric η_{ab} invariant. The action of a general infinitesimal reparametrization on this preferred Minkowskian frame, $x^a \rightarrow x^a - \varepsilon^a(x)$, induces a change of the metric *functions*

$$g_{ab}(x) \rightarrow g_{ab}(x) + \partial_a \varepsilon_b(x) + \partial_b \varepsilon_a(x) + O(\varepsilon^2) , \quad (6.2)$$

which can be reinterpreted as a redefinition of h_{ab} , rather than a reparametrization of the Minkowskian coordinates x^a . Therefore, the linearized gravitational field tensor h_{ab} inherits a *gauge redundancy*

$$h_{ab} \rightarrow h_{ab} + \partial_a \varepsilon_b + \partial_b \varepsilon_a \quad (6.3)$$

as the linearized approximation of the general coordinate invariance of the underlying theory.

We can now linearize Einstein's equations, starting from

$$\Gamma_{ab}^c = \Gamma_{ab}^{c(1)} + O(h^2) = \frac{1}{2} \eta^{cd} (\partial_a h_{bd} + \partial_b h_{ad} - \partial_d h_{ab}) + O(h^2) . \quad (6.4)$$

In the following, it will be convenient to raise and lower Lorentz indices with the Lorentz metric η_{ab} . The Ricci tensor linearizes as

$$R_{ab} = R_{ab}^{(1)} + O(h^2) = \partial_c \Gamma_{ab}^{c(1)} - \partial_a \Gamma_{bc}^{c(1)} + O(h^2) , \quad (6.5)$$

and the Einstein tensor is given by

$$G_{ab} = G_{ab}^{(1)} + O(h^2) = \frac{1}{2} \left(-\partial^2 \gamma_{ab} + \partial_a \partial^c \gamma_{cb} + \partial_b \partial^c \gamma_{ca} - \eta_{ab} \partial^c \partial^d \gamma_{cd} \right) + O(h^2) \quad (6.6)$$

in terms of the related field

$$\gamma_{ab} \equiv h_{ab} - \frac{1}{2} \eta_{ab} h , \quad h \equiv \eta^{ab} h_{ab} . \quad (6.7)$$

So the Einstein equation in the linearized approximation reads

$$G_{ab}^{(1)} = 8\pi G T_{ab} . \quad (6.8)$$

It is easy to check that this is none other than the Fierz–Pauli field equation for a massless symmetric tensor, coupled minimally to the energy-momentum tensor of matter. So, we recover the first step in the long-road program towards GR, if we did not fancy taking the shortcut through the Equivalence Principle.

The linearized Einstein tensor simplifies considerably by choosing an analog of the Lorenz gauge in electrodynamics,

$$\partial^a \gamma_{ab} = 0 , \quad (6.9)$$

in which the linearized Einstein's equations boil down to

$$-\partial^2 \gamma_{ab} = 16\pi G T_{ab} , \quad (6.10)$$

taking the form of a standard wave equation, with an equally standard solution in terms of retarded potentials

$$\gamma_{ab}(x)_{\text{retarded}} = 4G \int \frac{T_{ab}(t - |\vec{x} - \vec{y}|, \vec{y})}{|\vec{x} - \vec{y}|} d^3 y \quad (6.11)$$

for fields that vanish at infinity. Fields that do not vanish at infinity can be incorporated by adding solutions of the homogeneous vacuum equation, $\partial^2 \gamma_{ab} = 0$, i.e. gravitational waves propagating in *vacuo* at the speed of light.

6.1.2 Newtonian and Post-Newtonian

Assuming that the energy-momentum of the sources is time-independent and non-relativistic, i.e. velocities of massive objects are small: $|T^{00}| \gg |T^{0i}| \gg |T^{ij}|$, we approximate the solution of (6.11) by $\gamma_{ij} \approx 0 \approx \gamma_{0i}$, and the dominant component is

$$\gamma_{00}(\vec{x}) \approx 4G \int d^3y \frac{T_{00}(\vec{y})}{|\vec{y} - \vec{x}|}. \quad (6.12)$$

Since $T_{00} \approx \rho_m$ for non-relativistic sources, the matter *mass* density, we have a direct relation to the Newtonian potential ϕ_N ,

$$\gamma_{00} \approx -4\phi_N. \quad (6.13)$$

Using now $\gamma = -h = -\gamma_{00}$ and $h_{ab} = \gamma_{ab} - \frac{1}{2}\eta_{ab}\gamma$ we obtain for the original metric perturbation

$$h_{00} \approx \frac{1}{2}\gamma_{00}, \quad h_{0i} \approx 0, \quad h_{ij} \approx \delta_{ij}h_{00},$$

which gives back the known asymptotic form of the metric perturbation,

$$h_{00} \approx -2\phi_N = \frac{2GM}{|\vec{x}|}, \quad M \equiv \int d^3x \rho_m(\vec{x}),$$

previously derived from the non-relativistic limit of the geodesic equation. Hence, we have found the following asymptotic form of the metric as sourced by a non-relativistic system of mass M :

$$ds^2 = -\left(1 - \frac{2GM}{|\vec{x}|} + \dots\right) dt^2 + \left(1 + \frac{2GM}{|\vec{x}|} + \dots\right) d\vec{x}^2, \quad (6.14)$$

valid for $|\vec{x}| \gg 2GM$.

An important observation regarding the asymptotic form (6.14) is the possibility of reading off the mass of the field source by inspection of the long-distance tails of the deviation of the metric from the Minkowski form. This is the GR analog of Gauss' theorem, and it will be a recurring theme in what follows.

The metric (6.14) is a more accurate one than the strict Newtonian form (3.79), including the effect of purely spatial curvature in addition to the temporal curvature. This is enough to get the leading effects of light bending right. Still, to get the correct perihelion precession for Mercury one needs to go to order $(GM/r)^2$ in g_{00} . This was done by Einstein in November 1915 a few weeks before Schwarzschild discovered his complete nonlinear metric.¹ To understand why the leading post-Newtonian effects regarding the orbit of Mercury require going one step further than the leading linearized approximation (6.14), consider a general metric *ansatz* of the form

$$ds^2 = -(1 + 2\phi + 2\beta\phi^2)dt^2 + (1 - 2\gamma\phi + \zeta\phi^2)d\vec{x}^2, \quad (6.15)$$

with β, γ, ζ some constants. This captures the Newtonian approximation when $\phi = -GM/|\vec{x}|$ represents the Newtonian potential and parametrizes the leading corrections up to, and including, order $(GM/|\vec{x}|)^2$. We can now consider a test particle freely falling in this metric, and

¹The g_{00} component in the ordinary Schwarzschild metric does not contain $(GM/r)^2$ terms, but notice that (6.14) is written in *isotropic* coordinates because of the presence of the $d\vec{x}^2$ term with the definition of radius as $|\vec{x}|$. So we should rather compare (6.14) with the large distance expansion of (5.80) which yields a correction $-2G^2M^2/|\vec{x}|^2$ to g_{00} .

expand the Lagrangian out to next-to-leading order in the non-relativistic approximation. Since ϕ is the Newtonian potential we have $v^2 \sim \phi$ in typical bound orbital motions. Therefore, keeping terms to $O(v^4)$ requires that we also keep the $O(\phi^2)$ and $O(v^2\phi)$ terms, leading to

$$L = -m\sqrt{-g_{ab}\frac{dx^a}{dt}\frac{dx^b}{dt}} \approx -m + \frac{1}{2}m\vec{v}^2 + \frac{1}{8}m\vec{v}^4 - m\phi + \frac{1}{2}m(1-2\beta)\phi^2 - (\frac{1}{2} + \gamma)m\phi\vec{v}^2 + \dots, \quad (6.16)$$

so that the $O(\phi^2)$ term in the time-time component g_{00} of the metric (6.15) does contribute to the relativistic corrections. Notice, however, that the term $\zeta\phi^2$ in the spatial components of the metric only contributes to order $\phi^2\vec{v}^2$ and therefore can be dropped at the level of accuracy respected in (6.16). The $O(\vec{v}^4)$ and $O(\phi^2)$ terms are respectively the leading relativistic corrections to the kinetic and potential energy of the particle. On the other hand, the $O(\phi\vec{v}^2)$ term represents qualitatively new forces that depend on velocities.

In the analysis of precision experimental tests, one usually parametrizes the deviation from Einstein's theory by the so-called post-Newtonian parameters which represent the asymptotic metric *ansatz*

$$ds^2 \rightarrow -\left(1 - \frac{2GM}{|\vec{x}|} + \beta \frac{2G^2M^2}{|\vec{x}|^2} + \dots\right) dt^2 + \left(1 + \gamma \frac{2GM}{|\vec{x}|} + \dots\right) d\vec{x}^2 + \dots \quad (6.17)$$

We can quote the equivalent presentation of this metric in the 'standard' or Schwarzschild frame, rather than the 'isotropic' frame. The standard frame is characterized by defining the radial coordinate r so that the area of spatial spheres at radius r is $4\pi r^2$. So, the standard presentation of the post-Newtonian spacetime (6.17) is

$$ds^2 \rightarrow -\left(1 - \frac{2GM}{r} + 2(\beta - \gamma) \frac{G^2M^2}{r^2} + \dots\right) dt^2 + \left(1 + \gamma \frac{2GM}{r} + \dots\right) dr^2 + r^2 d\Omega^2 + \dots \quad (6.18)$$

The prediction of GR is $\beta = \gamma = 1$. Recalculating classic tests for the metric (6.18) one finds a factor of $(1 + \gamma)/2$ with respect to the GR prediction, for the bending of light and the echo delay. The precession of perihelia is corrected by a factor of $(2 + 2\gamma - \beta)/3$. On the other hand, the gravitational redshift remains the same as in GR to this order in the gravitational potential. Together, these solar-system precision tests bound $|1 - \gamma|$ and $|1 - \beta|$ to be at most of order 10^{-5} .

Parametrizing Extensions of GR

The reader may be surprised by our insistence on checking the deviations from GR at the 10^{-5} level through the post-Newtonian parameters γ and β , given that the Principle of Equivalence is tested to the 10^{-15} level. The reason is that the minimal Einstein theory is not the only one that is compatible with the Principle of Equivalence. What this principle tests is the universality of free fall, that is to say, the universality of the coupling of matter to gravitation. In Einstein's language, this means that the effect of gravity on matter is equivalent to postulating the existence of a metric that couples minimally to matter at the Lagrangian level, at least to leading order in the long-distance expansion, but this universal metric may not be determined by the standard Einstein's equations of minimal GR. An example that illustrates the point is the addition of a small correction to the gravitational dynamics by postulating a scalar field φ

with gravitational-strength couplings to matter, such as the theory (3.1) with action

$$S = \frac{1}{16\pi\tilde{G}} \int \sqrt{-\tilde{g}} \left(\tilde{R} - 2\tilde{g}^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi \right) + \int \sqrt{-\tilde{g}} \left(\mathcal{L}_m(\tilde{g}_{\mu\nu}, \Psi) + \alpha \varphi \tilde{g}_{\alpha\beta} T^{\alpha\beta} \right), \quad (6.19)$$

where $\tilde{R}$ is the Ricci scalar for the metric $\tilde{g}_{\mu\nu}$ that has a standard Hilbert action. The coupling to matter of the new ‘gravitational’ degree of freedom φ is modulated by a small dimensionless parameter α and we have allowed for a parametric difference between the Newton’s constant appearing in the Lagrangian, $\tilde{G}$, and the one measured in, say, a Cavendish experiment, G . This scalar coupling redefines the free-fall metric by a conformal rescaling $\tilde{g}_{\mu\nu} \rightarrow g_{\mu\nu} = (1 + \alpha\varphi)^2 \tilde{g}_{\mu\nu}$. For point particles this is exact, since $\alpha \int \sqrt{-\tilde{g}} \varphi \tilde{g}_{\alpha\beta} T^{\alpha\beta} = -\alpha \sum_p m_p \int d\tilde{\tau}_p \varphi$ turns the particle action into $-\sum_p m_p \int d\tilde{\tau}_p (1 + \alpha\varphi) = -\sum_p m_p \int d\tau_p$, while for general matter it holds to leading order in α . So, the Principle of Equivalence holds with respect to the rescaled metric $g_{\mu\nu}$, but Einstein’s equations are now extended by the presence of φ .² Examining a vacuum solution for the coupled system of $\tilde{g}_{\mu\nu}$ and φ , with asymptotic behavior $g_{00} \sim -1 + \frac{2GM}{r}$ at large r , one finds an effective metric where matter ‘falls’, $g_{\mu\nu}$, of the form (6.18) with post-Newtonian parameters

$$\gamma = 1 - 2\alpha^2 + O(\alpha^4), \quad \beta = 1 - \frac{1}{2}\alpha^4 + O(\alpha^6), \quad (6.20)$$

and a phenomenological Newton’s constant given by $G = \tilde{G}(1 + \alpha^2)$. The value of β is sensitive to the form of the coupling at second order in φ : for instance, the famous Jordan–Brans–Dicke theory has an exponential coupling $g_{\mu\nu} = e^{2\alpha\varphi} \tilde{g}_{\mu\nu}$, giving the same γ , but $\beta = 1$ exactly.

So we see how this theory can satisfy Einstein’s Principle of Equivalence to the 10^{-15} level and yet have post-Newtonian corrections of order 10^{-5} . In view of (6.20), those corrections are mostly felt in the γ parameter, governing light bending and echo delay, which are the effects for which the scalar does not contribute at all, and bound the dimensionless strength parameter α to be at most of order 10^{-3} . Various non-linear generalizations of such scalar-metric theories exist, among which the Jordan–Brans–Dicke theory played an important historical role. Another important lesson is that (6.19) gives a simple example of a theory that violates the so-called ‘Strong Principle of Equivalence’, by which we mean that universality of free fall includes extended systems with significant gravitational self-energy. Near a heavy body, the self-energy has contributions from the gradients of the new φ field, and not just from the $\tilde{g}_{\mu\nu}$ metric. A measure of the violation of the Strong Principle of Equivalence is given to post-Newtonian level by the so-called Nordtvedt parameter³

$$\eta_N = 4\beta - \gamma - 3, \quad (6.21)$$

which vanishes for GR and is of order $2\alpha^2$ for the scalar-metric theory discussed here.

²The action (6.19) in terms of $\tilde{g}_{\mu\nu}$ is known as the ‘Einstein-frame action’. Equivalently, we can rewrite the whole action in terms of the metric $g_{\mu\nu}$ felt by matter. The Hilbert term then becomes $\sqrt{-g} R / 16\pi\tilde{G}(1 + \alpha\varphi)^2$, plus a modified kinetic term for φ , so that we would be working with a φ -dependent Newton constant, a form of the action known as the ‘Jordan frame’.

³This formula can be derived within an extension of the effective Lagrangian (6.16) to a system of many gravitating bodies. In GR, this many-body Lagrangian was first obtained by Einstein, Infeld and Hoffmann in 1938. Its post-Newtonian generalization, and the formula for η_N , are due to Nordtvedt (1968).

Problem: Scalar-Tensor Theory

Consider the scalar-tensor theory (6.19) and derive the post-Newtonian parameters (6.20) of the Jordan-frame metric. Embed this model into the Jordan–Brans–Dicke family of theories (more precisely, into its generalization with a field-dependent Brans–Dicke parameter) and present the resulting Lagrangians in both Einstein and Jordan frame.

6.1.3 Dipole Tails from a Little Rotation

Coming back to the safe realm of GR, we have seen that the leading Newtonian approximation to (6.11) yields a long-distance metric field $h_{00} \sim 2GM/|\vec{x}|$. This kind of approximation can be systematized by performing a *multipole expansion* of the integral, i.e. we consider sources with effectively ‘compact’ support over a region of average size $\ell \ll |\vec{x}|$ and expand

$$\frac{1}{|\vec{x} - \vec{y}|} = \frac{1}{|\vec{x}|} + \frac{\vec{x} \cdot \vec{y}}{|\vec{x}|^3} + \frac{3(\vec{x} \cdot \vec{y})^2 - |\vec{x}|^2 |\vec{y}|^2}{2|\vec{x}|^5} + \dots$$

Monopole terms are of $O(1/|\vec{x}|)$, dipole terms of $O(1/|\vec{x}|^2)$, quadrupole terms of $O(1/|\vec{x}|^3)$, and so on. This expansion can be applied to all components of the metric field γ_{ab} . For simplicity, we shall continue with our assumption that T_{ab} is time-independent, although this hypothesis can be lifted by a more detailed treatment of the multipole expansion. At any rate, any source can be regarded as stationary in time if the time resolution of the measurements average out fast motions, so we can regard a source stationary if the time-dependence is lost to us ‘for all practical purposes’.

For stationary T_{ab} , the conservation equation $\partial_a T^{ab} = 0$ implies $\sum_k \partial_k T_{ak} = 0$. This leads to some useful identities for the spatial integrals of the energy-momentum tensor. Denoting $\int d^3y \equiv \int_{\mathbb{R}^3}$, we have

$$\int_{\mathbb{R}^3} T_{aj} = \int_{\mathbb{R}^3} \sum_k \partial_k (T_{ka} x_j) = 0, \quad (6.22)$$

where we have neglected the boundary terms in the last step, under the assumption that T_{ab} has compact support. Setting $a = 0$ in this equation we obtain that the total momentum vanishes $P_i = \int_{\mathbb{R}^3} T_{0i} = 0$. With $a = i$, we see that the volume average of the stresses T_{ij} also vanishes. This implies that the monopole approximations to γ_{0i} and γ_{ij} both vanish, and these terms are dominated by the dipole contribution. Since $h_{0i} = \gamma_{0i}$ and $h_{ij} = \gamma_{ij} - \frac{1}{2} \delta_{ij} \gamma$, with $\gamma = -\gamma_{00} + \text{dipole terms}$, we see that h_{ij} is dominated by the monopole contribution to γ_{00} .

Hence, in order to capture the leading behavior of the metric perturbation, it suffices to consider the monopole approximation to γ_{00} and the dipole approximation to γ_{0i} . Under these conditions, the monopole and dipole approximations to γ_{0i} are

$$h_{0i}(\vec{x}) = \gamma_{0i}(\vec{x}) = \frac{4G}{|\vec{x}|} \int_{\mathbb{R}^3} T_{0i} + \frac{4G}{|\vec{x}|^3} \sum_j x_j \int_{\mathbb{R}^3} y_j T_{0i} + \dots$$

The first monopole term, proportional to the total momentum of the system, vanishes for time-independent sources. More generally, it can be eliminated by adjusting the frame to be at rest with respect to the sources. Then, as already stated above, the leading contribution to h_{0i} is the dipole term. We can simplify it by using again the time-independence condition to write

$$0 = \int_{\mathbb{R}^3} y_i y_j \sum_k \partial_k T_{0k} = - \int_{\mathbb{R}^3} (y_j T_{0i} + y_i T_{0j}),$$

so that the dipole term is purely antisymmetric in the indices i, j . Finally, using that the angular momentum is defined as

$$J_k \equiv \frac{1}{2} \sum_{ij} \epsilon_{ijk} J_{ij} \equiv -\frac{1}{2} \sum_{ij} \epsilon_{ijk} \int_{\mathbb{R}^3} (y_i T_{0j} - y_j T_{0i}) , \quad (6.23)$$

(the minus sign is there because we have lowered the time component of the energy-momentum tensor) we see that we can read off the angular momentum of the source system from the $O(1/|\vec{x}|^2)$ tail of the off-diagonal metric components,

$$h_{0i}(\vec{x}) = \frac{2G}{|\vec{x}|^3} \sum_{j,k} \epsilon_{ijk} x_j J_k . \quad (6.24)$$

Hence, in *stationary situations*, the long-distance behavior of the asymptotically flat spacetime is controlled by the total mass and angular momentum of the sources ‘inside’ by the following $|\vec{x}| \rightarrow \infty$ asymptotics:

$$ds^2 \rightarrow - \left(1 - \frac{2GM}{|\vec{x}|} + \dots \right) dt^2 + \left(\frac{4G}{|\vec{x}|^3} (\vec{x} \times \vec{J}) + \dots \right) \cdot d\vec{x} dt + \left(1 + \frac{2GM}{|\vec{x}|} + \dots \right) d\vec{x}^2 . \quad (6.25)$$

6.1.4 Gravitomagnetism

The off-diagonal terms in (6.25) have a characteristic effect on test particles. As we have seen in (3.78), the effect of a term $2\vec{h} d\vec{x} dt$ in the metric can always be reinterpreted as inducing a ‘Coriolis-like’ force with effective angular velocity $\frac{1}{2} \vec{\partial} \times \vec{h}$. More precisely, we have a ‘Coriolis field’, since the effective angular velocity has a non-trivial space dependence. Indeed, if we rerun the analysis of the particle action in (6.16) by including such an off-diagonal term in the metric we find a correction to the effective particle Lagrangian of the form $\delta L = m \vec{h} \cdot \vec{v}$, to leading order in $\vec{h}$. This has the same form as a coupling of a charged particle to an electromagnetic vector potential, so that we may regard such coupling as ‘gravitomagnetic’. Using the explicit form of $\vec{h} = 2G(\vec{x} \times \vec{J})/|\vec{x}|^3$, we can write the effective interaction in the form of a ‘magnetic moment’ correction:

$$\delta L = m \vec{h} \cdot \vec{v} = -\vec{\mu}_J \cdot \vec{L} , \quad \vec{\mu}_J \equiv \frac{2G}{|\vec{x}|^3} \vec{J} . \quad (6.26)$$

As a result, we find the so-called Lense–Thirring precession effect on the orbital momentum, as a direct analog of the Larmor precession in a magnetic field. To derive it, notice that a correction $\delta L = m \vec{h} \cdot \vec{v}$ to the Lagrangian produces, to leading order in $\vec{h}$, a correction $\delta H = -\vec{p} \cdot \vec{h}$ to the Hamiltonian. Using then the Hamilton equations one obtains

$$\frac{d\vec{L}}{dt} = \frac{\partial H}{\partial \vec{p}} \times \vec{p} - \vec{x} \times \frac{\partial H}{\partial \vec{x}} = \frac{2G}{|\vec{x}|^3} \left(-(\vec{x} \times \vec{J}) \times \vec{p} + \vec{x} \times (\vec{J} \times \vec{p}) \right) ,$$

which further reduces to

$$\frac{d\vec{L}}{dt} = \vec{\mu}_J \times \vec{L} . \quad (6.27)$$

We find that orbital planes undergo a precession induced by a spin-orbit coupling which is entirely analogous to the one operating between the nuclear spin and the electron’s orbital motion in atoms. This effect is also responsible for a small precession of perihelia of the same

order of magnitude. When applied to the motion of gyroscopes, we obtain the same physics described by the spin free-fall equation (3.87).

The effective Coriolis field induced by a rotating body produces a ‘frame dragging’ effect, namely inertial bodies tend to co-rotate with a rotating heavy object, as if the heavy object was ‘stirring along’ the fabric of space, thus dragging along inertial motions of test particles. This is reminiscent of the old critique of Newton’s rotating bucket by Ernst Mach and it is usually quoted as evidence that GR incorporates at least some aspects of Mach’s views about the foundations of mechanics.

Problem: Newton’s Bucket

Consider a rotating hollow sphere of mass M and radius R . Compute the leading order h_{0i} components of the metric inside the sphere and interpret them in terms of a frame dragging effect (this was done by Thirring in 1918). Discuss the lessons to be learned in the context of Mach’s critique of Newton’s bucket example.

Problem: Spin-Induced Orbital Precession

Consider the secular effect of the sun’s spin on the orbit of a planet. In addition to the Larmor-like precession of the orbital plane obtained in (6.27), one can derive a precession of perihelia within the plane of the orbit, by studying the evolution of the Runge–Lenz vector

$$\vec{\mathcal{A}} = \frac{1}{m} \vec{p} \times \vec{L} - \frac{GMm}{|\vec{x}|} \vec{x} ,$$

which is accidentally conserved under the purely Newtonian Kepler motion, staying constant in the orbital plane and directed towards the perihelion. Therefore, the two vectors $\vec{L}$ and $\vec{\mathcal{A}}$ together determine the overall orientation of the orbit.

Compute the time derivative of the Runge–Lenz vector to obtain

$$\frac{d\vec{\mathcal{A}}}{dt} = \frac{2G}{|\vec{x}|^3} \vec{J} \times \vec{\mathcal{A}} + \frac{6G}{m|\vec{x}|^5} (\vec{J} \cdot \vec{L}) (\vec{x} \times \vec{L}) .$$

The first term gives the Larmor-like precession. Upon further averaging over a full revolution, show that

$$\left\langle \frac{d\vec{V}}{dt} \right\rangle_{\text{secular}} = \vec{\Omega}_{\text{LT}} \times \vec{V} ,$$

where $\vec{V}$ is either $\vec{L}$ or $\vec{\mathcal{A}}$, and the Lense–Thirring angular velocity is given by

$$\vec{\Omega}_{\text{LT}} = \frac{2G|\vec{J}|}{a^3(1-e^2)^{3/2}} (\vec{n}_J - 3\vec{n}_L(\vec{n}_J \cdot \vec{n}_L)) ,$$

where a is the semimajor axis of the ellipse and e its eccentricity, while $\vec{n}_J$ and $\vec{n}_L$ are unit vectors in the $\vec{J}$ and $\vec{L}$ directions, respectively.

We find that the full orbit as a whole precesses around the Lense–Thirring vector $\vec{\Omega}_{\text{LT}}$.

Problem: Precession of Orbiting Gyroscopes

Consider a spherical gyroscope in orbit. Starting from the term $-\frac{3}{2}m\phi\vec{v}^2$ in (6.16), integrate it over the particles forming the gyroscope to obtain an effective Lagrangian

$$\delta L_{\text{eff}} \approx \frac{3GM}{2} \int d^3z \rho(\vec{z}) \frac{(\vec{v}_0 + \vec{\omega} \times \vec{z})^2}{|\vec{x}_0 + \vec{z}|}.$$

where $\vec{v}_0$ and $\vec{x}_0$ are the velocity and position of the gyro's center of mass, with mass density $\rho(\vec{z})$, and $\vec{\omega}$ is its angular velocity. We further consider the cross-term proportional to $\vec{v}_0 \cdot (\vec{\omega} \times \vec{z})$ and use the dipole approximation

$$\frac{1}{|\vec{x}_0 + \vec{z}|} \approx \frac{1}{|\vec{x}_0|} - \frac{\vec{z} \cdot \vec{x}_0}{|\vec{x}_0|^3} + \dots$$

Calculate the integral using

$$\int d^3z \rho(\vec{z}) z_i z_j = \frac{1}{2} I \delta_{ij},$$

with I the gyro's moment of inertia, so that its spin is $\vec{S} = I\vec{\omega}$. Then, prove that the effective Lagrangian reduces to

$$\delta L_{\text{geo}} \approx -\vec{\Omega}_{\text{geo}} \cdot \vec{S},$$

where the *geodetic* frequency is given by

$$\vec{\Omega}_{\text{geo}} \approx \frac{3GM}{2|\vec{x}_0|^3} (\vec{x}_0 \times \vec{v}_0),$$

and induces the so-called *geodetic* precession of the top's spin

$$\frac{d\vec{S}}{dt} = \vec{\Omega}_{\text{geo}} \times \vec{S}.$$

There is also a *hyperfine* effect (also referred to as Lense–Thirring effect), coupling directly the gyro's spin $\vec{S}$ to the spin of the gravitational field, $\vec{J}$, induced by the “magnetic” term $2G\vec{v} \cdot (\vec{x} \times \vec{J})/|\vec{x}|^3$, i.e.

$$\delta L = 2G \int d^3z \rho(\vec{z}) \frac{\vec{J} \cdot ((\vec{v}_0 + \vec{\omega} \times \vec{z}) \times \vec{x})}{|\vec{x}|^3},$$

with $\vec{x} = \vec{x}_0 + \vec{z}$. Perform this integral in the dipole approximation to obtain

$$\delta L_{\text{hyper}} \approx -\vec{\Omega}_{\text{hyper}} \cdot \vec{S},$$

where

$$\vec{\Omega}_{\text{hyper}} = \frac{3G\vec{x}_0(\vec{x}_0 \cdot \vec{J})}{|\vec{x}_0|^5} - \frac{G\vec{J}}{|\vec{x}_0|^3}.$$

The geodetic precession for the case of satellites around Earth is in the ball park of 10 seconds of arc per year. The hyperfine precession is smaller by a factor

$$\left| \frac{\vec{\Omega}_{\text{hyper}}}{\vec{\Omega}_{\text{geo}}} \right|_{\oplus} \sim 6.5 \times 10^{-3}.$$

6.1.5 Gravitational Energy

In a strict sense, the basic equation of linearized GR (6.8) is inconsistent with any non-trivial self-gravity effects. The linearized Einstein tensor happens to be transverse:

$$\partial^a G_{ab}^{(1)} = 0 , \quad (6.28)$$

which implies conservation of the matter energy momentum alone: $\partial^a T_{ab} = 0$. But this implies that particles contributing to T_{ab} necessarily follow geodesics of the zeroth-order metric η_{ab} , instead of the perturbed one $\eta_{ab} + h_{ab}$. This is a manifestation of the old inconsistency of the Fierz–Pauli theory that we described already. The solution involved correcting the Fierz–Pauli dynamics adding non-linearities in the h_{ab} field. Now, with the benefit of hindsight, we can perform this task in one step, by a suitable generalization of (6.11) to an *exact* equation equivalent to Einstein’s equations.

Under the basic decomposition $g = \eta + h$, we write the full Einstein tensor as

$$G_{ab}(\eta + h) = G_{ab}^{(1)}(h) + G'_{ab}(\eta + h) , \quad (6.29)$$

where G'_{ab} is simply the deviation of the exact Einstein tensor from the linear approximation. By definition, the small field expansion of G'_{ab} starts at *quadratic* order in h_{ab} . Defining now

$$t_{ab}(h) \equiv -\frac{1}{8\pi G} G'_{ab}(\eta + h) \quad (6.30)$$

we see that Einstein’s equations can be exactly rewritten as

$$G_{ab}^{(1)} = 8\pi G \mathcal{T}_{ab} , \quad (6.31)$$

where

$$\mathcal{T}_{ab} \equiv T_{ab} + t_{ab} \quad (6.32)$$

is a sort of *total* energy momentum tensor of matter and gravitation. This definition has t_{ab} obviously playing the role of the energy-momentum ‘tensor’ of the gravitational field, and is completely tied to the particular background flat spacetime and the family of coordinate systems in which its metric takes the Minkowskian simple form. The quotations here refer to the fact that t_{ab} is not a true tensor with respect to general diffeomorphisms of the full spacetime. It is only a Lorentz tensor with respect to the flat background spacetime. One should not become mystified about this fact, since the whole perturbative construction singles out the Minkowski metric η_{ab} as special, and the equations are written in manifestly non-covariant form.

Now, in the situation in which the complete solution for γ_{ab} vanishes at infinity, the equation (6.11) still holds exactly with the replacement of T_{ab} everywhere by $\mathcal{T}_{ab}$

$$\gamma_{ab}(x)_{\text{retarded}} = 4G \int d^3y \frac{\mathcal{T}_{ab}(t - |\vec{y} - \vec{x}|, \vec{y})}{|\vec{x} - \vec{y}|} . \quad (6.33)$$

Notice that now $\mathcal{T}_{ab}$ depends on γ_{ab} , so that this is an integral equation for γ_{ab} , containing exactly the same information as Einstein’s equations (plus the condition that γ_{ab} vanishes at infinity, preserving the asymptotic flatness). The systematics of the weak-field expansion can be obtained from the iterative solution of this integral equation. Namely, the n -th order term in the evaluation of h_{ab} via eq. (6.33) uses the $(n - 1)$ -th order evaluation of the gravitational

stress-energy tensor t_{ab} , plus the matter energy-momentum tensor calculated in the $(n - 1)$ -th order background metric, after solving the matter equations. In particular, this construction only requires that γ_{ab} is a weak field at large distances, since (6.33) is valid for arbitrarily strong fields which still preserve the condition of asymptotic flatness. Hence, the iterative weak-field solution of (6.33) can be understood as a gradual reconstruction of the field ‘from outside in’.

The standard local conservation law in Minkowski space also holds *exactly* in terms of the complete tensor:

$$\partial_a \mathcal{T}^{ab} = 0 , \quad (6.34)$$

as a result of the exact transversality of the linearized Einstein tensor, $\partial^a G_{ab}^{(1)} = 0$.

This iterative scheme shows precisely how the Lorentz-invariant theory of a symmetric tensor field (the Fierz–Pauli scheme) must be corrected by nonlinear terms in order to achieve consistency with energy-momentum conservation (6.34). Hence, in hindsight we have found that Einstein’s theory is the minimal consistent theory of a symmetric tensor field.

For sufficiently localized systems, with T_{ab} of compact support, and in the absence of radiation, one finds that $\gamma_{ab} = O(1/|\vec{x}|)$ at spatial infinity. Then $\mathcal{T}_{ab} \sim \partial\gamma\partial\gamma = O(1/|\vec{x}|^4)$ and its integral over a spatial section converges. In this situation, we can derive a useful formula (due to Arnowitt, Deser and Misner) for the total energy in asymptotically flat spacetimes. In the transverse gauge $\partial^a \gamma_{ab} = 0$, the exact Einstein equations (6.31) can be written as $-\partial^2 \gamma_{ab} = 16\pi G \mathcal{T}_{ab}$, so that

$$E_{\text{ADM}} = \int_{\mathbb{R}^3} \mathcal{T}_{00} = -\frac{1}{16\pi G} \int_{\mathbb{R}^3} \partial^2 \gamma_{00} . \quad (6.35)$$

By repeated use of the transversality condition, $\partial^a \gamma_{ab} = 0$, we can eliminate the time derivatives to obtain

$$E_{\text{ADM}} = \frac{1}{16\pi G} \sum_{i,j} \oint_{S_\infty^2} dS^i (\partial_j \gamma_{ij} - \partial_i \gamma_{00}) = \frac{1}{16\pi G} \sum_{i,j} \oint_{S_\infty^2} dS^i (\partial_j h_{ij} - \partial_i h_{jj}) . \quad (6.36)$$

Once we have obtained this formula in the transverse gauge, one can directly see that it is gauge invariant under those transformations of the form (6.3) that preserve the condition of asymptotic flatness, namely those such that $\partial_i \varepsilon_j + \partial_j \varepsilon_i = O(1/|\vec{x}|)$.

This formula shows that the total energy in stationary, asymptotically flat spacetimes only depends on the asymptotic properties of the metric, and can be obtained as an integral over the *celestial sphere* at infinity S_∞^2 , a result known as the Arnowitt–Deser–Misner (ADM) expression.⁴ Notice that the ADM energy does contain the gravitational self-energy, since it is the integral of $T_{00} + t_{00}$, including the energy stored in the gravitational field through t_{ab} . If we evaluate (6.36) for the asymptotic metric (6.25) we learn that $E_{\text{ADM}} = M$ in this case, which coincides with the effective mass measured by a probe Newtonian particle, from the asymptotic form of g_{00} , so we may as well read the ADM mass from the leading $2GM/r$ correction to the temporal component of the metric. This result can be generalized to obtain similar expressions for the total momentum and angular momentum of localized self-gravitating systems. The answer is also equivalent to reading off the total angular momentum contained in $\mathbb{R}^3$ from (6.25).

⁴An important theorem by Schoen and Yau establishes that $E_{\text{ADM}} \geq 0$ for any smooth, asymptotically flat, solution of Einstein’s equations, provided $T_{\mu\nu}$ satisfies appropriate positivity conditions.

Komar Energy

We have seen how the ADM mass, which includes contributions from self-gravity effects, can be read off from the asymptotic tail of the metric in an asymptotically flat solution. There is an alternative version of this statement that does not make use of the auxiliary flat background spacetime that we introduced via the split $g_{ab} = \eta_{ab} + h_{ab}$, or the corresponding pseudo-tensor $\mathcal{T}_{ab}$, and only employs the complete metric $g_{\alpha\beta}$ that solves the full non-linear Einstein's equations. The necessary assumption on the exact spacetime is that it has a timelike isometry, i.e. a Killing vector field ξ_t^μ which asymptotically looks like the standard Minkowski vector field ∂_t . So the price we pay for disposing of the auxiliary scaffolding is that we have to restrict to exactly isometric spacetimes.

Given a Killing vector ξ^μ let us define the antisymmetric tensor

$$K_{\mu\nu} = \nabla_\mu \xi_\nu - \nabla_\nu \xi_\mu . \quad (6.37)$$

Just as the electromagnetic tensor, it is constructed as an exact two-form, with vanishing exterior derivative $\epsilon^{\mu\nu\rho\sigma} \nabla_\nu K_{\rho\sigma} = 0$, the analog of the homogeneous Maxwell equations. As in electrodynamics, this identity only implies that the ‘magnetic’ flux of $K_{\mu\nu}$ through any closed two-surface vanishes. The interesting quantity is the ‘electric’ flux, whose behavior is controlled by the analog of Gauss’ law, namely the divergence of the two-form:

$$\nabla^\mu K_{\mu\nu} = -2R_{\nu\alpha} \xi^\alpha , \quad (6.38)$$

by simply applying the Ricci identity (4.11) and the Killing equation $\nabla_{(\alpha} \xi_{\beta)} = 0$. In particular, the two-form is divergence-free in any vacuum region, $R_{\mu\nu} = 0$, so that its electric flux through a two-sphere surrounding all the sources does not depend on the size or shape of the two-sphere, and equals its value on the celestial sphere S_∞^2 . Using Einstein's equations we can further write

$$\nabla^\mu K_{\mu\nu} = -16\pi G \left(T_{\nu\alpha} - \frac{1}{2} g_{\nu\alpha} T \right) \xi^\alpha . \quad (6.39)$$

Consider now a spacelike hypersurface Σ^3 with unit future-directed timelike normal vector τ^μ . Since $\nabla^\mu \nabla^\nu K_{\mu\nu} = 0$, the right hand side of (6.39) is a covariantly conserved current, so that its integral over Σ^3 does not depend on the particular choice of hypersurface. This means that any spacetime with a Killing vector field ξ^μ defines a conserved charge (we shall call it the Komar charge)

$$\mathcal{K}_\xi = 16\pi G \int_{\Sigma^3} \left(T_{\mu\alpha} - \frac{1}{2} g_{\mu\alpha} T \right) \tau^\mu \xi^\alpha . \quad (6.40)$$

The conserved current defined by the right hand side of (6.39) is similar, but *different* from the previously known current $-T^{\mu\nu} \xi_\nu$ that we associated to a Killing vector. Both currents are conserved, but (6.39) has a crucial extra property from its construction as the divergence of an antisymmetric field, namely we can express its charge (6.40) as an integral over the boundary of the hypersurface, i.e. the celestial sphere. Applying the divergence theorem to $\nabla^\mu K_{\mu\nu}$, after pulling it back onto Σ^3 , one finds the boundary expression for the Komar charge

$$\mathcal{K}_\xi = - \int_{\Sigma^3} \nabla^\mu K_{\mu\nu} \tau^\nu = - \int_{S_\infty^2} K_{\mu\nu} r^\mu \tau^\nu = -2 \int_{S_\infty^2} \nabla_{[\mu} \xi_{\nu]} r^\mu \tau^\nu , \quad (6.41)$$

where r^μ is the outgoing unit normal to asymptotic 2-spheres that converge to S_∞^2 and is also orthogonal to τ^μ . Nothing in the discussion assumed that the Killing vector was timelike, so

we can apply this formalism to define Komar versions of the energy as well as the angular momentum contained in spacetime. Evaluating the surface integral at the celestial sphere for the case of the standard asymptotic metric (6.25) one finds $8\pi G M = \mathcal{K}_{\xi_t}$ and $16\pi G J = -\mathcal{K}_{\xi_\phi}$ where ξ_ϕ is the Killing vector for standard rotations in the azimuthal angle ϕ and J is the angular momentum about the polar axis. This fixes the normalization so that we may write the so-called Komar formulae

$$\begin{aligned} M &= 2 \int_{\Sigma^3} (T_{\mu\nu} - \tfrac{1}{2} g_{\mu\nu} T) \xi_t^\mu \tau^\nu = -\frac{1}{4\pi G} \int_{S_\infty^2} \nabla^{[\mu} \xi_t^{\nu]} r_\mu \tau_\nu , \\ J &= - \int_{\Sigma^3} (T_{\mu\nu} - \tfrac{1}{2} g_{\mu\nu} T) \xi_\phi^\mu \tau^\nu = \frac{1}{8\pi G} \int_{S_\infty^2} \nabla^{[\mu} \xi_\phi^{\nu]} r_\mu \tau_\nu . \end{aligned} \quad (6.42)$$

6.2 Gravitational Radiation

We consider now situations where the sources have a non-trivial time dependence, leading to *radiation*-type gravitational fields. We first assume that the energy-momentum distribution of the sources has compact support, and the radiation is analyzed *in vacuo*, far away from such sources. This means that the radiation field γ_{ab} solves the free wave equation far away from the sources, together with the transversality condition:

$$\partial^2 \gamma_{ab} = \partial^a \gamma_{ab} = 0 . \quad (6.43)$$

The gauge redundancy (6.3) acts on γ_{ab} as

$$\gamma_{ab} \rightarrow \gamma_{ab} + \partial_a \varepsilon_b + \partial_b \varepsilon_a - \eta_{ab} \partial^c \varepsilon_c , \quad (6.44)$$

and the transversality condition $\partial^a \gamma_{ab} = 0$ is preserved by all transformations of the form (6.44) with harmonic vector fields $\partial^2 \varepsilon_a = 0$. Hence, out of the six degrees of freedom left by (6.43) we may still eliminate four, leaving just two non-redundant degrees of freedom in γ_{ab} . We can easily demonstrate this in momentum space, since an arbitrary solution of $\partial^2 \gamma_{ab} = 0$ is a superposition of plane waves $\tilde{\gamma}_{ab}(p) \exp(ipx) + \text{c.c.}$ with $p^2 = 0$. Choosing the frame so that $p = (\omega, 0, 0, \omega)$ the transversality conditions reduce to

$$p^a \tilde{\gamma}_{ab} = 0 = \omega(\tilde{\gamma}_{0b} + \tilde{\gamma}_{3b}) . \quad (6.45)$$

The gauge redundancy (6.3) in momentum space reads

$$\tilde{h}_{ab} \rightarrow \tilde{h}_{ab} + ip_a \tilde{\varepsilon}_b + ip_b \tilde{\varepsilon}_a . \quad (6.46)$$

The four components $\tilde{\varepsilon}_b$ can be used to set $\tilde{h}_{0a} = 0$ which, together with (6.45) may be used to enforce the vanishing of all components of $\tilde{\gamma}_{ab}$ and $\tilde{h}_{ab}$ except for the two-dimensional submatrix with components along the (12)-plane, subject to the extra condition of vanishing trace. We refer to such a standard form of the fluctuating metric tensor as the ‘transverse-traceless’ gauge, $h_{ab}^{\text{TT}} = \gamma_{ab}^{\text{TT}}$. Such a matrix can be written as a linear combination of the Pauli matrices σ_3 and σ_1 . The component proportional to σ_3 is called the (+) wave polarization, whereas the component proportional to σ_1 is usually denoted ($\times$) and corresponds to the same polarization after a 45-degree rotation.

6.2.1 Generation

We can estimate the typical amplitude of a gravitational wave detected far away from a source. We begin by noticing that, because of the transversality constraint $\partial^a \gamma_{ab} = 0$, the time-dependent part of γ_{0a} is determined in terms of the time-dependence of the purely spatial ones γ_{ij} , so we concentrate on the retarded solution for those:

$$\gamma_{ij}(t, \vec{x}) = 4G \int d^3y \frac{T_{ij}(t - |\vec{x} - \vec{y}|, \vec{y})}{|\vec{x} - \vec{y}|} . \quad (6.47)$$

Evaluating this integral in the monopole approximation we find

$$\gamma_{ij}(x) \approx \frac{4G}{|\vec{x}|} \int_{\mathbb{R}^3} T_{ij} \Big|_{\text{retarded}} \quad (6.48)$$

with the retarded time given by $t_r = t - |\vec{x}|$. In this expression, we are neglecting the effects of the dipole corrections to the retarded time in (6.47), i.e. the terms linear in $\vec{y}$ in

$$t - |\vec{x} - \vec{y}| = t - |\vec{x}| + \frac{\vec{x} \cdot \vec{y}}{|\vec{x}|} + O(y^2)$$

These terms arise in the Taylor expansion of the energy-momentum tensor as

$$T_{ij}(t - |\vec{x} - \vec{y}|, \vec{y}) = T_{ij}(t_r, \vec{y}) + \frac{\vec{x} \cdot \vec{y}}{|\vec{x}|} \partial_t T_{ij}(t_r, \vec{y}) + \dots$$

The time derivative of T_{ij} determines the frequency of the radiation, so that this term is negligible in the ‘small emitter limit’, $\ell/\lambda \ll 1$, namely the radiation has long wavelength compared to the size ℓ of the source.

Using now the conservation equation $\partial^b T_{ba} = -\partial_t T_{0a} + \sum_k \partial_k T_{ka} = 0$ and repeated integration by parts we can obtain the identity

$$\int_{\mathbb{R}^3} T_{ij} = \frac{1}{2} \partial_t^2 \int_{\mathbb{R}^3} T_{00} y_i y_j ,$$

where we neglect all surface integrals on account of the compact support of the sources’ energy-momentum. Hence we finally obtain

$$\gamma_{ij}(t, \vec{x}) = \frac{2G}{|\vec{x}|} \frac{d^2 \mathcal{Q}_{ij}}{dt^2} \Big|_{\text{retarded}} , \quad (6.49)$$

where the derivatives are evaluated at the ‘point-like’ retarded time $t_r = t - |\vec{x}|$ and we have defined

$$\mathcal{Q}_{ij} \equiv \int_{\mathbb{R}^3} T_{00} y_i y_j , \quad (6.50)$$

the quadrupole moment of the source’s energy density. We thus conclude that gravitational waves in the low-frequency approximation, $\omega \ell \ll 1$, are generated by the quadrupole mass distribution, unlike electromagnetic waves, which are generated by charge dipole moments. This fact is largely responsible for the smallness of gravitational radiation. Since the quadrupole moment is of order $\mathcal{Q} \sim M \ell^2$ for a system of mass M and size ℓ , the metric perturbation at large distances is of order

$$h \sim \frac{GMv^2}{|\vec{x}|} , \quad (6.51)$$

so that we have the usual relativistic suppression of order v^2 with respect to the static Newtonian term.

Radiated Power

The energy carried off by such gravitational waves can be captured by measuring the flux through a large sphere at large distances from the source. In particular, the rate of energy loss is given by the *luminosity*

$$L_G = \frac{dE_{\text{rad}}}{dt} = \int_{S_\infty^2} \sum_i n_i t^{0i}, \quad (6.52)$$

where $n_i = x_i/|\vec{x}|$ is the unit radial vector on the celestial sphere S_∞^2 .

To estimate (6.52) we must evaluate t^{0i} by expanding (6.30) to second order in the metric perturbation h_{ab} . This leads to very complicated expressions, but the physical content may be exposed by a series of approximations. First, we consider a TT wave traveling in the x^3 direction, i.e. depending on just two functions $\gamma_+(t - x^3)$ and $\gamma_\times(t - x^3)$ and evaluate a coarse-grained version of t^{03} by averaging over observation times that are long compared to the inverse characteristic frequency of the wave. This results in a fair amount of cancellation, leaving a rather simple expression in terms of the two TT polarizations:

$$t^{03} \Big|_{3\text{-plane \& time av}} = \frac{1}{16\pi G} \left((\partial_t \gamma_+)^2 + (\partial_t \gamma_\times)^2 \right) \Big|_{3\text{-plane \& time av}}. \quad (6.53)$$

For the particular wave we are considering, we can write the same expression in the more symmetric fashion

$$t^{03} \Big|_{3\text{-plane \& time av}} = \frac{1}{32\pi G} \sum_{ij} \partial_t \gamma_{ij}^{\text{TT}} \partial_t \gamma_{ij}^{\text{TT}} \Big|_{3\text{-plane \& time av}}. \quad (6.54)$$

In taking the remaining average over all directions it is useful to remove the transversality constraint, since this obviously depends on the direction. The no-trace condition can still be enforced in a rotationally-invariant way by considering the trace-free matrix $\gamma_{ij}^{(s)} \equiv \gamma_{ij} - \frac{1}{3} \delta_{ij} \delta^{kl} \gamma_{kl}$ with five degrees of freedom. Hence, by substituting $\gamma^{(s)}$ in place of γ^{TT} in (6.54) we implement the averaging, up to a factor of 2/5, i.e.

$$\left\langle \sum_i n_i t^{0i} \right\rangle_{\text{sphere \& time av}} \approx \frac{2}{5} \cdot \frac{1}{32\pi G} \left\langle \sum_{ij} \partial_t \gamma_{ij}^{(s)} \partial_t \gamma_{ij}^{(s)} \right\rangle_{\text{sphere \& time av}}. \quad (6.55)$$

Finally, using the long-distance expression for the radiation field (6.49) and integrating over the celestial sphere we find for the luminosity

$$L_G = \frac{G}{5} \sum_{i,j} \left\langle \frac{d^3 \mathcal{Q}_{ij}^{(s)}}{dt^3} \frac{d^3 \mathcal{Q}_{ij}^{(s)}}{dt^3} \right\rangle_{\text{coarse \& retarded}}, \quad (6.56)$$

where $\mathcal{Q}_{ij}^{(s)} = \mathcal{Q}_{ij} - \frac{1}{3} \delta_{ij} \delta^{kl} \mathcal{Q}_{kl}$ is the traceless quadrupole moment of the matter distribution. The power radiated at a typical frequency $\omega \sim v/\ell$ by a system of typical masses M is of order

$$L_G \sim GM^2 \ell^4 \frac{v^6}{\ell^6} \sim \frac{G^4 M^5}{\ell^5} \sim M \frac{v^8}{\ell}, \quad (6.57)$$

A useful characterization of amount of energy loss in gravitational waves is the following. Given the luminosity power $L_G = dE_{\text{rad}}/dt$, we can consider the amount of energy lost in the time

that a single wave crest is emitted, $\omega^{-1} \sim \ell/v$, i.e. $L_G \ell/v$ and compare it with the gravitational self-energy, which is of order $|E_G| \sim GM^2/\ell$. Then, we find

$$\frac{\omega^{-1} \cdot L_G}{|E_G|} \sim v^5 \sim \phi^{5/2}, \quad (6.58)$$

where ϕ is the average value of the gravitational potential. If the system is not too relativistic, most of the self-energy loss can be interpreted as the ‘decay’ of typical orbits,

$$L_G = \frac{dE_{\text{rad}}}{dt} = -\frac{dE_G}{dt} \approx -\frac{GM^2}{\ell^2} \frac{d\ell}{dt}.$$

So the system ‘shrinks’ by gravitational wave emission at a rate

$$\frac{d\ell}{dt} \approx \frac{L_G \cdot \ell}{E_G} \sim -v^6 \sim -\phi^3. \quad (6.59)$$

Equivalently, the rate of period decay for internal motions is $dT/dt \sim -v^5$.

Suppose now that we have a binary system such as a pair of black holes or neutron stars in approximately circular orbit with $v^2 \sim GM/R$. Since the period is given by $T \sim GM/v^3$, we have $dT/dt \sim -v^5 \sim -(GM/T)^{5/3}$, which gives the law of decay of the period:

$$T \sim (GM)^{5/8} (t_c - t)^{3/8}. \quad (6.60)$$

This period decay implies a corresponding frequency increase, giving the gravitational wave signals from spiraling binaries a characteristic ‘chirp’.

The energy loss by gravitational radiation, proportional to $\phi^{5/2}$, is quite small unless we have sufficiently relativistic systems. The famous binary pulsar PSR1913+16 discovered by Hulse and Taylor has provided such a system for decades now, fulfilling one of the most impressive tests of Einstein’s theory so far (down to the 10^{-3} level in a mildly relativistic regime). The era of gravitational wave astronomy started in 2015 with the first detection of gravitational waves by LIGO, coming from a distant (1300 million light years) binary black hole collision.

Problem: Gravitational Radiation from Binaries

Consider a binary with unequal masses m_1 and m_2 on a circular orbit, slowly shrinking because of gravitational wave emission.

- (a) Show that, at fixed orbital frequency, the radiated power and the rate of change of the frequency depend on the masses only through the *chirp mass*

$$M_{\text{chirp}} \equiv \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}}.$$

- (b) Integrate the frequency evolution. Obtain the exact form of the chirp law (6.60), the accumulated phase, and the number of cycles spent between two frequencies.
- (c) Compute the two polarizations h_+ and $h_\times$ seen by an observer at distance r whose line of sight makes an angle ι with the orbital angular momentum. Check that their energy flux, integrated over the sphere, reproduces the radiated power.

- (d) Which properties of the signal are *not* fixed by M_{chirp} ? Estimate the time spent and the number of cycles above 10 Hz for a binary neutron star and for a binary black hole similar to the first one detected by LIGO.

6.2.2 Detection

Gravitational waves are capable of transferring energy to any detector sensitive to gravitational fields. For example, the nonrelativistic limit of the geodesic deviation equation yields

$$\frac{d^2 \ell_j}{dt^2} \approx \sum_k \ell^k R_{j00k} \approx \sum_k \frac{1}{2} \frac{d^2 h_{jk}^{\text{TT}}}{dt^2} \ell^k, \quad (6.61)$$

so that gravitational waves induce non-zero curvature and produce accelerations on the relative positions $\vec{\ell}$ of test particles.

The characteristic stretching of a ball of test particles, measured by $\Delta \ell / \ell$, is proportional to the amplitude of the metric perturbation $|h_{ij}|$. In Michelson-type interferometers such as LIGO, the test particles are heavy suspended mirrors whose distance is monitored by a laser beam. One has two perpendicular arms of equal length and a laser beam that is split between two components, each one of them traveling through one of the arms, and subsequently recombined for interference. Any relative variation in the distance between the two arms is detected through a variation of the interference pattern.

We can describe the salient features of such LIGO-like signals by solving the equation (6.61) in the realistic approximation that $|h|$ is very small. Then, we can approximate it by

$$\frac{d^2 \ell_j}{dt^2} \approx \sum_k \frac{1}{2} \frac{d^2 h_{jk}^{\text{TT}}}{dt^2} \ell_0^k, \quad (6.62)$$

where ℓ_0^k is the initial equilibrium separation. This leads to the simple solution

$$\ell_j(t) \approx \sum_k \left(\delta_{jk} + \frac{1}{2} h_{jk}^{\text{TT}}(t) \right) \ell_0^k. \quad (6.63)$$

Let us now consider the two arms of an interferometer with very similar (but not necessarily equal) lengths L_1 and L_2 and furthermore adapt the reference frame along their orthogonal directions. The variation of the arm lengths induced by the passing gravitational wave is given by

$$\frac{\delta L_1}{L_1} = \frac{1}{2} h_{11}(t), \quad \frac{\delta L_2}{L_2} = \frac{1}{2} h_{22}(t), \quad (6.64)$$

where we suppress the TT label on the metric perturbation. Considering now a laser pulse going through the first arm and back, it will accumulate a phase $\Delta \phi_1 \approx 2\omega L_1(1 + \frac{1}{2} h_{11}(t))$, and similarly for the pulse along the second arm $\Delta \phi_2 \approx 2\omega L_2(1 + \frac{1}{2} h_{22}(t))$, with ω the angular frequency of the light. Here we are assuming for simplicity that the travel time along the arms is much shorter than the period of the gravitational wave, namely $\omega_{GW} L \ll 1$. Suppose we arrange the apparatus so that the interference is completely destructive when $L_1 = L_2$ and in the absence of a gravitational wave. Then the amplitude after recombination must be proportional to

$$A \propto e^{i\Delta \phi_1} - e^{i\Delta \phi_2}, \quad (6.65)$$

so that it vanishes for trivial phase displacements. Then, the intensity of light recorded after recombination is proportional to

$$|A|^2 \propto |e^{i(\Delta\phi_1 - \Delta\phi_2)} - 1|^2 \approx |\Delta\phi_1 - \Delta\phi_2|^2 \approx 4\omega^2(L_1 - L_2)^2 + 4\omega^2L(L_1 - L_2)(h_{11}(t) - h_{22}(t)), \quad (6.66)$$

where $L = (L_1 + L_2)/2$ and we have neglected the term $\omega^2L^2(h_{11} - h_{22})^2$, quadratic in the extremely small gravitational wave amplitude. This output measurement allows us to read the modulation proportional to $h_{11}(t) - h_{22}(t)$ and literally draw a picture in real time of this combination of gravitational wave amplitudes. We see the convenience of keeping $L_1 - L_2$ small but non-zero, so that the modulated term is linear in the metric perturbation. Had we arranged for perfect cancellation in the absence of a gravitational wave, by choosing $L_1 = L_2$, the modulated term would have been quadratic in the gravitational wave amplitude, an extremely small number.




Figure 6.1: (a) Sketch of a Michelson interferometer, seen from above, with a gravitational wave propagating along x^3 , orthogonal to the plane of the arms. A laser beam is split into the two arms, of lengths L_1 and L_2 , by the beam splitter. The two beams are reflected back by the suspended end mirrors, which act as test masses, and recombined towards the photodetector. The input mirrors turn each arm into a Fabry-Pérot cavity, which effectively lengthens the arms. The dashed mirrors show the displaced positions for a ‘+’ wave with $h_{11} = -h_{22} > 0$, with $\delta L_{1,2}$ given by (6.64) (displacements exaggerated). The modulated output intensity is the term linear in h in (6.66). (b) A ring of free test particles in the (x^1, x^2) plane during one period of a wave $h_{+, \times} \propto \cos \omega t$, following (6.63), with the two arms superimposed. The dashed circle is the unperturbed ring. A ‘+’ wave stretches one arm while compressing the other, as in (a) at $\omega t = 0$. A ‘ $\times$ ’ wave only rotates the arms and leaves their lengths unchanged at first order, so an interferometer with arms along x^1 and x^2 is blind to this polarization.

We see that the signal is better the longer the arms (provided they are not much larger

than ω_{GW}^{-1}), a fact that justifies the use of Fabry–Pérot cavities that effectively make the arms longer by arranging for many reflections of the laser beam before recombination. At any rate, we see that such an interferometer has some blind directions, namely those for which the wave components along the two arms cancel one another, as seen in the frame of the interferometer.

This is an extremely simplified account of the famous measurements by Advanced LIGO which discovered gravitational waves in 2015, an event that could only be understood in terms of merger event of black holes of about 30 solar masses, occurring about 1300 million light-years away, thus inaugurating the era of gravitational wave astronomy. The stretching sensitivity that was achieved in order to perform this feat is a staggering $\delta L/L \sim |\Delta h| \sim 10^{-21}$.

Chapter 7

Cosmological Spacetimes

As we map the universe on scales of order Gpc or higher, we see no preferred direction in space and all our observations of the distribution of galaxies, and the properties of the Cosmic Microwave Background (CMB), are consistent with the idea that we sit at no privileged place in space, namely the universe looks *homogeneous* and *isotropic* on those large scales. Galaxy clusters are ‘standard observers’ which ‘see’ the universe as isotropic and no particular cluster is special. Below scales of a few hundred Mpc, galaxies may be ‘falling’ predominantly in certain directions inside the cluster. The discussion of cosmology starts with the characterization of spacetime geometries that are compatible with these constraints.

7.1 FLRW spacetimes

The mathematical characterization of homogeneity and isotropy is that the spacetime admits a foliation by spacelike hypersurfaces Σ_t on which there is an isometry group mapping every point to any other point, and moreover, around a given point P , no preferred vector field exists on the tangent space of Σ_t at P . Geodesics orthogonal to Σ_t are those of ‘standard observers’ such as galaxy clusters. Choosing the proper time of these standard observers as the time coordinate, i.e. Gaussian normal coordinates, we have $ds^2 = -dt^2 + d\ell_{\Sigma_t}^2$. We have already seen global de Sitter spacetime with metric

$$ds_{\text{dS}}^2 = -dt^2 + R^2 \cosh^2(t/R) d\ell_{\mathbb{S}^3}^2 . \quad (7.1)$$

as a prototype of a spacetime whose spatial sections are *homogeneous* and *isotropic*. Here, Σ_t are 3-spheres of radius $R(t) = R \cosh(t/R)$. Thus, one considers the natural generalization of (7.1) that retains the homogeneity and isotropy properties, resulting in the so-called Friedmann–Lemaître–Robertson–Walker (FLRW) spacetimes,

$$ds^2 = -dt^2 + R(t)^2 d\ell_{\mathbf{K}}^2 , \quad (7.2)$$

where $d\ell_{\mathbf{K}}^2$ stands for the spatial metric of a three-manifold that satisfies the criteria of homogeneity and isotropy and has fixed units of curvature. The overall size and curvature of this manifold is controlled by the function $R(t)$ which is to be determined by Einstein’s equations. It can be shown that the spatial section of constant curvature and homogeneity can only be one of three options: a 3-sphere in the case of positive curvature, a 3-hyperboloid in the case of

negative curvature, and flat $\mathbb{R}^3$ for zero curvature. We can write the three types of metric at once by the following polar parametrization

$$d\ell_{\mathbf{K}}^2 = \frac{dr^2}{1 - Kr^2} + r^2 d\Omega^2, \quad (7.3)$$

where $K = 1$ for the positive curvature $\mathbf{S}^3$, $K = 0$ for flat $\mathbb{R}^3$, and $K = -1$ for the 3-hyperboloid $\mathbf{H}^3$ of negative curvature. Homogeneity and isotropy of $\mathbb{R}^3$ is obvious. The other two 3-manifolds of constant curvature can be obtained by the equations for the 3-surface

$$Kx_1^2 + x_2^2 + x_3^2 + x_4^2 = K, \quad (7.4)$$

embedded in $\mathbb{R}^4$ with metric $ds_{\mathbb{R}^4}^2 = Kdx_1^2 + dx_2^2 + dx_3^2 + dx_4^2$. This confirms the homogeneity of $\mathbf{S}^3$ and $\mathbf{H}^3$.

The case $K = 1$ yields model universes like dS, in that the spatial sections are compact, whereas the models $K = 0, -1$ correspond to non-compact (so-called *open*) universes. What particular model is realized in nature, as well as the precise features of the ‘size’ function $R(t)$ depends on the Einstein equations.

It is important to notice that our ‘FLRW parametrization’, given by the metrics (7.2) and (7.3), may suffer from some ambiguities when it comes to global issues. A famous example is that of the ‘steady state metric’,¹ one of the presentations of de Sitter spacetime:

$$ds^2 = -d\bar{t}^2 + e^{2\bar{t}/R} d\vec{y}^2, \quad (7.5)$$

where $\vec{y} \in \mathbb{R}^3$. This is a particular case of FLRW with $K = 0$. However, there is change of variables revealing that it is identical to the $z_0 + z_4 > 0$ domain of the complete dS manifold. Hence, a FLRW manifold with $K = 0$ can be a ‘patch’ of another FLRW manifold with $K = 1$.

Problem: The Many Faces of de Sitter

Find proper domains of the global de Sitter manifold and appropriate coordinate systems that make them into FLRW cosmologies with $K = 0$ (flat de Sitter or ‘steady-state’ metric) and $K = -1$ (open de Sitter).

7.2 The Standard Cosmological Model

A good physical model of the universe in different epochs of its history is given in terms of three components: pressure-less matter, or ‘dust’, and a cosmological constant as a dominant component today, and radiation as a dominant component in the past. All situations can be modeled as a perfect fluid with energy momentum tensor $T_{00} = \rho$ and $T_{ij} = pg_{ij}$, with $p = 0$ for matter, $p = \rho/3$ for radiation, and $p = -\rho$ for vacuum energy (cosmological constant). We can summarize the three components by the equation of state $p = w\rho$, with three values of w .

¹The *steady state* universe of Gold, Hoyle and Bondi was once a fashionable model of cosmology, now long forgotten since the discovery of the cosmic microwave background radiation.

If the components or ‘species’ are approximately non-interacting, we can write $\rho = \sum_s \rho_s$ and $p_s = w_s \rho_s$ for each component.

Then, the Einstein’s equations can be reduced to the time-time component (so-called Friedmann equation)

$$\left(\frac{dR}{dt}\right)^2 + K = \frac{8\pi G}{3} \rho R^2, \quad (7.6)$$

and the space-space components

$$\frac{d^2 R}{dt^2} = -\frac{4\pi G}{3}(\rho + 3p) R. \quad (7.7)$$

The second equation can be traded for that of local energy conservation, also equivalent to the Bianchi identities $\nabla_\mu T^{\mu\nu} = 0$, or

$$\frac{d}{dt}(\rho R^3) = -p \frac{d}{dt}(R^3). \quad (7.8)$$

This equation is valid for each component separately, provided they are non-interacting. Using $p_s = w_s \rho_s$ we can solve it as $\rho_s \propto R^{-3(1+w_s)}$. The Friedmann equation is usually written in terms of the Hubble parameter $H \equiv R^{-1} dR/dt$, as

$$H^2 = -\frac{K}{R^2} + \frac{8\pi G}{3} \rho. \quad (7.9)$$

Another useful definition is the so-called density parameter

$$\Omega_s = \frac{8\pi G \rho_s}{3H^2}, \quad \Omega = \sum_s \Omega_s, \quad (7.10)$$

so that $\Omega > 1$ corresponds to $K = +1$, $\Omega < 1$ corresponds to $K = -1$, whereas $\Omega = 1$ describes a flat universe with $K = 0$. At present, Ω is believed to be well approximated by the sum of three components:

$$\Omega_0 = \Omega_b + \Omega_{\text{dm}} + \Omega_{\text{de}}, \quad (7.11)$$

where $\Omega_b \approx 0.05$ is the component in baryonic matter, $\Omega_{\text{dm}} \approx 0.27$ is the dark matter component and $\Omega_{\text{de}} \approx 0.68$ is the ‘cosmological constant’ component. The big surprise of the last decades is the fact that Ω_{de} is actually dominant. Hence, $\Omega_0 \approx 1$ within experimental errors, and the universe appears to have flat, $K = 0$ spatial sections.

Inserting the solution of the energy conservation equation $\rho_s = \rho_{s,0}(R_0/R)^{3(1+w_s)}$ into the Friedmann equation, we may recast it in the form

$$\frac{1}{2} \left(\frac{da}{dt}\right)^2 + V_{\text{eff}}(a) = 0. \quad (7.12)$$

where $a = R/R_0$, the quantities with zero subscript denoting the “present” values. The Friedmann equation is thus formally equivalent to a mechanical zero-energy motion of a particle in a potential

$$V_{\text{eff}}(a) = \frac{K}{2R_0^2} - \sum_s \frac{4\pi G \rho_{s,0}}{3} \frac{1}{a^{1+3w_s}} = \frac{H_0^2}{2} \left(\Omega_0 - 1 - \sum_s \frac{\Omega_{s,0}}{a^{1+3w_s}} \right). \quad (7.13)$$

The regions where $V_{\text{eff}} < 0$ determine the range of $R(t)$. In general, if one considers all three types of ‘species’ to be present with non-zero values (positive energy density of nonrelativistic matter and radiation and positive or negative vacuum energy), one sees that radiation always dominates near $R = 0$, whereas the cosmological constant controls the large R behavior, the ‘dust’ and ‘curvature’ components only shaping the transient effects at intermediate sizes. For $\Lambda < 0$, the effective potential V_{eff} is monotonically increasing and the universe always bounces between a *bang* and a *crunch*, quite independently of the sign of K and the ordinary matter content. For $\Lambda > 0$, the potential V_{eff} has always a “ Λ ” shape (a single local maximum) and the qualitative behavior depends on whether the potential is positive, zero or negative at the maximum. This is controlled by the actual values of the relative energy densities, as well as the sign of K . If $\max(V_{\text{eff}}) > 0$, one possibility is that the universe expands to a maximum size and bounces back to a crunch, but another possibility is that it starts contracting from $R = \infty$ in the infinite past down to a minimum size and then re-expands again, a deformation of the dS solution. The famous static Einstein universe that started modern cosmology in 1917 corresponds to $\max(V_{\text{eff}}) = 0$ and it is clearly unstable. Finally, if $K \leq 0$ (and still $\Lambda > 0$) then $\max(V_{\text{eff}}) < 0$ and the universe expands forever from zero size to infinite size. This picture may have some important variations if we remove (with fine tuning) the two asymptotically dominating species (radiation at small R and/or vacuum energy at large R).

A useful physical parameter that can be directly related to observations is the *deceleration* parameter

$$q \equiv -\frac{R d^2 R/dt^2}{(dR/dt)^2} = \frac{1+3\bar{w}}{2} \Omega, \quad (7.14)$$

where we have used (7.7) and defined the average equation-of-state parameter $\bar{w} = p/\rho$. For positive values of Ω , the expansion of the universe will accelerate if $\bar{w} < -1/3$, and will decelerate otherwise. Splitting in terms of species, we have

$$q = \sum_s \frac{1+3w_s}{2} \Omega_s. \quad (7.15)$$

At present, $q_0 \approx \frac{1}{2} \Omega_{\text{matter}} - \Omega_{\text{de}}$, and the measurement of q_0 supports our determination of Ω_{de} as a dominant component.

The connection with observation depends on the phenomenon of cosmological *red shift*. There are many ways of obtaining this very basic relation. Here we just give a shortcut based on the observation that (7.2) is conformally related to a static spacetime, i.e. defining a new time variable $\eta = \int^t dt'/R(t')$ we can write any Robertson–Walker spacetime as

$$ds^2 = R(\eta)^2 (-d\eta^2 + d\ell_{\mathbf{K}}^2).$$

Hence, frequencies $\omega \sim d/dt$ are related to frequencies measured in the static spacetime $d/d\eta$ by a factor of $R(t)$, which is the only overall time-dependent scale of the spacetime. An equivalent argument in terms of particle paths is the following: if two photons are sent from point P_1 to point P_2 with a conformal time separation $\Delta\eta_1$ at emission, they arrive at P_2 with the same conformal time separation, $\Delta\eta_2 = \Delta\eta_1$, since the null geodesics can be computed in the *static* metric $d\tilde{s}^2 = -d\eta^2 + d\ell_{\mathbf{K}}^2$. Then the corresponding proper times are related by

$$\frac{\Delta\tau_1}{\Delta\tau_2} = \frac{R(\eta_1)}{R(\eta_2)}.$$

Thus, we have the homogeneity property $\lambda(t) \sim R(t)$ for the wavelength of any electromagnetic wave that solves Maxwell's equations in the FLRW metric. Defining the redshift coefficient z in terms of the relative wavelength increase between emission and reception, we find

$$z \equiv \frac{\lambda_0 - \lambda_e}{\lambda_e} = \frac{\lambda_0}{\lambda_e} - 1 = \frac{R_0}{R_e} - 1 . \quad (7.16)$$

Furthermore, using the linear approximation of $R(t)$ for nearby galaxies, $R(t_0) \approx R_e + \frac{dR}{dt}(t_0 - t_e)$. Since the proper distance to nearby galaxies is given by $D \approx t_0 - t_e$, we find the approximate law

$$z \approx H_0 D , \quad H_0 \approx 0.23 \text{ Gpc}^{-1} , \quad (7.17)$$

originally tested by Hubble when the expansion of the universe was discovered back in 1929 (in the units used commonly by astronomers, $H_0 \approx 67\text{--}73 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the spread reflecting the current ‘Hubble tension’ between CMB and distance-ladder determinations). A more careful measurement of z for far away galaxies allows us to determine H_0 (to linear order) and q_0 (to first order in deceleration), determining the type of FLRW model that describes our universe.

Problem: Hubble's Law

Derive a more precise version of Hubble's law that is sensitive to the deceleration parameter, q_0 , in terms of the *luminosity distance* $D_L^2 = L/4\pi F$, where L is the absolute luminosity of a source and F is the flux of light at reception. Using the definition of luminosity distance and the geodesic equation for photons in a FLRW model, obtain the relation

$$D_L = H_0^{-1} \left(z + \frac{1}{2} (1 - q_0) z^2 + \dots \right) ,$$

which determines H_0 and q_0 from the directly measurable z and D_L .

Horizon and Flatness Problems

FLRW models starting with a *bang* at $t = 0$ pose the following question: is it possible that not all of $\mathbf{K}$ comes into causal contact in the time elapsed since the bang? This of course depends on the time passed since $t = 0$, the rate of expansion and the geometry of $\mathbf{K}$. The trajectories of light rays satisfy $d\ell/d\eta = \pm 1$, where ℓ is a (comoving) distance measured with the metric $d\ell_{\mathbf{K}}^2$ and η is the conformal time defined in the previous subsection. Hence, the causal structure is essentially determined by that of an effective two-dimensional Minkowski spacetime parametrized by (η, ℓ) . There are horizons if the past-directed light cones from a point P do not cover the entire $\mathbf{K}$ before the ‘initial’ time η_* is reached.

Since

$$\eta - \eta_* = \int_0^t \frac{dt'}{R(t')} , \quad (7.18)$$

such ‘particle horizons’ will never occur if $\eta_* = -\infty$, i.e. if this integral diverges at the lower limit. Conversely, if $\eta_* > -\infty$ (such as all FLRW models that are dominated by radiation or dust near $t = 0$), then there is a horizon if $\eta_P - \eta_* < \ell_{\max}(\mathbf{K})$, where $\ell_{\max}(\mathbf{K})$ is the maximum

distance in $\mathbf{K}$ (measured with $d\ell_{\mathbf{K}}^2$) length covered by a light ray in $\mathbf{K}$ (this is only relevant for compact $\mathbf{K}$). There is also a global notion of horizon, associated to the whole future history of a particle worldline, called the ‘event horizon’: the boundary of the set of events that can ever send signals to the observer. It exists whenever the conformal time $\int^\infty dt/R(t)$ converges towards the future.




Figure 7.1: *Local conditions at points P and Q could not be causally connected in the time elapsed since the big bang.*

The behavior of (7.18) at the lower limit can be further elucidated by rewriting it in the form

$$\eta - \eta_* = \frac{1}{R_0} \int_0^a \frac{da'}{a' da'/dt} = \frac{1}{R_0} \int_0^a \frac{da'}{a' \sqrt{-2V_{\text{eff}}(a')}} .$$

Hence, η_* is finite if $|V_{\text{eff}}(a)|$ grows as a negative power of a as $a \rightarrow 0$ (recall that V_{eff} is always negative for the actual solutions). Examining the dependence of V_{eff} on a we see that horizons will be present for $\bar{w} > -1/3$, i.e. for any cosmology that starts decelerating right from the bang. On the other hand, for cosmologies that start out (marginally) accelerating, $\bar{w} \leq -1/3$, the bang occurs at minus infinity in conformal time, and there are no horizons.

Our model universe as we have described it so far, with radiation domination at early times, does have horizons. This poses the question of why we see a homogeneous universe after all, since no causal processes could achieve this homogeneity by standard local interactions. An interesting way out of this conundrum is the idea of *inflation*, consisting in the possibility that a phase of large accelerated expansion exists before the radiation-domination era. Such a phase gives a large contribution to (7.18) and thus pushes η_* to more negative values. This means that a phase of inflation can be used to remove horizons.

FLRW models with flat spatial sections correspond to $\Omega = 1$ at all times. We can ask how natural is this condition as an asymptotic behavior as $t \rightarrow 0$ or $t \rightarrow \infty$ in expanding models with a bang. From the previous equations we have

$$\Omega = 1 + \frac{K}{H^2 R^2} . \quad (7.19)$$

Using the definition of H we have $H^2 R^2 = (dR/dt)^2 = R_0^2 (da/dt)^2 = -2R_0^2 V_{\text{eff}}(a)$. Hence we find

$$\Omega - 1 = \frac{K}{2R_0^2 |V_{\text{eff}}(a)|} . \quad (7.20)$$

Since the monotonicity of V_{eff} is associated with the accelerating/decelerating character of the universe, we see that accelerating phases ($\bar{w} < -1/3$ or increasing $|V_{\text{eff}}|$) drive Ω to unity, whereas decelerating phases ($\bar{w} > -1/3$ or decreasing $|V_{\text{eff}}|$) drive Ω *away* from unity.

Our universe has been dominated by radiation and dust till relatively recently, i.e. it has been decelerating for most of its history. Hence, the fact that Ω_0 is so close to unity implies that it was much closer at early times. In the radiation era, the value of Ω was thus extremely *fine-tuned*, an initial condition without a clear explanation within the standard cosmological model. Inflation provides an interesting perspective on this so called *flatness problem*. Since accelerated universes have $\Omega = 1$ as a future attractor, an early phase of inflation is also able to dynamically tune $\Omega \approx 1$, setting the initial conditions of the radiation era.

Chapter 8

Black Holes

A black hole is a region of spacetime from which nothing, not even light, can escape to infinity. The simplest example is provided by the Schwarzschild solution (5.45) when it is continued inside its Schwarzschild radius. In this chapter we first study the global structure of the Schwarzschild spacetime, then argue that black holes are an unavoidable outcome of gravitational collapse, and generalize to charged and rotating holes. Black holes are now known to exist with masses ranging from stellar scale to billions of solar masses. They are central to the developments of astrophysics, cosmology and fundamental physics.

8.1 The Schwarzschild Black Hole

We write the Schwarzschild metric in terms of the Schwarzschild radius $r_s = 2GM$,

$$ds^2 = -\left(1 - \frac{r_s}{r}\right) dt^2 + \frac{dr^2}{1 - r_s/r} + r^2 d\Omega^2. \quad (8.1)$$

The Schwarzschild radius has an elementary Newtonian interpretation: the escape velocity from the surface of a spherical body of radius r , $v_{\text{esc}} = \sqrt{2GM/r}$, reaches the speed of light precisely at $r = r_s$. This led Michell and Laplace to speculate, back in the XVIII century, about the existence of ‘dark stars’. Recall that, by Birkhoff’s theorem, (8.1) is the unique vacuum solution with $SO(3)$ symmetry, the condition of stationarity being redundant. Hence, the exterior of a spherical star that pulsates, or even collapses, in vacuum is always described by the *static* metric (8.1).

There is an apparent singularity at $r = r_s = 2GM$, about 3 km for $M = M_\odot$. On the other hand, the local physics, as diagnosed by an infalling observer (a *frefo*), remains regular there. In particular, tidal effects are controlled by the Riemann tensor which, in such *frefo* frame, has components of order $R_{\mu\nu\rho\sigma} \sim r_s/r^3$. A coordinate-invariant measure is the Kretschmann scalar

$$R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} = \frac{12 r_s^2}{r^6}, \quad (8.2)$$

which is finite at $r = r_s$ and only singular at $r = 0$. The tidal acceleration across a body of size ℓ at the Schwarzschild radius is $a_{\text{tidal}} \sim \ell r_s/r^3|_{r=r_s} = \ell/r_s^2$, which is small for large black holes. Hence, $r = r_s$ is only a *coordinate* singularity of (8.1), whereas $r = 0$ is a genuine curvature singularity. In any case, something quite peculiar must happen for $r < r_s$, since g_{00} and g_{rr}

change sign, suggesting that the radial coordinate becomes a time-like coordinate, so that the singularity at $r = 0$ must be interpreted as a ‘moment’ in time, rather than a ‘place’ in space. A precise statement along these lines can be made by finding a coordinate chart that smoothly passes to the interior region. We will begin with a local analysis and later proceed to a more powerful global description.

Schwarzschild and Rindler

There is a simple way of exhibiting a *frefo* near the Schwarzschild radius, which is actually global over all the sphere at $r = r_s$. Denote $f(r) = 1 - r_s/r$ and expand to first order in a Taylor series:

$$f(r) = 2\kappa(r - r_s) + \dots, \quad \kappa = \frac{1}{2r_s} = \frac{1}{4GM}, \quad (8.3)$$

where we have neglected terms of quadratic order in $r - r_s$. The quantity κ is known as the ‘surface gravity’ and will be shortly interpreted as an inverse ‘relaxation’ time. The metric near r_s can be approximated as

$$ds^2 \approx -2\kappa(r - r_s)dt^2 + \frac{dr^2}{2\kappa(r - r_s)} + r_s^2 d\Omega^2, \quad (8.4)$$

or, defining the proper-radius coordinate $\ell = \sqrt{2(r - r_s)/\kappa}$, we find¹

$$ds^2 \approx -\kappa^2 \ell^2 dt^2 + d\ell^2 + r_s^2 d\Omega^2. \quad (8.5)$$

Metrically, this is the direct product of a sphere of radius r_s and two-dimensional Rindler spacetime, which coincides with (3.16) under $\kappa = g$ and $\kappa\ell = e^{\kappa\xi}$ (notice however that our t coordinate here maps to the η coordinate there). Therefore, we see that *fido* observers at fixed ℓ , or fixed r , have uniform Rindler acceleration of magnitude $g_\ell = 1/\ell = \sqrt{\kappa/2(r - r_s)}$. Since this acceleration diverges as $r \rightarrow r_s$, we see that one needs infinite forces to stay put at $r = r_s$, suggesting that the singularity at $r = r_s$ is just a coordinate one, whose physical interpretation is the mere impossibility of realizing physical static *fidors* at fixed- $r \rightarrow r_s$.

Rindler spacetime is just a reparametrized version of Minkowski, as we can make explicit by defining

$$X^\pm = T \pm X = \pm \ell e^{\pm \kappa t}, \quad (8.6)$$

in terms of which

$$ds^2 \approx -dX^+ dX^- + r_s^2 d\Omega^2 = -dT^2 + dX^2 + r_s^2 d\Omega^2. \quad (8.7)$$

The region $r \geq r_s$ of the near- r_s Schwarzschild spacetime corresponds to the intersection of the regions $X^+ \geq 0$ and $X^- \leq 0$. In principle, there is no difficulty in extending the geometry to the region $X^- > 0$ as would correspond to the free-fall trajectory of a particle in a radial plunge. Any such particle follows locally a trajectory $X^- = \alpha(X^+ - X_0^+)$ with $\alpha > 0$ in Minkowski coordinates, and has no trouble in continuing into the $X^- > 0$ region. Since this was a radial plunge, such continued trajectory must enter the $r < r_s$ ‘interior’.

¹A similar analysis obtains the same result for a general metric of the form $ds^2 = -f(r)dt^2 + dr^2/g(r) + r^2 d\Omega^2$ for any functions f, g having a non-degenerate zero at the same point, i.e. $f(r) = f'_s(r - r_s) + \dots$ and $g(r) = g'_s(r - r_s) + \dots$. Any such metric looks like Rindler near r_s with $2\kappa = \sqrt{f'_s g'_s}$.




Figure 8.1: The near-horizon region of the Schwarzschild geometry in the Minkowski coordinates (8.6). Each point represents a sphere of radius r_s , cf. (8.7). The exterior $r > r_s$ is the wedge $X^+ > 0$, $X^- < 0$, bounded by the horizon $r = r_s$ on the null lines $X^\pm = 0$. Static fiducials at fixed ℓ , or fixed r , follow the hyperbolae $X^+X^- = -\ell^2$, with an acceleration $1/\ell$ that diverges at the horizon. Lines of constant t go through the origin. A radially plunging particle, $X^- = \alpha(X^+ - X_0^+)$ with $\alpha > 0$, is a straight timelike line. It crosses the future horizon $X^- = 0$ at $X^+ = X_0^+$ and continues smoothly into the region $X^- > 0$, the interior $r < r_s$.

Solving for the radial plunge path as a function of the proper height $\ell(t)$ and the asymptotic time one finds

$$\ell(t) = \frac{X_0^+ e^{-\kappa t}}{1 + \alpha^{-1} e^{-2\kappa t}} \approx X_0^+ e^{-\kappa t}. \quad (8.8)$$

The finite-distance fall towards $r = r_s$ corresponds to an exponentially stretched time in the asymptotic time variable, with characteristic e-folding time κ^{-1} , giving the announced interpretation of κ as an inverse relaxation time.

Problem: Rindler and Regular Horizons

Consider a metric of the form

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{g(r)} + r^2 d\Omega^2$$

with f, g positive for $r > r_s$ and having a non-degenerate zero at $r = r_s$, i.e. $f(r) = f'_s(r - r_s) + \dots$ and $g(r) = g'_s(r - r_s) + \dots$. Show that any such metric looks near $r = r_s$ like the direct product of a sphere and two-dimensional Rindler with surface gravity $\kappa = \sqrt{f'_s g'_s}/4$.

8.1.1 The Kruskal Extension

An even more global description is possible by a clever change of variables due to Kruskal and Szekeres. This is suggested by the study of radial light rays, satisfying $ds^2 = d\Omega^2 = 0$. Writing

$$ds^2|_{d\Omega=0} = \left(1 - \frac{r_s}{r}\right) (-dt^2 + dr_*^2) , \quad dr_* = \frac{dr}{1 - r_s/r} , \quad (8.9)$$

we find the Regge–Wheeler *tortoise* coordinate

$$r_* = r + r_s \log \left(\frac{r}{r_s} - 1 \right) , \quad (8.10)$$

which maps $r \in (r_s, \infty)$ into $r_* \in (-\infty, +\infty)$, the horizon being sent to $r_* \rightarrow -\infty$. Radial null rays are simply lines at 45 degrees in the (t, r_*) plane. Introducing null coordinates $x^\pm = t \pm r_*$, the metric reads $ds^2 = -(1 - r_s/r) dx^+ dx^- + r^2 d\Omega^2$. Near $r \approx r_s$ we have $1 - r_s/r \approx e^{-1} \exp((x^+ - x^-)/2r_s)$, so that

$$ds^2 \approx -\frac{4r_s^2}{e} d \left(e^{x^+/2r_s} \right) d \left(-e^{-x^-/2r_s} \right) + r_s^2 d\Omega^2 , \quad (8.11)$$

which suggests the definition of the *Kruskal* coordinates

$$X^\pm = \pm 2r_s e^{\pm x^\pm/2r_s} . \quad (8.12)$$

In terms of these, the Schwarzschild metric takes the exact form (we use the same notation for $X^\pm$ as in the previous section, because they coincide near r_s)

$$ds^2 = -\frac{r_s}{r} e^{-r/r_s} dX^+ dX^- + r^2 d\Omega^2 , \quad (8.13)$$

where r is now regarded as a function of (X^+, X^-) , implicitly defined by

$$X^+ X^- = (2r_s)^2 \left(1 - \frac{r}{r_s} \right) e^{r/r_s} , \quad (8.14)$$

so that the curves of constant r are hyperbolae in the (X^+, X^-) plane. The Schwarzschild time is determined by

$$\frac{X^+}{X^-} = -e^{t/r_s} , \quad (8.15)$$

so that curves of constant t are straight lines through the origin. The metric (8.13) is manifestly regular at the Schwarzschild radius, which now corresponds to $X^+ = 0$ or $X^- = 0$, whereas the singularity $r = 0$ sits at the hyperbola $X^+ X^- = (2r_s)^2$.

The original Schwarzschild coordinates only cover the region I, defined by $X^+ > 0$, $X^- < 0$. The coordinates (X^+, X^-) can be extended to the whole domain $X^+ X^- < (2r_s)^2$, where (8.13) is a smooth solution of the vacuum Einstein equations. This *maximal extension* of the Schwarzschild spacetime contains four regions, depicted in Figure 8.2. Region III ($X^+ < 0$, $X^- > 0$) is a second copy of the asymptotically flat exterior. Region II ($X^\pm > 0$), with $r < r_s$, is the interior of the black hole, ending in the future singularity. Region IV ($X^\pm < 0$) is its time reverse, a *white hole* emanating from a past singularity. Since the metric (8.13) is conformal to $-dX^+ dX^-$ in the radial directions, radial light rays travel at 45 degrees in the Kruskal diagram,




Figure 8.2: The Kruskal diagram of the maximally extended Schwarzschild spacetime. Each point represents a two-sphere of radius $r(X^+, X^-)$. The diagonals $X^\pm = 0$ are the horizons $r = r_s$, curves of constant r are hyperbolae and lines of constant t go through the origin. The wiggly lines are the singularities at $r = 0$. The local radial light cone is shown in region I.

making the causal structure transparent. In particular, the lines $X^\pm = 0$ are null surfaces, the *horizons*, and any future-directed causal curve entering region II can never leave it: it necessarily ends at the singularity in finite proper time.

The $t = 0$ slice corresponds to the straight line $X^+ = -X^-$, crossing regions I and III through the origin. Its intrinsic geometry is that of two asymptotically flat spaces $\mathbb{R}^3$ connected by a ‘throat’ whose minimal sphere has radius r_s (see Figure 8.3). This so-called Einstein–Rosen bridge is *not* traversable in a causal way, since an observer leaving region I towards region III necessarily crosses the future horizon and ends at the singularity. On the other hand, two observers jumping from regions I and III can meet inside region II, before being destroyed at the singularity.

The global picture allows us to be more precise in the interpretation of κ as an inverse relaxation time. Let us consider an infalling antenna sending out photons in radial outgoing trajectories as it radially plunges towards the horizon at r_s . The photons travel outwards along a line of constant X^- , so that the interval ΔX^- between pulses can be read either in terms of the near-horizon quantities or the asymptotic ones. Let the proper time between pulses be $\Delta\tau = \Delta X^-/\sqrt{\alpha}$ for a near horizon trajectory $X^- = \alpha(X^+ - X_0^+)$. Since $X^- = -\kappa^{-1}e^{-\kappa x^-} = -\kappa^{-1}e^{-\kappa(t-r_*)}$, an asymptotic observer at fixed large radial distance with $r_* = R_* \sim R$ will receive the pulses separated by a Δt interval given by

$$\Delta t = \sqrt{\alpha} \Delta\tau e^{\kappa(t-R_*)} . \quad (8.16)$$

Equivalently, if $\Delta\tau \sim 1/\nu_e$ and $\Delta t \sim 1/\nu_r$ for ν_e and ν_r the emission and reception frequencies, we find

$$\nu_r(t) \propto \nu_e e^{-\kappa t} , \quad (8.17)$$




Figure 8.3: *Embedding diagram of the $t = 0$ slice of the Kruskal spacetime, showing the Einstein–Rosen bridge connecting the two asymptotic regions, I and III.*

an expression that shows in a transparent way how the surface gravity κ is interpreted as an inverse relaxation time: all processes occurring in $O(1)$ intervals of time for an infalling observer near the horizon, slow their rate exponentially fast, with characteristic time κ^{-1} , as seen by the asymptotic observer. For astrophysical black holes, this is quite a fast time scale in the ballpark of tens of microseconds, times $M/M_\odot$. This is the dynamical meaning of the usual statement that ‘black holes have no hair’, namely, any structures hovering near the horizon are absorbed in such a time scale, leaving a clean near-horizon geometry.

Killing Vectors and Trapped Surfaces

The Schwarzschild spacetime has four Killing vectors: three rotational ones ξ_Ω , generating $SO(3)$, and the time translation $\xi_t = \partial/\partial t$. The rotational Killing vectors are spacelike everywhere, whereas ξ_t satisfies

$$\xi_t^2 = g_{tt} = -\left(1 - \frac{r_s}{r}\right), \quad (8.18)$$

so that it is timelike in regions I and III and spacelike in regions II and IV. Hence, the geometry is *static* in the exterior regions but, in regions II and IV, where r becomes a timelike coordinate, the geometry is time dependent, rather like an anisotropic *cosmological* spacetime which collapses to (or expands from) a singularity. Since the worldline of a stationary observer must be tangent to a timelike Killing vector, there are no stationary observers in regions II or IV. Using (8.15) we find that, in Kruskal coordinates,

$$\xi_t = \frac{1}{2r_s} \left(X^+ \frac{\partial}{\partial X^+} - X^- \frac{\partial}{\partial X^-} \right), \quad (8.19)$$

which has exactly the form of a Lorentz boost in the (X^+, X^-) plane. We obtain again the result that, near the horizon, the Kruskal diagram looks just like Minkowski space, with the static observers in region I playing the role of the uniformly accelerated observers in the Rindler wedge (cf. Figure 3.1).

In our previous discussion of the near-horizon limit, we found that the surface gravity κ gave an important relation between local near-horizon rates and the asymptotic point of view. It is

interesting that the Killing vector ξ_t can be used to give a more invariant characterization of the surface gravity. By direct computation, we can check that κ satisfies

$$\xi_t^\mu \nabla_\mu \xi_t|_{r \rightarrow r_s} = \kappa \xi_t|_{r \rightarrow r_s}, \quad \kappa^2 = \lim_{r \rightarrow r_s} \partial_\mu N(\xi_t) \partial^\mu N(\xi_t), \quad (8.20)$$

where $N(\xi_t) = \sqrt{-\xi_t^2}$ is the norm of the Killing vector. We will find that these, more covariant definitions of κ , do generalize to other black holes beyond Schwarzschild.

A more invariant characterization of the black hole interior uses the notion of trapped surfaces. Consider a two-sphere S_r^2 at some point of the Kruskal diagram, and the two spherical light fronts emitted from it, one ingoing (γ_{in}) and one outgoing (γ_{out}). In region I, the area of the outgoing front increases towards the future while that of the ingoing front decreases, as expected in flat space. However, in region II the two fronts move to smaller values of r (see Figure 8.4), and both spherical fronts decrease their area towards the future:

$$A(S_{\text{in}}^2), A(S_{\text{out}}^2) < A(S_r^2).$$

Such a sphere is said to be *future trapped*. All spheres in region II are future trapped, whereas all spheres in region IV are *past trapped*. More generally, a compact spacelike two-surface is trapped when the expansions of both families of future-directed null geodesics orthogonal to it are negative.




Figure 8.4: Radial light rays emitted from a sphere S_r^2 outside the horizon, where the surfaces of constant r are timelike, and inside the horizon, where they are spacelike. In the second case, both light fronts move to smaller r and the sphere is trapped.

8.1.2 Global Causal Structure

The global causal structure of a spacetime is best visualized in terms of a device called a *Penrose diagram*, which brings infinity to a finite coordinate distance by means of a conformal transformation that preserves the light cones. For the Schwarzschild spacetime we define

$$X^\pm = 2r_s \tan \tilde{X}^\pm, \quad \tilde{X}^\pm \in (-\pi/2, \pi/2), \quad (8.21)$$

so that the metric takes the form

$$ds^2 = -f(\tilde{X}^+, \tilde{X}^-) d\tilde{X}^+ d\tilde{X}^- + r^2(\tilde{X}^+, \tilde{X}^-) d\Omega^2, \quad f = \frac{4r_s^3}{r} \frac{e^{-r/r_s}}{\cos^2 \tilde{X}^+ \cos^2 \tilde{X}^-}. \quad (8.22)$$

Since the conformal factor f is irrelevant for the causal structure, we can represent the spacetime by plotting the auxiliary two-dimensional metric $d\tilde{s}_{1+1}^2 = -d\tilde{X}^+ d\tilde{X}^-$, in which radial light rays still travel at 45 degrees, and all of infinity is mapped to the boundaries $\tilde{X}^\pm = \pm\pi/2$. The singularity at $X^+ X^- = 4r_s^2$ is mapped to $\tan \tilde{X}^+ \tan \tilde{X}^- = 1$ or, equivalently $\tilde{X}^- = \frac{\pi}{2} - \tilde{X}^+$, a straight spacelike line in the diagram (and similarly for the past singularity). The result is Figure 8.5. Each asymptotic region has a future null infinity $\mathcal{J}^+$, where outgoing light rays end, a past null infinity $\mathcal{J}^-$, where ingoing light rays originate, a spatial infinity i^0 and future and past timelike infinities $i^\pm$, where timelike geodesics end and originate, respectively.




Figure 8.5: Penrose diagram of the maximally extended Schwarzschild spacetime. The future and past horizons of region I are denoted $\mathcal{H}^\pm$. A curve of constant $r > r_s$ and a curve of constant t are shown in region I.

For comparison, the analogous construction for Minkowski spacetime starts from $r_* = r$ and $x^\pm = t \pm r$, which we compactify as

$$\frac{x^\pm}{\ell} = \tan \tilde{X}^\pm, \quad (8.23)$$

where ℓ is an arbitrary reference length scale. The metric becomes

$$ds^2 = \frac{\ell^2}{\cos^2 \tilde{X}^+ \cos^2 \tilde{X}^-} \left(-d\tilde{X}^+ d\tilde{X}^- + \frac{1}{4} \sin^2(\tilde{X}^+ - \tilde{X}^-) d\Omega^2 \right), \quad (8.24)$$

and the condition $r \geq 0$ restricts the diagram to $\tilde{X}^+ \geq \tilde{X}^-$, the origin $r = 0$ being the vertical line $\tilde{X}^+ = \tilde{X}^-$. The result is the triangle of Figure 8.6.

Cosmic Censorship

While unphysical, it is interesting, for illustration, to consider the Penrose diagram of a negative-mass Schwarzschild metric:

$$ds^2 = -\left(1 + \frac{r_s}{r}\right) dt^2 + \frac{dr^2}{1 + r_s/r} + r^2 d\Omega^2. \quad (8.25)$$




Figure 8.6: Penrose diagram of Minkowski spacetime. Each point represents a two-sphere, except for the left vertical boundary $r = 0$.

Now $g_{tt} < 0$ and $g^{rr} > 0$ for all $r > 0$, so there is no horizon, and the Kruskal step (8.12) is not needed: the Schwarzschild chart already covers the whole spacetime. (Formally, $r_s \rightarrow -r_s$ turns (8.14) into $X^+X^- = 4r_s^2(1 + r/r_s)e^{-r/r_s}$, which never vanishes.) The tortoise coordinate

$$r_* = r - r_s \log \left(1 + \frac{r}{r_s} \right), \quad r_* \simeq \frac{r^2}{2r_s} \quad (r \ll r_s), \quad (8.26)$$

maps $r \in (0, \infty)$ onto $r_* \in (0, \infty)$. In contrast with (8.10), where the horizon is pushed to $r_* \rightarrow -\infty$, the point $r = 0$ now sits at a *finite* value of r_* . We can therefore compactify $x^\pm = t \pm r_*$ exactly as for Minkowski spacetime, $x^\pm = r_s \tan \tilde{X}^\pm$, which gives the analog of (8.22)

$$ds^2 = -\frac{r_s^2(1 + r_s/r)}{\cos^2 \tilde{X}^+ \cos^2 \tilde{X}^-} d\tilde{X}^+ d\tilde{X}^- + r^2(\tilde{X}^+, \tilde{X}^-) d\Omega^2. \quad (8.27)$$

The condition $r \geq 0$ is again $\tilde{X}^+ \geq \tilde{X}^-$, and the diagram is the Minkowski triangle (Figure 8.7). The crucial difference is that the vertical boundary $\tilde{X}^+ = \tilde{X}^-$ is now a curvature singularity, with $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} = 12r_s^2/r^6$ and a diverging conformal factor in (8.27). Being timelike and not hidden by any horizon, it is a *naked* singularity. Ingoing radial light rays from $\mathcal{J}^-$ reach it at finite affine parameter, and outgoing rays emitted arbitrarily close to it reach $\mathcal{J}^+$. The negative mass causes ordinary positive-mass particles of energy $E = m\varepsilon$ to be *repelled* by the singularity, bouncing at $r_{\min} = r_s/(\varepsilon^2 - 1)$.

Such a geometry cannot be the exterior of matter satisfying the dominant energy condition, since the positive energy theorem then guarantees $E_{\text{ADM}} \geq 0$. Although negative-mass Schwarzschild is a very unphysical metric, we will see that rotating black holes that are ‘over-spun’ beyond a certain limit would correspond to spacetimes sharing this same global structure,




Figure 8.7: Penrose diagram of the negative-mass Schwarzschild spacetime (8.25). The triangle coincides with that of Minkowski spacetime (Figure 8.6), but the left boundary $r = 0$ is now a timelike curvature singularity (wiggly line), visible from $\mathcal{J}^+$. Shown are a curve of constant r , a curve of constant t , an ingoing radial light ray ending on the singularity, an outgoing one emanating from it, and a radial timelike geodesic repelled at $r_{\min} = r_s/(\epsilon^2 - 1)$.

featuring ‘naked singularities’. The expectation that no such processes can take place starting from physical initial conditions is the content of the ‘cosmic censorship’ conjecture by Penrose.

8.2 Gravitational Collapse

The maximally extended Schwarzschild spacetime is an eternal black hole as seen from outside, with a white hole in its past and a second asymptotic region. We now show that these exotic features are absent in more realistic black holes formed by gravitational collapse.

8.2.1 Imploding Shells

The simplest model of gravitational collapse is an imploding spherical shell of photons, such as those produced by a spherical laser pulse converging on a point. By Birkhoff’s theorem, the spacetime outside the null shell is a portion of the Schwarzschild geometry with mass parameter given by the total ADM mass of the photon shell, whereas the spacetime inside the shell is flat (this is the relativistic version of Newton’s shell theorem). The complete spacetime is obtained by gluing a portion of the Minkowski Penrose diagram (the region before the shell arrives) to a portion of the Schwarzschild Penrose diagram (the region after the shell passes) along the null trajectory of the shell, which is fully known by symmetry and the invariant speed of light. The result is depicted in Figure 8.8.

The region of spacetime from which no signal can reach $\mathcal{J}^+$ is, by definition, the black hole. It lies ‘inside’ the *event horizon*, defined as the boundary of the causal past of future null infinity, $\mathcal{H}^+ = \partial[\text{Past}(\mathcal{J}^+)]$. Notice that the event horizon starts *in the flat region*, before




Figure 8.8: Penrose diagram of the collapse of a null spherical shell. The region inside the shell is flat, whereas the region outside the shell is Schwarzschild. The event horizon $\mathcal{H}^+$ is the dashed line, the trapped region is shaded.

the shell has passed, since its definition refers to the whole future history of the spacetime. The region between the horizon and the singularity, after the passage of the shell, is filled with trapped spheres. We define the *apparent horizon* as the boundary of the trapped region. In the case of a single shell, apparent and event horizons coincide in the Schwarzschild portion of the diagram. A key lesson is that the white hole and the parallel universe are gone! Only regions I and II of the maximally extended Schwarzschild spacetime survive in a physical collapse.

The difference between apparent and event horizons is illustrated in the case of two successive shells, of energies M_1 and M_2 (Figure 8.9). The apparent horizon jumps discontinuously, from zero to $r = 2GM_1$ when the first shell passes, and from $r = 2GM_1$ to $r = 2G(M_1 + M_2)$ when the second shell passes. On the other hand, the event horizon grows continuously and monotonically, anticipating the future arrival of the shells. Choosing a family of Cauchy surfaces Σ_τ , we may define the horizon area as a function of time,

$$A_{\mathcal{H}}(\tau) = \text{Vol}(\mathcal{H} \cap \Sigma_\tau) .$$

A famous theorem by Hawking states that the area of the event horizon never decreases,

$$\delta A_{\mathcal{H}^+} \geq 0 , \quad (8.28)$$

provided the matter satisfies the null energy condition (and there are no naked singularities)². Both the event and the apparent horizon areas can decrease when the null energy condition is violated, as happens with the quantum effects discussed below (Hawking radiation). Classically,

²The null energy condition states that the matrix elements of the matter energy-momentum tensor along null directions are non-negative, i.e. $T_{\mu\nu}k^\mu k^\nu \geq 0$ for any null k^μ

the apparent horizon can jump discontinuously (for ideal, delta-function shells), whereas the event horizon grows continuously. If the black hole settles down to a stationary state, so that there is a timelike Killing vector ξ_t outside, the apparent horizon approaches the event horizon.




Figure 8.9: Collapse of two successive null shells. Left: the event horizon (thick gray line) and the apparent horizon (dotted line). Right: area of the event horizon and the apparent horizon as a function of time.

Hence, in order to locate horizons in stationary spacetimes it is enough to locate apparent horizons, i.e. the topological spheres whose outgoing light front has marginal (vanishing) expansion. Given the areal radius r , the apparent horizon is where the surfaces of constant r become null (cf. Figure 8.4), i.e. where the gradient $\partial_\mu r$ is a null vector:

$$g^{\alpha\beta} \partial_\alpha r \partial_\beta r = g^{rr} = 0. \quad (8.29)$$

For the Schwarzschild metric, $g^{rr} = 1 - r_s/r$ and we recover $r = r_s$.

Summarizing, for the eternal Schwarzschild black hole we have the following equivalent characterizations of the horizons:

$$\begin{aligned} \mathcal{H}^+ &= \partial [\text{Past}(\mathcal{J}^+)] = \text{future event horizon}, \\ \mathcal{H}^- &= \partial [\text{Future}(\mathcal{J}^-)] = \text{past event horizon}, \\ \mathcal{H}^\pm &= \partial [\text{future/past trapped region}] = \text{future/past apparent horizon}, \quad g^{rr}|_{\mathcal{H}^\pm} = 0, \\ \mathcal{H}^+ \cup \mathcal{H}^- &= \partial [\text{stationary region}] = \text{Killing horizon}, \quad \xi_t^2|_{\mathcal{H}^\pm} = 0. \end{aligned}$$

In dynamical situations these notions differ, and the event horizon is the one with a global, teleological character, defined by the first two criteria. For non-static, but still stationary black holes, such as the Kerr rotating solutions, the third criterion is still valid but not the fourth one, which only characterizes the horizon of static black holes.

Although an imploding shell of photons provides a perfectly simple example of a collapse metric, it is hardly a very practical procedure to really engineer a black hole, much less a black hole of large proportions in the astrophysical sense. We could replace the photon shell by a shell of particles or a collapsing cloud of particles, which we may separate enough so that gravitation is the main force at play, say for instance a very large and dilute cloud of small asteroids. This

example is actually instructive in showing that black holes are inevitable, even without the need of invoking any kind of extreme conditions. Let the density of this cloud be ρ , as small as we like, and its size R , which gives a mass of order ρR^3 . Consider the Newtonian estimate for its Schwarzschild radius, $r_s \sim GM \sim G\rho R^3$. As we make the cloud larger and larger at fixed density, we can be sure that eventually $r_s(R) > R$, the crossover being the relativistic analog of the Jeans length $R_J \sim 1/\sqrt{G\rho}$. We thus conclude that the formation of sufficiently large black holes is inevitable given appropriate initial conditions.³ To put some numbers, a density of order $\rho \sim 10 \text{ kg/m}^3$, one hundred times smaller than that of water, we get $r_s \sim 3 \times 10^9 \text{ km}$ and $M \sim 10^9 M_\odot$, comparable to the size of the orbit of Neptune, and the mass is in the range of the largest supermassive black holes found at the centers of galaxies, such as M87* (the black hole at the center of our galaxy, Sgr A*, is lighter, with about $4 \times 10^6 M_\odot$).

Still, static and dilute spherical clouds that can collapse spherically are not easy to find out there as typical initial conditions. Smaller clouds tend to condense into stars that will have statistically distributed velocities, leading to virialized equilibria in which every particle moves in a statistical orbit in the field of all others. Collapse requires that the system sheds its angular momentum by radiative processes and this is not particularly fast for large clouds. A much better chance at setting up the conditions for gravitational collapse is to start from systems of larger density that are naturally spherical and are supported by an active agent that may one day falter. This is the case of stars.

8.2.2 Stars and Their Limits

Stars are self-gravitating balls of gas supported by pressure, either thermal pressure or the degeneracy pressure of a Fermi gas. Their energy as a function of their radius R is, in order of magnitude,

$$E(R) = E_{\text{gas}}(R) + E_G(R) = E_{\text{gas}}(R) - \frac{GM^2}{R} . \quad (8.30)$$

The equilibrium condition $\partial_R E(R) = 0$ gives

$$0 = \frac{GM^2}{R^2} + \frac{\partial E_{\text{gas}}}{\partial R} = \frac{GM^2}{R^2} + 4\pi R^2 \frac{\partial E_{\text{gas}}}{\partial V} = \frac{GM^2}{R^2} - 4\pi R^2 p_{\text{gas}} , \quad (8.31)$$

a version of the virial theorem. For thermal pressure, $p_{\text{gas}} = NT/V$, and we find $GM^2/R \sim NT$. Writing $M = Nm_p$ in terms of the proton mass we find

$$\frac{R_{\text{eq}}}{r_s} \sim \frac{m_p}{T} \sim \frac{\text{GeV}}{\text{keV}} \sim 10^6 , \quad (8.32)$$

where we have used the typical temperature of stellar cores, $T \sim 10^7 \text{ K} \sim \text{keV}$. Hot stars are very far from being black holes.

When the nuclear fuel is exhausted the star cools down and the pressure support must come from the degeneracy of fermions of mass m , such as electrons (in white dwarfs, $m = m_e$) or neutrons (in neutron stars, $m = m_n \approx m_p$). The number of fermions is of the order of the number of baryons $N = M/m_p$. They are packed at distances of the order of their de Broglie wavelength, $V \sim R^3 \sim N\lambda^3$, so that the Fermi momentum is $p_F \sim 1/\lambda \sim N^{1/3}/R$. The gas energy is then⁴ $E_{\text{gas}}(R) \sim N \cdot p_F^2/2m \sim N^{5/3}/mR^2$ in the non-relativistic case, and

³The fully relativistic treatment of this example in GR was given by Oppenheimer and Snyder in 1939.

⁴In this section we use natural units $\hbar = c = 1$.

$E_{\text{gas}} \sim N \cdot p_F \sim N^{4/3}/R$ in the relativistic case. The crossover between both behaviors takes place at $p_F \sim m$, i.e. at the radius $R_m \sim N^{1/3}/m$ (see Figure 8.10).




Figure 8.10: Left: energy of a degenerate Fermi gas as a function of the radius of the star. Center and right: total energy for masses below and above the critical mass.

At small radius, $R < R_m$, the total energy scales as $E(R) \sim (N^{4/3} - GM^2)/R$, and there are two regimes. For

$$M < M_c \sim \frac{m_{\text{Pl}}^3}{m_p^2}, \quad (8.33)$$

the degeneracy pressure wins at small R , and any putative stable minimum must lie in the large radius region $R_{\text{eq}} \geq R_m$ (Here $m_{\text{Pl}} = G^{-1/2} \sim 10^{19} \text{ GeV}$ is the Planck mass). On the other hand, for $M > M_c$ the total energy is negative and monotonically increasing, so that the system is unstable and collapses. Hence, M_c is the limiting mass for stability. Numerically, $m_{\text{Pl}}^3/m_p^2 \approx 1.8 M_\odot$, close to Chandrasekhar's precise value $1.4 M_\odot$ for white dwarfs.

On the stable branch, minimizing the energy in the non-relativistic regime,

$$E(R) \sim \frac{N^{5/3}}{m R^2} - \frac{GM^2}{R},$$

we obtain an equilibrium radius of order

$$R_{\text{eq}} \sim \frac{m_{\text{Pl}}^2}{m m_p^{5/3} M^{1/3}}, \quad (8.34)$$

a radius that decreases with increasing mass. The radius at the critical mass is $R_c = R_{\text{eq}}(M_c) \sim m_{\text{Pl}}/(m m_p)$, around 5000 km for white dwarfs and a few km for neutron stars. Comparing with the Schwarzschild radius,

$$\frac{R_{\text{eq}}}{r_s} \sim \frac{m_p}{m} \left(\frac{M_c}{M} \right)^{4/3}, \quad (8.35)$$

we see that neutron stars close to the critical mass are almost black holes, and GR must be used for a precise analysis (the Tolman–Oppenheimer–Volkoff equations).

The upshot is that M_c is of the order of a few solar masses. Even taking into account rotation and our uncertainties about the equation of state of nuclear matter, a very conservative bound for neutron stars is $M_c \lesssim 4 M_\odot$. Heavier compact remnants of stellar evolution have no option but to collapse into black holes.

Growth

Once formed, black holes grow by accretion. Infalling gas carries angular momentum and settles into a thin disk, where turbulent viscosity transports angular momentum outwards, so that matter spirals in through nearly circular orbits down to the ISCO and then plunges. The binding energy radiated on the way defines the efficiency $\eta = 1 - \varepsilon_{\text{ISCO}}$, with $\eta \simeq 0.06$ for Schwarzschild ($1 - \sqrt{8/9}$) and up to $\eta = 1 - 1/\sqrt{3} \simeq 0.42$ for a co-rotating disk around an extremal Kerr hole (see next section). The disk then shines with luminosity $L = \eta \dot{M}$, while the hole grows at the rate $\dot{M}_{\text{BH}} = (1 - \eta)\dot{M}$.

The luminosity is capped by radiation pressure: Thomson scattering of photons off electrons, which drag the protons along electrostatically, balances gravity at the Eddington luminosity. We can estimate its value as follows. Let the rate of energy radiated per unit area, at distance r , be $L/4\pi r^2$. In natural units ($c = 1$) this coincides with the flux of momentum per unit area because for photons, energy equals momentum. Since each electron presents an effective area given by the Thomson cross section $\sigma_T = (8\pi/3)\alpha^2/m_e^2$, the momentum transferred per unit time to an electron (and the proton that goes along for the ride) is $L\sigma_T/4\pi r^2$. Since this is an outward force, it equilibrates the gravitational force on the proton, GMm_p/r^2 , precisely at the Eddington luminosity

$$L_{\text{Edd}} = \frac{4\pi GMm_p}{\sigma_T} \simeq 1.3 \times 10^{31} \text{ W} \left(\frac{M}{M_\odot} \right). \quad (8.36)$$

Associating an effective black-body temperature to this luminosity, through the Stefan–Boltzmann form $L_{\text{Edd}} \sim 4\pi R_{\text{disk}}^2 \sigma_B T_{\text{eff}}^4$, one finds

$$T_{\text{eff}} \sim 1.5 \left(\frac{GM}{R_{\text{disk}}} \right)^{1/2} \left(\frac{M_\odot}{M} \right)^{1/4} \text{ keV},$$

explaining that accretion disks on compact objects such as neutron stars or solar-mass black holes radiate in X rays, whereas accretion disks on much larger black holes radiate on longer wavelengths.

Accretion at the Eddington rate yields exponential growth, $M(t) = M_0 e^{t/t_S}$, with the Salpeter time $t_S = \frac{\eta}{1-\eta} \frac{\sigma_T}{4\pi G m_p} \simeq 50 \text{ Myr}$ for $\eta = 0.1$. Quasars powered by black holes of $\sim 10^9 M_\odot$ are observed at $z \simeq 7.5$, when the universe was only $\sim 0.7 \text{ Gyr}$ old. Building them from stellar-mass seeds of $\sim 10^2 M_\odot$ requires $\log 10^7 \simeq 16$ e-folds, i.e. about 0.8 Gyr of uninterrupted Eddington-limited accretion, more than the available time. This tension suggests either heavy seeds of 10^4 – $10^5 M_\odot$ from the direct collapse of pristine gas clouds, or phases of super-Eddington accretion. The latter is possible because the Eddington bound assumes spherical symmetry, whereas in thick disks the photons can be advected into the hole. Recent detections by the James Webb Space Telescope of 10^7 – $10^8 M_\odot$ black holes at $z \gtrsim 10$ have sharpened the problem.

Problem: Charged Collapse

Consider a ball of radius R with electric charge Q and total energy M_∞ as measured from infinity, when the constituents are dispersed. At radius R the energy can be estimated as having contributions from rest mass, together with electrostatic and gravitational self-energies,

$$M(R) \approx M_\infty + \frac{Q^2}{4\pi R} - \frac{GM(R)^2}{R} . \quad (8.37)$$

- (a) Show that the charged ball will collapse only when $4\pi GM^2 > Q^2$. Even ignoring quantum effects, is this telling us anything about the possibility of viewing the electron as a tiny black hole?
- (b) The generalization of Schwarzschild's solution to a black hole carrying electromagnetic charge is known as the Reissner–Nordström solution, given by $ds^2 = -f(r)dt^2 + dr^2/f(r) + r^2d\Omega^2$ with

$$f(r) = 1 - \frac{2GM}{r} + \frac{GQ^2}{4\pi r^2} ,$$

with a radial electric field $E_r = Q/4\pi r^2$ (magnetic versions of these solutions do exist as well). Find the location of the horizon as a function of M and Q . Compute the surface gravity κ and see how it behaves in the ‘extremal’ limit $4\pi GM^2 \rightarrow Q^2$. Show that the proper distance to the horizon diverges in the extremal limit.

- (c) Show that surpassing the bound into $4\pi GM^2 < Q^2$ would produce a solution with a ‘naked’ singularity at $r = 0$ (a naked singularity is one which is causally connected with arbitrarily far away places).
- (d) Suppose you have a Schwarzschild black hole and you want to charge it by dropping protons from a long distance, starting at rest. Express the maximal value Q/M that you can get this way in terms of fundamental constants of nature. Can you make an extremal black hole by shooting particles with large inward velocity? Can you overcharge the black hole above the extremality limit?

8.3 Rotating Black Holes

In astrophysical situations where black holes are found, it is expected that they form from systems that carry non-zero angular momentum. More generally, we can always give angular momentum to a Schwarzschild black hole by throwing any object that is not falling on an exactly radial plunge. The asymptotic metric picks up spin-dependent terms of universal nature, as we have seen in (6.25) and this has characteristic ‘gravitomagnetic’ effects, also known as ‘frame-dragging’ effects, that are detectable on the orbital dynamics and the behavior of spinning tops. Given the astrophysical importance of rotating black holes, it is remarkable that a complete vacuum solution of Einstein's equations with the interpretation of a rotating black hole was found in closed form by Kerr in 1963.

8.3.1 The Kerr Metric

The Kerr solution is parametrized by the ADM mass M and the total angular momentum J and can be written in the so-called Boyer–Lindquist coordinates as follows ⁵

⁵A very similar solution known as the Kerr–Newman black hole gives the most general stationary, asymptotically flat black-hole solution of the Einstein–Maxwell equations, parametrized by mass, angular momentum and

$$ds^2 = -\frac{\rho^2 \Delta}{\Sigma} dt^2 + \frac{\Sigma}{\rho^2} \sin^2 \theta (d\phi - \omega dt)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 , \quad (8.38)$$

where $a = J/M$ is the angular momentum per unit mass. Further definitions include

$$\Delta = r^2 - 2GMr + a^2 , \quad \rho^2 = r^2 + a^2 \cos^2 \theta , \quad \Sigma = (r^2 + a^2)^2 - a^2 \Delta \sin^2 \theta , \quad (8.39)$$

and

$$\omega \equiv -\frac{g_{t\phi}}{g_{\phi\phi}} = \frac{2GMa r}{\Sigma} \quad (8.40)$$

gives a position-dependent ‘Coriolis field’ which represents the frame dragging or, equivalently, a local rotation of the coordinate frame.

The horizons are surfaces $r = \text{const}$ which are null, i.e. $\partial_\mu r$ is a null vector and, according to (8.29), $g^{rr} = \Delta/\rho^2 = 0$. Hence, the horizons sit at the roots of Δ ,

$$r_{\pm} = GM \pm \sqrt{(GM)^2 - a^2} , \quad (8.41)$$

with the extremality bound now being $(GM)^2 \geq a^2 = J^2/M^2$. Black holes spinning above this bound would have no smooth horizon and the singularity at $r = 0$ would be exposed in the sense that light rays starting from $r = 0$ would be able to reach arbitrarily far away, the causal structure being similar to that of negative-mass Schwarzschild as studied in Section 8.1.2.

Since $\Sigma(r_+) = (r_+^2 + a^2)^2$, the area of the event horizon is

$$A_{\mathcal{H}} = \int_{r=r_+} \sqrt{g_{\theta\theta} g_{\phi\phi}} d\theta d\phi = 4\pi (r_+^2 + a^2) . \quad (8.42)$$

Furthermore, evaluating the Coriolis field $\omega(r, \theta)$ at the horizon we find

$$\omega(r_+, \theta) = \Omega_{\mathcal{H}} = \frac{a}{r_+^2 + a^2} = \frac{a}{2GMr_+} , \quad (8.43)$$

what we may call the ‘angular velocity’ of the horizon.

The First Law

Directly manipulating the explicit expressions for the horizon location (8.41) and its area (8.42) we can derive the so-called Christodoulou–Ruffini relation

$$M^2 = \frac{A_{\mathcal{H}}}{16\pi G^2} + \frac{4\pi J^2}{A_{\mathcal{H}}} = M_{\text{irr}}^2 + \frac{J^2}{4G^2 M_{\text{irr}}^2} , \quad (8.44)$$

where the *irreducible mass* is defined by $M_{\text{irr}}^2 = A_{\mathcal{H}}/16\pi G^2$. This expression characterizes the part of the total energy, M , that is due to angular momentum, leaving an ‘irreducible’ component, M_{irr} , which would persist even if the hole would somehow lose its rotation. By explicit computation one further finds

$$\left. \frac{\partial M}{\partial A_{\mathcal{H}}} \right|_J = \frac{\kappa}{8\pi G} , \quad \left. \frac{\partial M}{\partial J} \right|_{A_{\mathcal{H}}} = \Omega_{\mathcal{H}} ,$$

charge. In these notes we will ignore the charge parameter, since one does not expect to find significantly charged black holes in nature. But charged black holes play a very important role as theoretical laboratories for various fundamental questions relating to black holes and quantum mechanics.

so that we can write

$$\delta M = \frac{\kappa}{8\pi G} \delta A_{\mathcal{H}} + \Omega_{\mathcal{H}} \delta J, \quad (8.45)$$

the *first law* of black hole dynamics, governing the energy balance of perturbations to the rotating black hole, distinguishing between the ‘kinetic’ part controlled by the angular momentum and the ‘irreducible’ part proportional to the area of the horizon.

The Kepler Problem

The Kepler problem around a Kerr black hole is altered by the frame dragging effect, allowing for the existence of highly bound orbits, provided they are co-rotating with the black hole, and conversely preventing binding for counter-rotating ones. An analysis quite similar to that of the Schwarzschild black hole gives, for equatorial orbits, an effective one-dimensional problem

$$\frac{1}{2} \dot{r}^2 + V_{\text{eff}}(r) = \varepsilon_{\text{eff}}, \quad (8.46)$$

where $\varepsilon_{\text{eff}} = \frac{1}{2}(\varepsilon^2 - 1)$, with $\varepsilon = E/m$ the conserved orbital energy per unit rest mass and

$$V_{\text{eff}}(r) = -\frac{GM}{r} + \frac{\ell^2 - a^2(\varepsilon^2 - 1)}{2r^2} - \frac{GM(\ell - a\varepsilon)^2}{r^3}, \quad (8.47)$$

and $\ell = L/m$ the orbital angular momentum per unit rest mass.

The most important new qualitative fact about the orbital dynamics around a Kerr black hole is that the ‘innermost stable circular orbits’ (ISCO) can exist much deeper into the near-horizon region than they do in the Schwarzschild case. Circular orbits satisfy $V_{\text{eff}} = \varepsilon_{\text{eff}}$ and $V'_{\text{eff}} = 0$, where ε also appears inside the potential. In the extremal case, $a = GM$, these conditions can be solved in closed form. Writing $r = GMx^2$, the co-rotating circular orbits have

$$\varepsilon = \frac{x^2 + x - 1}{x^{3/2}\sqrt{x+2}}, \quad \ell = GM \frac{x^3 + x^2 + x - 1}{x^{3/2}\sqrt{x+2}}, \quad (8.48)$$

and one checks that $V''_{\text{eff}} > 0$ for all $x > 1$. Co-rotating circular orbits are therefore stable all the way down to $r = r_+ = GM$, which is the ISCO. (The ISCO, the photon orbit and the horizon all sit at $r = GM$ in Boyer–Lindquist coordinates, but they remain distinct: the radial coordinate degenerates in the extremal throat.) As $x \rightarrow 1$ we get $\varepsilon \rightarrow 1/\sqrt{3}$. Gas spiraling slowly from infinity through circular orbits down to the ISCO thus radiates the fraction

$$\eta = 1 - \varepsilon_{\text{ISCO}} = 1 - \frac{1}{\sqrt{3}} \simeq 0.42 \quad (8.49)$$

of its rest mass, compared with $1 - \sqrt{8/9} \simeq 0.06$ for Schwarzschild. The same orbit has $\ell = 2GM\varepsilon$. Accretion from the ISCO thus keeps an extremal hole extremal, since $\delta(GM^2 - J) = m(2GM\varepsilon - \ell) = 0$. Counter-rotating orbits are instead pushed out to $r_{\text{ISCO}} = 9GM$, with $\varepsilon_{\text{ISCO}} = \sqrt{25/27}$ and $\eta \simeq 0.04$.⁶

⁶In practice the spin of an accreting hole saturates slightly below extremality, at $a \simeq 0.998 GM$, because the hole preferentially captures disk photons with negative angular momentum (Thorne, 1974). This lowers the maximal efficiency to $\eta \simeq 0.3$.

Problem: Where Is the Extremal ISCO?

According to (8.48), the ISCO of an extremal Kerr hole sits at the Boyer–Lindquist radius $r = GM = r_+$, seemingly *on* the horizon. To clarify this, consider a near-extremal hole with $a = GM(1 - \delta)$, $\delta \ll 1$.

- Using the condition for marginally stable prograde circular orbits, $r^2 - 6GMr + 8a\sqrt{GMr} - 3a^2 = 0$, show that $r_{\text{ISCO}} \simeq GM [1 + (4\delta)^{1/3}]$, whereas $r_+ \simeq GM (1 + \sqrt{2\delta})$.
- Compute the proper circumference of the ISCO and of the equator of the horizon. Show that the latter equals $4\pi GM$ for *any* spin, and that the former tends to the same value as $\delta \rightarrow 0$. Is this a good test of whether the ISCO falls onto the horizon?
- Compute the proper radial distance between the horizon and the ISCO, along the equatorial plane of a $t = \text{const}$ slice, and show that it diverges as $(GM/6) \log(1/\delta)$. Compare with the photon circular orbit, $r_{\text{ph}} \simeq GM (1 + 2\sqrt{2\delta/3})$, and interpret the result in terms of the intrinsic geometry of the equatorial plane near $r = GM$.

Problem: Maximal Time Dilation on Kerr Orbits

An astronaut orbits a Kerr black hole of mass M and spin a on a circular equatorial geodesic of radius r . Her clock runs slow relative to static clocks at infinity by the factor $\gamma_\infty \equiv dt/d\tau = u^t$.

- From the radial component of the geodesic equation and the normalization $u^2 = -1$, show that the angular velocity of circular orbits is $\Omega = d\phi/dt = \pm\sqrt{GM}/(r^{3/2} \pm a\sqrt{GM})$, with the upper (lower) sign for co-rotating (counter-rotating) orbits, and that

$$\gamma_\infty(r) = \frac{r^{3/2} \pm a\sqrt{GM}}{r^{3/4} \sqrt{r^{3/2} - 3GM} \pm 2a\sqrt{GM}}.$$

- What is the largest time dilation factor attainable on circular geodesics? Show that $\gamma_\infty(r)$ decreases monotonically with r for all circular orbits outside the horizon.
- Restrict now to *stable* circular orbits. Is the ISCO the stable orbit with maximal time dilation? Using the marginal stability condition $r^2 - 6GMr \pm 8a\sqrt{GMr} - 3a^2 = 0$, express γ_∞ at the ISCO as a function of r_{ISCO} alone, and study its dependence on a . How close to extremality must the hole spin to make one hour in orbit last seven years at infinity?

8.3.2 The Ergosphere

The Kerr metric has two Killing vectors, $\xi_t = \partial_t$ and $\xi_\phi = \partial_\phi$. The first one is timelike at infinity, $\xi_t^2 = g_{tt} \rightarrow -1$, but it becomes spacelike already *outside* the event horizon. The condition $\xi_t^2 = g_{tt} = 0$ reads $r^2 - 2GMr + a^2 \cos^2 \theta = 0$, which defines the surface

$$r_{\mathcal{E}}(\theta) = GM + \sqrt{(GM)^2 - a^2 \cos^2 \theta}, \quad (8.50)$$

touching the horizon at the poles, $\theta = 0, \pi$. The region $r_+ < r < r_{\mathcal{E}}(\theta)$, where $\xi_t^2 > 0$, is called the *ergosphere*, $\mathcal{E}$ (Figure 8.11). In the ergosphere there are no static observers: nothing can stay at rest with respect to the asymptotic frame, no matter how powerful its engines are. This is an extreme form of the *dragging of inertial frames* by the rotating mass, a phenomenon that realizes some of the ideas of Mach's principle.




Figure 8.11: *Left: the ergosphere $\mathcal{E}$ of a rotating black hole, in a section containing the rotation axis. Right: stationary observers must rotate with angular velocities in the interval $\Omega_- < \Omega < \Omega_+$.*

There are, however, stationary observers, whose four-velocity is proportional to a Killing vector,

$$u^\mu = \frac{dx^\mu}{d\tau} = \alpha \left(\xi_t^\mu + \Omega \xi_\phi^\mu \right) ,$$

with α fixed by $u^2 = -1$. Such an observer has $dt/d\tau = \alpha$ and $d\phi/d\tau = \alpha\Omega$, so that $d\phi/dt = \Omega$, i.e. it rotates rigidly with angular velocity Ω as seen from infinity. The four-velocity must be timelike, $(\xi_t + \Omega \xi_\phi)^2 = \xi_t^2 + 2\Omega \xi_t \cdot \xi_\phi + \Omega^2 \xi_\phi^2 < 0$, which requires $\Omega_- < \Omega < \Omega_+$ with

$$\Omega_{\pm} = \omega \pm \frac{\sqrt{\Delta} \rho^2}{\Sigma \sin \theta} . \quad (8.51)$$

In the ergosphere $\Omega_- > 0$: all stationary observers co-rotate with the hole. As $r \rightarrow r_+$, $\Delta \rightarrow 0$ and the interval shrinks to a single value, $\Omega_{\pm} \rightarrow \Omega_{\mathcal{H}}$, with

$$\Omega_{\mathcal{H}} = \omega(r_+) = \frac{a}{r_+^2 + a^2} , \quad (8.52)$$

showing that $\Omega_{\mathcal{H}}$ represents physically the angular velocity of the horizon. Accordingly, the Killing vector

$$\xi_{\mathcal{H}} = \xi_t + \Omega_{\mathcal{H}} \xi_\phi \quad (8.53)$$

is null on $\mathcal{H}^+$ and timelike in the whole exterior of the black hole. It may be used to give a definition of surface gravity using the general relations (8.20), in terms of this timelike Killing vector that is available in the exterior of the black hole, namely we can compute κ as ⁷

$$\nabla_{\xi_{\mathcal{H}}} \xi_{\mathcal{H}} = \kappa \xi_{\mathcal{H}} , \quad \kappa = \frac{r_+ - GM}{r_+^2 + a^2} . \quad (8.54)$$

⁷By passing to a non-rotating frame near the horizon, $\phi \rightarrow \phi - \Omega_{\mathcal{H}} t$ one can check that κ admits the usual definition in terms of derivatives of the metric components, and furthermore it has the same interpretation of an inverse-relaxation time for near-horizon processes.

Interestingly, $\kappa \rightarrow 0$ in the extremal limit.

8.3.3 The Penrose Process and the Second Law

The energy of a particle with four-momentum p^μ , as measured at infinity, is $E = -p_\mu \xi_t^\mu$. Inside the ergosphere, ξ_t is spacelike, and E can be negative for physical, future-directed, timelike or null momenta. Consider a particle with momentum p_0 entering the ergosphere, where it splits into two, $p_0 = p_1 + p_2$, with $E_0 = E_1 + E_2$. If $E_1 < 0$, the second particle escapes to infinity with $E_2 > E_0$. Hence, energy has been extracted from the black hole (Figure 8.12).




Figure 8.12: *The Penrose process, seen in the equatorial plane. A particle splits inside the ergosphere, one fragment falling into the horizon with negative energy, the other escaping to infinity with more energy than the original particle.*

The negative energy particle must fall into the hole, since it cannot exist outside the ergosphere. It changes the mass and angular momentum of the hole by $\delta M = E_1 < 0$ and $\delta J = L_1 = p_1 \cdot \xi_\phi$. Since p_1 is a future-directed causal vector crossing $\mathcal{H}^+$, and $\xi_{\mathcal{H}}$ is future-directed and null on $\mathcal{H}^+$, we must have $p_1 \cdot \xi_{\mathcal{H}} \leq 0$. Hence,

$$0 \geq p_1 \cdot (\xi_t + \Omega_{\mathcal{H}} \xi_\phi) = -E_1 + \Omega_{\mathcal{H}} L_1 = -\delta M + \Omega_{\mathcal{H}} \delta J ,$$

which, using the first law (8.45) implies

$$\delta A_{\mathcal{H}} = \frac{8\pi G}{\kappa} (\delta M - \Omega_{\mathcal{H}} \delta J) \geq 0 . \quad (8.55)$$

This is a remarkable result. It shows that we may extract energy from a rotating black hole, $\delta M < 0$, but only by slowing down its rotation, $\delta J < 0$ and the process always ends up with a bigger black hole $\delta A_{\mathcal{H}} > 0$. The maximal extractable energy is $M - M_{\text{irr}}$, which is about 29% of the total mass for an extremal Kerr black hole. This is potentially a gigantic amount of energy and, in fact, an electromagnetic version of the Penrose process, known as the Blandford–Znajek mechanism, is conjectured to power relativistic jets in black holes with highly magnetized accretion disks, such as those found at the center of the M87 galaxy or distant quasars.

The statement that $\delta A_{\mathcal{H}} \geq 0$ is a general theorem, proved by Hawking, which is actually valid in fully dynamical situations, only assuming the null energy condition and ‘cosmic censorship’

(the absence of naked singularities), and it is known as the *second law* of black-hole dynamics. The second law is arguably the most important result in classical black hole theory and it uses many of the same arguments that are responsible for the singularity theorems of Penrose and Hawking.

Problem: Cosmic Censorship

Consider small perturbations δM and δJ of an extremal Kerr black hole. Show that the first and second laws prevent the possibility that such perturbations could be used to over-spin the black hole into a solution that violated the bound $GM^2 \geq J$, which formally would have a naked singularity.

A simple model for such a perturbation is the attempt at over-spinning an extremal Kerr of mass M , by sending in a particle of rest mass m , energy εm at infinity and angular momentum $m\ell$. Show that one needs $\ell > 2GM\varepsilon$ in order to have a chance at achieving the over-spinning. On the other hand, show that the maximum of the effective potential in the marginal case $\ell = 2GM\varepsilon$ just manages to prevent absorption of such a particle.

The second law also puts constraints on the amount of energy that can be radiated in gravitational waves in a black-hole merger episode. Let two black holes with parameters (M_1, J_1) and (M_2, J_2) merge into a third black hole of parameters (M_3, J_3) . The parameters are constrained by the second law: $A_3 \geq A_1 + A_2$. The largest amount of energy in gravitational waves is obtained by starting with a configuration with the largest available energy for extraction (maximizing rotational energy of the initial black holes makes them extremal). Similarly, a final Schwarzschild hole ensures that no more energy is left for extraction. This implies that we must have $M_{\text{irr}} = M/\sqrt{2}$ for the initial black holes and $M_{\text{irr}} = M$ for the final one, so that the efficiency η is bounded by

$$\eta = \frac{M_1 + M_2 - M_3}{M_1 + M_2} = 1 - \frac{M_3}{M_1 + M_2} \leq 1 - \frac{\sqrt{M_1^2 + M_2^2}}{\sqrt{2}(M_1 + M_2)} \leq \frac{1}{2},$$

where the last bound is saturated for equal initial masses. A similar argument for Schwarzschild black holes yields a maximum efficiency of $1 - 1/\sqrt{2} \approx 29\%$. So, we find that non-rotating black holes are not completely inert as energy sources after all! At any rate, the efficiencies found in actual mergers are far from these maximal bounds, staying well below 10%.

Another version of the Penrose process is its wave incarnation, known as *superradiance*. Suppose that a classical wave of frequency ω and angular momentum L (measured in the same direction as the spin of the black hole) impinges on a Kerr black hole of mass M and angular momentum J . If we view the wave as a coherent state of quantum bosonic particles, their scattering will produce variations δM and δJ which will add up coherently for the ensemble of particles. Therefore, we will have approximately $\delta M/\delta J \sim \omega/L$. Applying then the first law we will have

$$\frac{\kappa}{8\pi G} \delta A_{\mathcal{H}} = \delta M - \Omega_{\mathcal{H}} \delta J = \delta M \left(1 - \frac{L}{\omega} \Omega_{\mathcal{H}} \right).$$

Now, if the second law is valid, this quantity is non-negative, which implies that, for any $\omega < L\Omega_{\mathcal{H}}$ we have $\delta M < 0$, implying that the outgoing wave is amplified and we extract energy from the hole. We can interpret superradiance as a form of ‘stimulated emission’, in an analogy with

atomic physics. The impinging wave activates the ergosphere, which causes the black hole to de-excite into a lower energy state. If this analogy is correct, we should also expect ‘spontaneous emission’ by quantum effects. This is precisely what Hawking radiation is.

8.4 Bekenstein and Hawking

The first and second laws of black hole dynamics evoke the corresponding laws of thermodynamics, and this is no coincidence. The first law ⁸

$$\delta M = \frac{\kappa}{8\pi G} \delta A_{\mathcal{H}} + \Omega_{\mathcal{H}} \delta J \quad (8.56)$$

suggests the identification of M with the internal energy of a thermodynamic system, the right hand side representing the distinction between ‘work’ (the rotational energy that we may affect from outside) and ‘heat’, an irreducible component proportional to the area term. Furthermore, perturbations relax exponentially, in a time scale of order κ^{-1} , a dissipative behavior characteristic of thermodynamical systems with κ replaced by the temperature. The second law $\delta A_{\mathcal{H}} \geq 0$ is also in line with the second law of thermodynamics if we make the associations

$$\frac{A_{\mathcal{H}}}{G} \longrightarrow S, \quad \kappa \longrightarrow T. \quad (8.57)$$

Further evidence comes from the fact that κ is constant over a horizon, emulating the *zeroth law*, namely that temperature is constant throughout a region in thermal equilibrium. Even the fact that zero absolute temperature is not reachable in finite-time processes (*third law*) can be interpreted as the ‘cosmic censorship’ phenomenon (the dynamical resistance of black holes to admit processes that would push them beyond extremality). Together, these statements constitute the ‘thermodynamic analogy’. The question is whether there is anything more than a mere analogy.

8.4.1 Black Hole Entropy

In the early 70’s, Bekenstein gave arguments in favor of these identifications, using the information-theoretical interpretation of entropy, as a measure of ‘hidden information’, and the arguments used quantum theory in a crucial way. The basic idea is to consider the absorption of a single quantum degree of freedom (a particle) by a black hole.

For simplicity we consider here the case of neutral particles on a Schwarzschild black hole. If the energy of this particle is ω , its degree of localization in space can be no better than $\hbar/\omega$, according to Heisenberg’s relation. Suppose that we only know that this particle has been absorbed by the black hole, so that the hidden information has increased by one bit, $\delta S = \log 2$. It follows that the energy of this particle must have been $\omega \sim \hbar/r_s$. The reason is that a particle of larger energy is localized on distances smaller than r_s and, when captured, we can know for example whether it entered through the north or south pole, or perhaps a few degrees southwest, etc. Since we are only interested in minimal hidden information, we must arrange for the particle to be delocalized on the scale of the horizon, so we cannot know ‘where precisely’ it entered. On

⁸Adding charge, thus considering the Kerr–Newman solutions, one simply adds a term $\Phi_{\mathcal{H}} \delta Q$ to the right hand side, where $\Phi_{\mathcal{H}}$ is the electrostatic potential at the horizon.

the other hand, the energy cannot be smaller than $\hbar/r_s$, because in that case the wavelength of the particle is larger than the hole itself, and the probability of capture becomes very small.

Hence, we conclude that single-bit entropy increments are associated to a single-quantum capture of energy $\omega \sim \hbar/r_s \sim \hbar/GM$. The corresponding increment in mass is $\delta M \sim \omega$ and the increment in area:

$$\delta A \sim \delta(G^2 M^2) \sim G^2 M \delta M \sim G r_s \omega = O(G\hbar) .$$

In this way we establish $\delta S \sim \delta A/G\hbar$, and therefore $S \sim A/G\hbar$ when ‘building’ the black hole from such optimal captures. In other words, we learn that the area increases by one Planck unit when the entropy increases by one bit. In particular, the black hole has ‘infinite entropy’ in the classical limit ($\hbar \rightarrow 0$). This agrees with the idea that, according to the no-hair theorem, all detailed information about the formation of the hole is hidden from the classical description, which only depends on three parameters: M, Q, J .

Since $A \sim (GM)^2$ for a Schwarzschild black hole, a relation of the form $S \sim A/G\hbar$ automatically *implies* that the black hole has a natural temperature obtained from the classic definition

$$T = \left(\frac{\partial S}{\partial E} \right)^{-1} ,$$

which gives $T \sim \hbar/GM \sim \hbar\kappa$ after the natural identification $E = Mc^2$.

8.4.2 Hawking Radiation

The Penrose process has a quantum analog. Pairs of particles can be spontaneously created from the vacuum with total energy $E_+ + E_- = 0$, provided one of them has negative energy, $E_- < 0$. Since $E = -p_\mu \xi_t^\mu$ must be positive wherever ξ_t is timelike, the negative energy partner must end up in a region where ξ_t is spacelike, i.e. inside the horizon or inside the ergosphere. The positive energy partner may escape to infinity as radiation of frequency $\omega > 0$, while the hole loses mass $\delta M = -\omega$. For the Kerr black hole, the pair creation can occur outside $\mathcal{H}^+$, inside the ergosphere, in which case the negative energy particle is captured by the hole in an essentially classical process that respects the second law $\delta A_{\mathcal{H}} \geq 0$. However, if the pair creation occurs right at $\mathcal{H}^+$, one of the particles can be nucleated inside the horizon, in which case the loss of mass by the hole is intrinsically quantum mechanical and can violate the second law, in particular, it can work for an ordinary Schwarzschild black hole.

Let the nucleated particle-antiparticle pair have rest mass m , so that the pair begins ‘real’ life separated by a typical proper distance of order $\hbar/m$. This corresponds to a radial height r_m satisfying

$$\frac{\hbar}{m} \sim \int_{r_s}^{r_m} \frac{dr}{\sqrt{1 - r_s/r}} ,$$

leading to $r_m - r_s \sim \hbar^2 \kappa / m^2$. The asymptotic energy of such a particle at maximal radial height r_m is of order

$$E = -m u_\mu \xi_t^\mu = m \sqrt{1 - r_s/r_m} \sim m \sqrt{\kappa(r_m - r_s)} \sim \hbar \kappa ,$$

where we made no attempt at fixing $O(1)$ factors. Notice that the asymptotic energy is independent of the particle rest mass. From the point of view of an asymptotic observer, the near horizon region is populated by particles of typical asymptotic energy $\hbar\kappa$ which are confined by the gravitational potential. We can expect to have a thermal gas at effective temperature $T_H \sim \hbar\kappa$ which slowly leaks out by tunneling through the gravitational potential barrier. We

thus expect a thermal emission spectrum at temperature T_H , modulated by a grey-body factor from the tunneling.

A famous calculation by Hawking in 1974, using quantum field theory, gives

$$T_H = \frac{\hbar\kappa}{2\pi} = \frac{\hbar}{8\pi GM} , \quad (8.58)$$

where the second expression assumes a Schwarzschild black hole. The expression in terms of the surface gravity is universal for any Kerr–Newman black hole. The luminosity in Hawking radiation is then given by

$$L_H = \frac{dE_{\text{rad}}}{dt} = \int \omega d\omega \frac{d\langle n_\omega \rangle}{dt} = \int \frac{d\omega \omega |T(\omega)|^2}{e^{\omega/T_H} - (-1)^F} . \quad (8.59)$$

Here $F = 0, 1$ distinguishes Bose and Fermi statistics. For a Kerr black hole, the Boltzmann suppression $\exp(-\omega/T_H)$ is replaced with $\exp(-(\omega - L\Omega_H)/T_H)$, from the fact that particles with angular momentum L are emitted with an extra energy $L\Omega_H$ from the frame dragging effect, so the purely thermal excitation energy is smaller by this amount. Incidentally, we see that bosons will undergo Bose-condensation in the vicinity of the horizon for $\omega < L\Omega_H$, another avatar of the superradiance phenomenon.

The precise determination of the normalization factor in (8.58) fixes the absolute normalization of the entropy to be

$$S_{BH} = \frac{A_H}{4\hbar G} , \quad (8.60)$$

a formula known as the Bekenstein–Hawking entropy formula. A fundamental problem in theoretical physics is to account for this thermodynamic entropy by counting quantum microstates. Such an entropy is equivalent to one bit of information per Planck area element of the horizon, so that one needs a full theory of quantum gravity to account for it. Furthermore, the dependence on an area rather than a volume is quite intriguing, the starting point of the so-called ‘holographic’ approach to quantum gravity.

There is little doubt about the importance of (8.58) and (8.60) for the foundations of physics. However, the practical impact on astrophysical black holes is less clear. The actual value of the Hawking temperature for astrophysical black holes is $T_H \sim 10^{-7}(M_\odot/M)$ Kelvin, well below the CMB temperature, so that astrophysical black holes absorb much more energy than they emit. In natural units with $\hbar = c = 1$, we can easily estimate the luminosity in Hawking radiation as $L_H \sim T_H^2 \sim A_H T_H^4$, the Stefan–Boltzmann law, even though the emitter has a size comparable to the typical wavelength, $\lambda \sim r_s \sim GM$. The lifetime of the hole is then

$$\tau_{BH} \sim \frac{M}{L_H} \sim r_s S_{BH} \sim \frac{S_{BH}}{T_H} , \quad (8.61)$$

i.e. $\tau_{BH} \sim G^2 M^3$, a gigantic time scale for any solar mass black hole, vastly larger than the age of the universe. It is said that the black hole *evaporates*, emitting one quantum of energy T_H every time interval of order T_H^{-1} , and a total number of $O(S_{BH})$ quanta over its lifetime.

The entropy of a black hole resulting from the collapse of a typical star made of N nucleons is $S_{BH} \sim Gm_p^2 N^2 \sim 10^{-38} N^2$, to be compared to the gas entropy for that star $S_{\text{gas}} \sim N$. Since $N \sim 10^{57}$ we see that gravitational collapse is an *extremely* irreversible process. The equilibrium between normal and collapsed entropies occurs for a mass of order $M_* \sim 10^{38} m_p \sim 10^{15}$ grams,

the mass of a typical mountain. Such black holes have size of the order of one fermi and temperatures of tens of MeV (of order m_p if the factor 8π in (8.58) is ignored). If they existed as ‘primordial’ black holes, somehow created at or near the Big Bang, they should be completing evaporation about *now*.

The greybody factors $|T(\omega)|^2$ suppress the emission of particles with large spin, and particles with mass $m > T_H$ are Boltzmann suppressed. Hence, large black holes radiate mostly neutrinos and photons (and gravitons). This implies that baryon number is violated. Consider a large body with baryon number $N_B \gg 1$ collapsing into a black hole. The hole radiates without returning the baryons until $T_H \sim m_p$, i.e. until its mass is down to $M_{\text{BH}} \sim m_{\text{Pl}}^2/m_p \sim 10^{38} m_p \sim M_*$. It is then impossible to return a baryon number $N_B \gg N_B(M_*) \sim 10^{38}$: black hole evaporation implies a macroscopic violation of global symmetries. To the extent that black holes at the Planck scale can be present in quantum gravitational fluctuations, we do not expect any theory of quantum gravity to have exact global symmetries.

Problem: Yellow Black Hole

Estimate the radiation power of a yellow black hole. Compare its entropy to that of the Sun.
